

*Climate of the Past Discussions* is the access reviewed discussion forum of *Climate of the Past*

# A studying of solar-ENSO correlation with southern Brazil tree-ring index (1955–1994)

N. R. Rigozo<sup>1,2,3</sup>, D. J. R. Nordemann<sup>3</sup>, M. Pereira de Souza Echer<sup>1</sup>, E. Echer<sup>3</sup>,  
and A. Prestes<sup>3</sup>

<sup>1</sup>Faculdade de Tecnologia Thereza Porto Marques – FAETEC, CEP 12308-320, Jacareí, Brazil

<sup>2</sup>LARAMG – Laboratório de Radioecologia e Mudanças Globais/Departamento de Biofísica e Biometria da Universidade do Estado do Rio de Janeiro (UERJ), Rio de Janeiro - RJ, Brazil

<sup>3</sup>Instituto Nacional de Pesquisas Espaciais – INPE, CP 515, 12201-970 São José dos Campos, Brazil

Received: 24 August 2005 – Accepted: 3 October 2005 – Published: 7 November 2005

Correspondence to: N. R. Rigozo (rodolfo@dge.inpe.br)

© 2005 Author(s). This work is licensed under a Creative Commons License.

215

## Abstract

Solar activity, volcanic aerosol, El Niño-Southern Oscillation and global temperature anomalies effects on Southern Brazil tree growth rings are presented through multiple linear analysis. Linear correlations were made on annual, 10 year running averages and band pass filter. For annual averages, the correlation coefficients were low, and the 5 10 years running average correlations the coefficient correlations were much higher. The multiple regression of 2 to 5 year band pass filter indicates that 60% of the variance in tree ring index was explained by volcanic eruptions, Southern Oscillation Index and temperature anomalies. The multiple regression of 10 year running averages indicates 10 that 84% of the variance in tree ring index was explained by solar activity and another time series. These results indicate that the effects of solar activity, volcanic eruptions, ENSO and temperature anomalies on tree rings are better seen on long timescales than volcanic eruption, ENSO and temperature anomaly.

## 1 Introduction

15 Around 76% of the surface of the Southern Hemisphere is covered by water. The interaction between ocean and atmosphere and specially the mechanisms of heat transport determine the patterns of present and past climates (Hamon and Godfrey, 1978). Due to this disparity in land area, the Southern Hemisphere climate is predominantly maritime than continental.

20 Much has been written about Northern Hemisphere climatic variations, but comparatively little is available on the Southern Hemisphere. A complete understanding of the climatic behavior of one hemisphere, however, is not possible without the knowledge of what happens in the other one (Pittock, 1978). A true understanding of climate change requires field experiments, empirical studies, and theoretical models. Theoretical models provide a guide for empirical studies. The models may predict previously 25 unsuspected relationships that experiments and empirical studies can either confirm

216

or disprove. Advancement requires the interaction between theory, experiment, and empirical models (Hoyt and Schatten, 1997).

The instrumental meteorological records available in the Southern Hemisphere are shorter, and more spatially concentrated than in the Northern Hemisphere. Very few Southern Hemisphere temperature records extended back into the nineteenth century (Jones et al., 1996). Working with such short timescales, trends lasting decades to centuries are difficult to evaluate with statistical confidence. One way to study climatic variations and solar variations before the previous century is to utilize indirect sources or “proxy data” such as tree-rings. Tree ring vary in width each year depending on variations in local temperature and local precipitation. Some tree-ring growths are mostly controlled by temperature and others by precipitation, but generally the growth control a mixture of the two (Fritts, 1976).

In this work we make a comparison between modeled and observed tree-ring index by accounting for additional natural forcings. That is important to evaluate the use of the tree-rings data as proxy data for climate and solar effects in Southern America.

## 2 Methodology of data analysis

Our study makes use of five observational data sets: Tree-ring index (Fig. 1a), Zürich Sunspot Number (Rz) (Fig. 1b), global temperature anomalies (Fig. 1c), Southern Oscillation Index (SOI) (Fig. 1d) with representative of the El Niño and La Niña events, and stratospheric aerosol optical depth measurements (Fig. 1e).

The trees used in this study were sampled from four locations in Rio Grande do Sul State, in Southern Brazil: nine samples from Canela (29°18'S, 50°51' W, 790 m a.s.l.), one sample from Nova Petrópolis (29°20'S, 51°10' W, 579 m a.s.l.), nine samples from São Francisco de Paula (29°25' S, 50o 24'W, 930 m a.s.l.) and two samples from São Martinho da Serra (29°30' S, 53°53' W, 484 m a.s.l.) (Rigozo, 1998). A total of 21 samples were used in this study. Trees were native *Araucaria* trees (*Araucaria angustifolia*) of ages ranging from 30 to 39 years. Trees of this species with longer lifetime exist but

217

because of the enforcement of laws on native species, it was not possible to sample older trees at the time of sampling. The methodology of polishing and treatment of the samples, as well as the digital image analysis of the scanned samples are described in Rigozo and Nordemann (1999) and Rigozo et al. (2001).

Every time series was detrended using an exponential function to eliminate growth trends and periods longer than 25 years. A growth index for each radius time series and then means by location and a global index for the region concerned were obtained.

Because long term climatic data for the collected samples are not available, our hypothesis about the climate response in the tree-ring data for this region is based on the global temperature anomalies. The annual global temperature anomalies come from about 4000 meteorological stations around the world. These data are an update of the analysis described by Hansen and Lebedeff (1987, 1988). These data are available in Goddard Institute for Space Studies (GISS).

The SOI is calculated from the monthly or seasonal fluctuations in the air pressure difference between Tahiti (20° S, 150° W) and Darwin (Australia, 10° S, 130° E). Positive values of SOI indicate La Niña events and negative values of SOI indicate El Niño events (Enfield, 1989; Neelin and Latif, 1997).

Sunspot data where obtain from The Sunspot Index Data Center – Brussels (World Data Center for the Sunspot Index). Sunspot number is the longest solar activity index available and it is a representative of the general state of solar activity (Hoyt and Schatten, 1997).

Volcanic eruption effect on the climate is represented by the stratospheric aerosol optical depth (SAOD) data set of Sato et al. (1993 and updates). The optical depth of the aerosol layer is considered to be the primary stratospheric aerosol climate forcing factor (Lacis et al., 1992; Sato et al., 1993). Major volcanic eruptions inject sulfur dioxide into the stratosphere, where, within a few weeks, it is converted to sulfate aerosol. This sulfate aerosol affects incoming solar radiation, reducing the amount reaching the earth's surface and thereby leading to a temperature reduction at the surface and throughout the lower troposphere (Mass and Portman, 1989; Dutton and Christy, 1992). The

218

SAOD data set used in our analysis contains yearly average south hemisphere SAOD values covering the period from 1955 through 1994 (Sato et al., 1993 and updates).

## 2.1 Analysis and results

In order to determine the effect of solar activity, temperature, ENSO and SAOD on tree-ring index, a multivariate linear regression approach is used. Initially spectral analysis in tree ring index time series is performed to determine the frequencies contained in the data. The spectral analysis results in tree ring time series are described. Figure 2 shows the amplitude as a function of frequency as determined by iterative regression analysis. It may be observed in the spectrum that there is no domination of long periods (>10 years) over the short ones (<10 years), or, in other terms, the long periods do not hide the low periods in the spectrum. The nature of this distribution suggests a favorable response on tree growth rate in South Brazil, to environmental factors for short and long periods. The tree-ring index from 1955 through 1994 is fit using the 10 years running average and 2–5 year band pass filter. These 10 year running averages and band pass filtered data were also applied to *Rz*, temperature anomaly, SOI and SAOD. The 8.2 yr periodicity was not included in the analyses because it is not caused by any natural phenomenon but it is caused man, because to every 8 years the reservations of tree from San Francisco of Paula and Canela in Brazil suffer cut of the forest to eliminate the sick trees in favor of the healthy ones (Rigozo, 1998).

The regression results are presented in Table 1 to original series, Table 2 for the 2 to 5 years band pass filter and Table 3 for the ten-year running averages. We found that the *Rz* in combination with temperature anomaly, ENSO and with SAOD original series explained only 17,8% of the variation in the tree ring index data, but using the band pass filter of 2 to 5 years on that data the combined influence of those phenomenon's for low periods on the tree ring index increases for 60,3% (Table 1). The influence of those phenomena in the variation of the tree ring index for longer periods than 10 years is very large. Using the 10 years running average on the data 84.2% of the variation in the tree ring index data is explained (Table 2).

219

The cross-correlation coefficients between tree ring index and the time series, for the average 10 years running average were: 0.70 to tree-ring index  $\times$  *Rz* with a lag of 1 year, 0.65 tree-ring index temperature anomaly with a lag of 1 year, -0.82 tree-ring index  $\times$  SOI index with a lag of 2 years and -0.77 tree ring index  $\times$  SAOD with a lag of -4 years. The cross-correlation tree ring index and the time series for the band pass filter of 2 to 5 years the were: 0.74 tree-ring index temperature anomaly with a lag of zero year, -0.45 tree-ring index  $\times$  SOI index with a lag of -3 years and 0.47 tree ring index  $\times$  SAOD with a lag of -2 years.

## 3 Discussion

Rigozo et al. (2004), in studies on solar activity in tree ring data from Concórdia-Brazil (27° S, 51° W), have found a lag of zero year for the 11 years period, between sunspot number and tree ring width, which indicates that in the Concórdia region the growth of tree seems to follow immediately the sunspot activity variability. Rigozo et al. (2002), still studying the solar variability effects in the tree-ring data, from Concórdia-Brazil, found in the cross-wavelet spectral analysis between the tree ring data and sunspot number a high amplitude for the period close to 11 years, during the time interval 1940 to 1970. This represents a correspondence in the response of the growth of the tree ring time series, for the interval of more intense solar activity. Murphy (1990), in studies on tree ring data from Australia, similarly have found low correlation coefficients between tree ring data and Sunspot number, values between 0.16 and 0.22. Murphy and Veblen (1992) also have found a correlation coefficient of 0.2 between tree ring data, from Colorado – USA, and sunspot number. They have attributed a strong 11-year periodic association between these two time series. The results found by us show a lag of zero year with a high positive correlation between the solar activity and tree ring data from Southern Brazil region. In the Fig. 3a–b show that the tree-ring index increases and decreases when the *Rz* increases and decreases. It is also observed in long tendency temperature anomaly (Fig. 3c), where the lag with tree ring index is one

220

year with a high positive correlation.

When the solar activity is maximum there is a higher radiative output of the Sun, at least as it was verified by satellite observations in the last two decades (Frölich and Lean, 1998). This higher radiative emission causes a heating in the most of Earth but it could make some regions to become colder. In the regions with heating during solar maximum, one may have two situations. In the warm latitudes, the heating might causes a smaller tree ring growth and an anti-correlation between tree ring width and sunspot number might be observed. In the cold latitudes, the heating might causes a higher tree ring growth and thus a positive correlation with sunspot number is observed. Thus, the regional/local conditions could influence on the climate response to the solar forcing (radiative) in this region.

The Southern Brazil is located in the temperate climate region. It presents rigorous winter and summer periods. Temperature and other climate data are not available for this region in the long period studied in this work (almost 200 years). One can only make some hypothesis about the climate response in this region to the solar forcing, based on the tree ring width variations. Thus, a small increase of the temperature in this area could favor in the growth of the trees, while a decrease of the temperature could cause an inhibition in the growth of the trees. The correlations obtained between tree ring date  $x Rz$ , for long periods ( $>10$  years), and tree ring dates  $x$  temperature anomaly for long and short periods (Fig. 4b), suggest that the activity solar cause warming, through a more radiation emission to every maximum of solar activity (Wilson and Hudson, 1988; Frölich and Lean, 1998), that is to say a temperature increase, in the south Brazil region, in which influence an increase in the growth of the tree-rings.

Figure 3d and multivariate linear regression (Table 2) show an anti-correlation between tree ring index and SOI index with a lag of 2 years. It represents that the tree rings present a greater growth in epochs of El Niño (Daí et al., 2000). The south of Brazil presents a larger amount of precipitation in epochs of El Niño. Trees have a larger amount of water available during summer causing a larger growth of its rings.

Figure 3e and multivariate linear regression (Table 2) show that the volcanic activity

presents a factor of negative influence in the tree rings growth for long periods. It is comprehensible because the main influence of the volcanic aerosols is of causing a decrease in the temperature of the planet (Mass and Portman, 1989; Dutton and Christy, 1992). This decrease of the temperature would cause an inhibition in the growth of the trees giving an anti-correlation between tree ring index and SAOD.

The signal of the multivariate linear regression to short periods (Fig. 4) show that the tree ring are more influenced by the temperature than the SOI index and SAOD. Temperature present a cross-correlation coefficient of 0.74 with a zero year lag. It shows that the tree ring growth is more sensitive to temperature, that a phenomenon of global scale, as the volcanic eruptions and the SOI Index.

Solar activity, global temperature anomalies, ENSO and volcanic forcing explain nearly four-fifth (83%) of the variation in the tree-ring tendency above ten years for the ten-year running averages. For the low periods, among 2 to 5 years, the global temperature anomalies, ENSO and volcanic forcing explain nearly two-thirds (60,3%) of the variation in the tree-ring data.

The correlations obtained between tree ring date  $x Rz$ , for long periods ( $>10$  years), and tree ring dates  $x$  temperature anomaly for long and short periods (Fig. 4b), suggest that the solar activity cause more warming, through more radiation emission at every solar activity maximum, that is to say a temperature increase, in Southern Brazil region, which causes an increase of tree growth ring. The multivariate linear regression (Table 2) showed an anti-correlation between tree ring index and SOI index so that that tree rings present a greater growth in epochs of El Niño. The volcanic activity presents a factor of negative influence in the tree rings growth for long periods inhibiting the growth of the trees, giving an anti-correlation between tree ring index and SAOD.

The signal of the multivariate linear regression to short periods (Table 1) shows that tree rings are more influenced by the temperature than the SOI index and SAOD. That shows that the tree ring growth is more sensitive to the temperature, than to volcanic eruptions and SOI Index.

*Acknowledgements.* Thanks to Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq for support granted to this research (Project APQ 474185/2003-6, PQ 300992/2003-3) and research fellowships (PDJ 150102/2005-4) and to FAPESP post doctoral research Project 02/12723-2.

## 5 References

- Dai, A. and Wigley, T. M. L.: Global Patterns of ENSO-induced Precipitation, *Geophys. Res. Lett.*, 27, 1283–1286, 2000.
- Dutton, E. G. and Christy, J. R.: Solar radiative forging at selected locations and evidence for global lower tropospheric cooling following the eruptions of El Chichón and Pinatubo, *Geophys. Res. Lett.*, 19, 2313–2316, 1992.
- Enfield, D. B.: El Niño, past and present, *Rev. Geophys.*, 27, 159–187, 1989.
- Fritts, H. C.: *Tree rings and climate*, Academic Press Inc., London, 1976.
- Frölich, C. and Lean, J.: The sun's total irradiance cycles, trends and related climate change uncertainties since 1976, *Geophys. Res. Lett.*, 97, 7579–7591, 1998.
- Hamon, B. V. and Godfrey, J. S.: The role of the oceans, in: *Climatic Change and Variability*, edited by: Pittock, A. B. et al., A Southern Perspective, Cambridge, Cambridge University Press, 31–52, 1978.
- Hansen, J. and Lebedeff, S.: Global surface air temperatures: Update through 1987, *J. Geophys. Res.*, 15, 323–326, 1988.
- Hansen, J. and Lebedeff, S.: Global trends of measured surface air temperature, *J. Geophys. Res.*, 92, 13 345–13 372, 1987.
- Hoyt, D. V. and Schatten, K. H.: *The role of the sun in climate change*, Oxford University Press, Oxford, 280 p., 1997.
- Jones, P. D., Raper, S. C. B., Goodess, C. M., Cherry, B. S. G., and Wigley, T. M. L.: A grid point surface air temperature data set for the Southern Hemisphere, DOE Technical Report N<sup>o</sup> TR027, Washington, D. C., US Dept. of Energy, Carbon Dioxide Research Division, 73 p., 1986.
- Lacis, A., Hansen, J., and Sato, M.: Climate forcing by stratospheric aerosols, *Geophys. Res. Lett.*, 19, 1607–1610, 1992.

223

- Mass, C. F. and Portman, D. A.: Major volcanic eruptions and climate: A critical evaluation, *J. Climate*, 2, 566–593, 1989.
- Murphy, J. O. and Veblen, T. T.: Proxy Data from Tree-ring time series for the eleven-year solar cycle, *Proceedings ASA*, 10, 64–67, 1992.
- Murphy, J. O.: Australian tree ring chronologies a proxy data for solar variability, *Proceedings ASA*, 8, 292–297, 1990.
- Neelin, J. D. and Latif, M.: El Niño dynamics, *Physics Today*, 32–36, 1998.
- Pittock, A. B.: An overview, in: *Climatic Change and Variability*, edited by: Pittock, A. B. et al., A Southern Perspective, Cambridge, Cambridge University Press, 1–6, 1978.
- Rigozo, N. R. and Nordemann, D. J. R.: Solar activity records in tree-rings, *Revista Geofísica – Instituto Panamericano de Geografía e Historia (México)*, 50, 163–170, Enero-Junio, 1999.
- Rigozo, N. R., Nordemann, D. J. R., Echer, E., Vieira, L. E. A., and Prestes, A.: Tree Ring Image Interactive Treatment, in: *6th Latin America Conference on Space Geophysics*, GA-26, Concepción, 2001.
- Rigozo, N. R., Nordemann, D. J. R., Echer, E., and Vieira, L. E. A.: Search for Solar Periodicities in Tree-Ring Widths from Concórdia (S.C., Brazil), *Pure Applied in Geophysics*, 161, 221–233, 2004.
- Rigozo, N. R., Nordemann, D. J. R., Echer, E., Zanandrea, A., and Gonzalez, W. D.: Solar variability effects studied by tree-ring data wavelet analysis, *Adv. Space Res.*, 29, 1985–1988, 2002.
- Rigozo, N. R.: *Registros da atividade solar e de outros fenômenos geofísicos em anéis de árvores*, PhD Thesis (Doctorate in Space Geophysics) – National Institute for Space Research, Brazil, 132 p., 1998.
- Sato, M., Hansen, J. E., McCormick, M. P., and Pollack, J. B.: Stratospheric aerosol optical depths, 1850–1990, *J. Geophys. Res.*, 98, 22 987–22 994, 1993 and updates.
- Wilson, R. C. and Hudson, H. S.: Solar luminosity variations in solar cycle 21, *Nature*, 332, 810–812, 1988.

224

**Table 1.** Multiple Regression Results: 2 to 5 years band pass filter.

R	0.777			
R-Square	0.604			
Prob>F	<0.0001			
Parameter	Value	Error	<i>t</i> -Value	Prob>  <i>t</i>
Intercept	5.0504E-4	0.01009	-0.05007	0.9604
Temperature	0.07035	0.00972	7.23747	<0.0001
SOI	-0.00448	0.00201	-2.22962	0.0321
SAOD	4.3191E-5	7.246E-4	0.05961	0.9528

**Table 2.** Multiple Regression Results to ten-year running averages.

R	0.917			
R-Square	0.842			
Prob>F	<0.0001			
Parameter	Value	Error	<i>t</i> -Value	Prob>  <i>t</i>
Intercept	11.88638	1.9253	6.17387	<0.0001
Sunspot	0.00211	4.769E-4	4.42098	<0.0001
Temperature	-0.77845	0.1366	-5.69957	<0.0001
SOI	-0.02877	0.0042	-6.88016	<0.0001
SAOD	-0.00513	7.493E-4	-6.84379	<0.0001




Fig. 1. Time series.

227




Fig. 2. Tree-ring index spectral analysis by iterative method.

228




Fig. 3. Ten-year running averages.

229




Fig. 4. 2 to 5 years band pass filter.

230