

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).
Please refer to the corresponding final paper in CP if available.

Vegetation history of Central Chukotka deduced from permafrost paleoenvironmental records of the El'gygytgyn Impact Crater

A. A. Andreev¹, E. Morozova², G. Fedorov², L. Schirrmeister³, A. A. Bobrov⁴,
F. Kienast⁵, and G. Schwamborn³

¹Institute of Geology and Mineralogy, University of Cologne, Zùlpicher St. 49a, 50674, Cologne, Germany

²Arctic and Antarctic Research Institute, Bering St. 38, St. Petersburg, 199397 Russia

³Alfred Wegener Institute for Polar and Marine Research, Department of Periglacial Research, Telegrafenberg A43, 14473 Potsdam, Germany

⁴Faculty of Soil Science, Moscow State University, Vorobiev Gory, 119899 Moscow, Russia

⁵Senckenberg, Research Institute and Natural History Museum, Research Station for Quaternary Paleontology, am Jakobskirchhof 4, 99423 Weimar, Germany

Received: 5 March 2012 – Accepted: 12 April 2012 – Published: 20 April 2012

Correspondence to: A. A. Andreev (aandreev@uni-koeln.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

1409

Abstract

Frozen sediments from three cores bored in permafrost surrounding of the El'gygytgyn Impact Crater Lake have been studied for pollen, non-pollen palynomorphs, plant macrofossils, and rhizopods. The palynological study of the cores contributes to a higher resolution of time intervals presented in a poor temporal resolution in the lacustrine sediments; namely the Allerød and succeeding periods. Moreover, permafrost records better reflect local environmental changes, thus, allowing more reliable reconstruction of the local paleoenvironments. The new data confirm that shrub tundra with dwarf birch, shrub alder and willow dominated in the lake surroundings during the Allerød warming. Younger Dryas pollen assemblages reflect abrupt changes to grass-sedge-herb dominated environments reflecting significant climate deterioration. Low shrub tundra with dwarf birch and willow dominate the lake vicinity at the onset of the Holocene. The finds of larch seeds indicate its local presence around 11 000 cal. yr BP and, thus a northward shift of treeline by about 100 km during the early Holocene thermal optimum. Forest tundra with larch and shrub alder stands grew in the area during the early Holocene. After ca. 3500 cal. yr BP similar-to-modern plant communities became common in the lake vicinity.

1 Introduction

El'gygytgyn Impact Crater is located in Central Chukotka, approximately 100 km north of the Arctic Circle (Fig. 1). The crater was formed 3.6 Myr ago (Gurov and Gurova, 1979; Layer, 2000). As inferred from geomorphologic research, the study area was never glaciated after the time of the impact ca. 3.6 Myr ago (e.g. Brigham-Grette et al., 2007 and references therein). Thus, the lake is probably the longest archive for Arctic terrestrial environmental and climate history. Elgygytgyn Late Quaternary lacustrine palynological records were first reported by Shilo et al. (2001) following more continuous and detailed records published by Lozhkin et al. (2007) and Matrosova (2009).

1410

The studied sediments comprise the oldest continuous Quaternary pollen record in the Arctic, which provides history of vegetation and climate changes since ca. 350 kyr.

Generally, pollen records from large and deep lake sediments are very valuable paleoenvironmental archives containing unique data about vegetation and climate history. However, such pollen records reflect predominately regional environmental changes because of the large input of long distance wind-transported pollen into the spectra. The Lake El'gygytgyn sediments, where the pollen from a several thousand square-kilometer source area is trapped, provide a reliable record of over-regional vegetation and climate changes as well (Lozhkin et al., 2007; Matrosova, 2009). The importance of such continuous long-term regional records is obvious. Nevertheless, the short-term palynological records reflecting the local paleoenvironmental dynamics are also highly desired. These records document predominately changes in local vegetation and may be compared with over-regional variations in order to better understand the role of local and regional vegetation in the paleobotanical records, which results in more reliable environmental reconstructions.

Palynological studies of surface samples from the study area essentially complement reliable reconstructions. A total of 56 surface sediment samples from Lake El'gygytgyn and 26 surface soil samples from the crater slopes have been recently studied (Matrosova et al., 2004; Matrosova, 2006, 2009; Glushkova et al., 2009). These studies demonstrate that pollen of trees and shrubs may reach up to 82 % of the recent lacustrine spectra although the only willow and dwarf birch stands grow in the crater in protected locations. The soil pollen spectra more reliably reflect the local vegetation but pollen of long-distance-transported taxa dominate even there (Matrosova, 2006; Glushkova et al., 2009). It is characteristic that pollen contents of *Pinus pumila* and *Alnus fruticosa*, species not growing in the crater vicinity may reach up to 15 and 37 % of the spectra consequently. Thus, by interpretations of fossil pollen assemblages it has to be taken in consideration that significant part of the pollen may have been originated from some dozens and even hundreds of kilometers away.

1411

This paper presents palaeoenvironmental and palaeoclimatic changes during the Lateglacial and Holocene inferred from permafrost pollen, plant macrofossil, and rhyzopod records from the permafrost surrounding of the El'gygytgyn Crater Lake. The Lateglacial/Holocene transition is considered as a unique period of intensive glaciation and deglaciation events accompanied by remarkable changes in global temperature, atmospheric circulation, air humidity, precipitation and vegetation changes (Johnsen et al., 1995; Stuiver et al., 1995; Blunier and Brook, 2001). Our studies of three permafrost cores add to a better understanding of paleoenvironmental changes during the time intervals, which are not well presented in a good temporal resolution in the lacustrine archive. A comparison of the palynological data from the new permafrost cores and previously studied exposures and lake cores were used to make a local chronostratigraphy scheme because of the partly insufficient geochronological datasets. Such comparison resulted in a more reliable reconstruction of vegetation and climate changes, especially during the transitional intervals from cold to warm periods.

2 Geographical setting

The El'gygytgyn Impact Crater is 18 km in diameter and holds a ca 170 m deep lake that has almost a round morphology with 11 km in diameter (Fig. 1). The crater is superimposed on the Anadyr lowland and was formed in an Upper Cretaceous volcanic plateau (Belyi, 1998). The crater rim comprises peaks between 600 and 930 m above sea level (a.s.l.), and the lake level is situated at 492 m a.s.l. Unconsolidated Quaternary permafrost deposits cover the crater bottom surrounding the lake. They show a distinctly asymmetrical distribution with a broad fringe of loose sediment that is 500 to 600 m wide in the north and west and only 10 to 20 m elsewhere around the lake (Fig. 1).

The study area belongs to the continuous permafrost zone with a mean annual ground temperature of -10°C at 12.5 m depth (Schwamborn et al., 2008). In 2003, the active layer was about 40 cm deep in peaty silts and reached 50 to 80 cm in sand,

1412

pebbles, and gravels. The region is characterized by extremely harsh climate with average annual air temperature ca. -10°C , mean July temperatures of 4 to 8°C and mean January temperatures of -32 to -36°C . The precipitation consists of 70 mm summer rainfall (June–September) and ca. 110 mm water equivalent of snowfall (Nolan and Brigham-Grette, 2007). Characterizing the modern climate around El'gygytgyn Crater Lake, an oceanic influence is distinctively expressed in decreasing summer temperatures and resulted in the thermal inertia of the crater area (Kozhevnikov, 1993). According to Kozhevnikov (1993), long-distance atmosphere convection bringing air masses from the south and north dominates at the lake area. These air masses bring tree and shrub pollen grains playing an important role in the recent pollen assemblages from long distances. This situation might also be occurred the past.

The modern treeline is positioned roughly 100 km to the south and west of the lake. However, single shrub alder stands were found only in approximately 10 km from the lake, in the Enmyvaam River valley (P. Minyuk, personal communication, 2009). The local vegetation has been well studied during last decades (e.g. Belikovich, 1988, 1989, 1994; Kozhevnikov, 1993; Belikovich and Galanin, 1994 and references therein). The study area belongs to the subarctic tundra zone and enters into a subzone of the southern subarctic tundra.

According to Belikovich (1994), ca. 40% of the area (low parts of smooth crater slopes and low lake terraces) are covered by hummock tundra with *Eriophorum vaginatum*, *E. callitrix*, *E. polystachion*, *Pedicularis pennellii*, *P. albolabiata*, *Carex rotundata*, *C. lugens*, *Salix fuscescens*, *S. reticulata*, *Senecio atropurpureus*, *Ledum decumbens*, *Andromeda polifolia*, and *Vaccinium uliginosum*; ca. 20% (low-middle parts of crater slopes) by moss-lichens tundra with *Cassiope tetragona*, *Rhododendron parvifolium*, *Senecio resedifolus*, *Ermania parryoides*, *Silene stenophylla*, *Dryas octopetala*, *Crepis nana*, *Potentilla elegans*, and *Androsace ochotensis*; ca. 15% (upper mountain plains) – by tundra with rare beds with *Salix phlebophylla*, *Pedicularis lanata*, *Artemisia furcata*, *Potentilla elegans*, *Eritrichium aretioides*, *Minuartia arctica*, *Potentilla uniflora*, *Arenaria capillaris*, *Poa pseudoabbreviata*, *Cardamine bellidifolia*,

1413

Saxifraga serpyllifolia, *Kobresia myosuroides*, and *Crepis nana*; ca. 10% – by nival vegetation with *Salix polaris*, *Cassiope tetragona*, *Carex tripartita*, *Phippsia algida*, *Koenigia islandica*, *Saxifraga hyperborea*, *Eritrichium villosum*, *Primula tschuktschorum*, *Hierochloa pauciflora*; ca. 10% – by meadow and shrubby tundra with *Artemisia arctica*, *Aconitum delphinifolium*, *Arctagrostis arundinacea*, *Carex podocarpa*, *Festuca altaica*, *Luzula multiflora*, *Senecio tundricola*, *Thalictrum alpinum*, *Veratrum oxysepalum*. Rare steppe-like communities with *Potentilla stipularis*, *Artemisia kruhseana*, *Myosotis asiatica*, *Saxifraga eschscholtzii*, *Papaver lapponicum*, *Senecio jacuticus*, *Woodsia ilvensis*, *Dianthus repens* can be found in rocky habitats. Along the Enmyvaam River and some large creeks grow low willow stands with *Salix tschuktschorum*, *S. saxatilis*, *Androsace ochotensis*, *Empetrum subholarcticum*, *Pleuropogon sabinii*, *Polemonium boreale*, *Beckwithia chamissonis*, *Saussurea tilesii*, *Lagotis minor*, *Pedicularis hirsuta* and meadow-shrub willow communities with *Salix alaxensis*, *S. krylovii*, *Deschampsia borealis*, *Chamerion latifolium*, *Equisetum variegatum*, *Stellaria fischerana*, *Potentilla hyperctica*, *Eutrema edwardsii*, *Cardamine blaisdellii*, *Trollius membranostylus*, *Polemonium acutiflorum*, *Parnassia kotzebuei*, *Poa paucispicula*.

3 Methods

A standard HF technique was used for pollen preparation (Berglund and Ralska-Jasiewiczowa, 1986). A tablet of *Lycopodium* marker spores was added to each sample for calculating total pollen and spore concentrations following Stockmarr (1971). Water-free glycerol was used for sample storage and preparation of the microscopic slides. Pollen and spores were identified at magnifications of $400\times$, with the aid of published pollen keys and atlases (Kupriyanova et al., 1972, 1978; Bobrov et al., 1983; Reille, 1992, 1995, 1998; Beug, 2004). In addition to pollen and spores a number of non-pollen-palynomorphs such as fungi spores, remains of algae and invertebrate, were also identified when possible and counted. These non-pollen-palynomorphs are also valuable indicators of past environments (e.g. van Geel, 2001 and references therein).

1414

At least 250 pollen grains were counted in each sample. The relative frequencies of pollen taxa were calculated from the sum of the terrestrial pollen taxa. Spore percentages are based on the sum of pollen and spores. The relative abundances of reworked taxa (mineralized pollen and spores of Tertiary and early Quaternary age) are based on the sum of pollen and redeposited taxa, the percentages of non-pollen palynomorphs are based on the sum of the pollen and non-pollen palynomorphs, and the percentages of algae are based on the sum of pollen and algae. The TGView software (Grimm, 2004) was used for the calculation of percentages and for drawing the diagrams. The diagrams were zoned by a qualitative inspection of significant changes in pollen associations, pollen concentrations and occurrence of particularly indicative taxa. The CorelDraw software was used for preparation of the final pollen diagrams.

In a depth of 146.5–151 cm in core P2, we detected a number of well preserved plant remains. They were picked using a stereomicroscope and identified by comparison with modern reference material from the Herbarium Senckenbergianum (IQW). Additionally, a *Carex* identification key (Egorova, 1999) has been used.

The core sediments were also studied for testate amoebae shells. The samples were sieved through a 0.5 mm mesh and testate amoebae shells were concentrated with a centrifuge. A drop of suspension was placed on the slide, and then glycerol was added. Normally, 5 slides were examined at X200-400 magnification with a light microscope.

A total of 33 AMS ^{14}C ages have been obtained from the studied deposits (Table 1). Plant macrofossils (i.e. grass remains) were picked from the cores P1 and P2 and the uppermost segment of 5011-3 for AMS radiocarbon dating. Because of the lack of plant remains in the lower part of core 5011-3 only bulk organic was dated there. AMS datings were done at the Leibniz Laboratory for Radiometric Dating and Stable Isotope Research (Christian Albrechts University, Kiel, Germany) and the Poznan Radiocarbon Laboratory (Adam Mickiewicz University, Poznan, Poland). Calibrated ages (cal. yr BP) were calculated using “CALIB Rev 6.1.0.” (Reimer et al., 2009).

1415

4 Results

4.1 P1 core

The first permafrost core (P1) has been extracted from a piedmont terrace about 1.7 km southeast of the lake (67°22'26" N, 172°13'10" E, Fig. 1) during the field work in summer 2003 (for details see Schwamborn et al., 2006). The study site is located on a slope exposed to the southwest with the angle of 5°. The vegetation cover at core site was relatively dense (ca. 80%).

The 5 m slope debris core mostly consists of a silty-to-sandy diamicton interpreted as a result from proluvial, colluvial and solifluctional deposition (Schwamborn et al., 2006). Prominent peaty layers interrupt the section between 330 and 220 cm core depth, which is also reflected in maximum values of total organic carbon (TOC on Fig. 2). Non-identified plant remains from several layers have been dated and show a correct depth-to-age relationship (Table 1, Fig. 2). The oldest date from 463 cm depth shows that the oldest core sediments are around 13 000 cal. yr BP old or slightly older.

Generally the P1 sequence is very rich in pollen and palynomorphs (Fig. 3). The studied pollen spectra can be subdivided into 5 pollen zones (PZ). PZ-I (ca. 495–430 cm) is dominated by *Cyperaceae*, *Poaceae*, *Betula* sect. *Nanae* and *Salix* pollen. PZ-II (ca. 430–380 cm) shows the significant increase of *Cyperaceae* pollen content, while *Betula* sect. *Nanae* content is decreased. PZ-III (ca. 380–330 cm) is notable for an increase in *Betula* sect. *Nanae* and appearance of small amounts of *Alnus fruticosa*. Pollen concentration is also increased in the upper part of the zone. The amounts of tree and shrub pollen (predominantly *Alnus fruticosa*) have a maximum in PZ-IV (ca. 330–265 cm). The pollen concentration is the highest in PZ-V (ca. 265–50 cm), which is notable for high amounts of *Betula* sect. *Nanae*, *Alnus fruticosa* and *Cyperaceae* pollen. Pollen of long-distance transported *Pinus*, *Larix* and *Picea* is also characteristic for this zone. PZ-V can be subdivided into 2 subzones, the upper one (50–0 cm) shows the higher contents of *Salix* pollen.

1416

The P1 has been also studied for rhizopods (Table 2). The only sphagnobiotic/hygrophilic *Heleopera petricola* v. *amethystea* pointing to a very wet environment has been found on the 463–473 cm depth. Mostly soil-eurybiotic (e.g. *Centropyxis aerophila*, *C. constricta*, *C. sylvatica*) and hydrophilic (*Diffflugia* and *Lagenodiffflugia*) species dominated the sediments between 334 and 223 cm. However, sphagnobiotic taxa (*Arcella*, *Heleopera*, *Nebela*, *Centropyxis aculeata*) are also common. The role of soil-eurybiotic species gradually increases in the upper part.

4.2 P2 core

The core has been retrieved at 12.5 km far from P1 across the lake to the north (67°32'50" N, 172°07'31" E, Fig. 1). The site is placed on a gently inclined (<3°) surface about 100 m from the north lake shoreline (for details see Schwamborn et al., 2008a). The surface is characterized by a boggy environment composed of a loamy substrate covered by grass tundra. Similarly like core P1 deposits, core P2 is composed of a silty-to-sandy diamicton deposition (Schwamborn et al., 2008b). The lower part of the core (510–250 cm) is interpreted as weathering debris of the local volcanic basement. The upper 250 cm consisted of proluvial slope wash out deposits. The lithological transition between the units is also very distinguishable by an increase of TOC contents (Fig. 2).

Non-identified plant remains found in the P2 deposits have also been dated and show a rather reliable depth-age relationship (Table 1, Fig. 2). Three radiocarbon dates from the sediments between 205 and 226 cm depth demonstrates that these sediments might be accumulated at about 14 000–12 400 cal. yr BP. The youngest date seems as the most reliable taking in consideration the comparison with other dated pollen records from the area (e.g. Matrosova, 2009; Glushkova et al., 2009 and references therein).

P2 core sediments are rich in pollen and palynomorphs except of the lowermost 170 cm. The studied pollen spectra can be subdivided into 6 PZ (Fig. 4). Sediments from PZ-I (ca. 510–350 cm) contain only single pollen grains of *Pinaceae*, *Betula* sect. *Nanae*, *Alnus fruticosa*, and *Cyperaceae*. Pollen concentration is slightly higher in the lowermost sample which contains few pollen of *Betula* sect. *Nanae*, *Alnus*

1417

fruticosa, *Pinus* s/g *Haploxyylon* and *Cyperaceae*. Pollen concentration is higher (up to 2650 grains/g) in PZ-II (ca. 350–330 cm), which is also notable for high content of *Lycopodium* and *Botrychium* spores. The lowermost PZ-I and PZ-II were not used for paleoenvironmental reconstructions because of very low pollen concentration in many samples which may lead to over-representing some taxa due to possible contamination or selective preservation of palynomorphs (e.g. abnormal presence of spores may indirectly point to it). Pollen concentration is much higher (up to 5800 grains/g) in PZ-III (ca. 330–265 cm), which is characterized by high pollen contents of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Cyperaceae*, *Poaceae*. Rather high amounts of *Pinus* s/g *Haploxyylon* and *Pinaceae* is also notable in this zone. The pollen concentration increases significantly (up to 35 700 grains/g) in PZ-IV (ca. 265–180 cm). *Betula* sect. *Nanae* and *Alnus fruticosa* pollen contents decreased dramatically at the beginning of the zone and gradually increased in the upper part. The zone can be subdivided in two subzones basing on the shrub pollen contents. Pollen concentration is the highest (up to 83 600 grains/g) in PZ-V (ca. 180–40 cm), which is dominated by pollen of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, *Cyperaceae*, and *Poaceae*. Additionally, on the 146.5–151 cm depth were found seeds and short purs of *Larix dahurica* as well as numerous utricle and nutlets of *Carex rostrata*. The uppermost PZ-VI (ca. 40–0 cm) is characterized by decreasing *Betula* sect. *Nanae* and *Alnus fruticosa* pollen contents, while *Cyperaceae*, *Pinus* s/g *Haploxyylon* and *Salix* pollen contents increased.

The P2 core has been also studied for rhizopods, but no shells were found there.

4.3 5011-3 core

The core has been drilled on the western margin of the crater (67°29'04" N, 171°56'40" E) at about 300 m west from the lake shore (Fig. 1). This 141.5 m long core was recovered during a drilling campaign in winter 2008 in the frame of the international ICDP funded project "El'gygytgyn Drilling Project" (Melles et al., 2011). The main objective of the coring was to extend the permafrost record back in time in order to better understand the interaction between catchment processes and lake sedimentation. The

1418

sediment core drilled in an alluvial fan consists of sediment layers of sandy gravel to gravelly sand and is interpreted to represent alternating subaerial and subaquatic parts of the fan. Occasionally intercalations of sandy beds occur, e.g. at 7, 9, 14.5, 18–19.5, 24, and 26 m. The modern setting of the coring site is placed in an alluvial-proluvial sediment fan and from aerial imagery it is obvious that the fan has a subaquatic prolon-
5 gation into the lake. In total, 12 samples from the core were AMS ^{14}C dated (Table 1). Although the non-identifiable plant remains (possibly grass roots) were picked through-
out the upper meter of the core and expected to provide reliable age control for studied
sediments, the ages appeared to be modern reflecting the presence of modern plant
10 roots in the active layer. The bulk AMS ^{14}C dates from some selected horizons (Ta-
ble 1) did not provide reliable ages either. These ages are not in a chronological order
reflecting the reworked character of TOC in the samples. The ages are obviously too
old taking in consideration the comparison with other dated pollen records from the
area (e.g. Matrosova, 2009; Glushkova et al., 2009; P1 and P2 records). Therefore, an
15 age estimations for the 5011-3 core are based on a comparison with the dated pollen
sequences from the area.

Generally, the upper 9 m of 5011-3 sediments are rich in pollen and palynomorphs, but only single pollen grains were found below this depth except sediments between 19.8 and 19.3 m (Fig. 5). The studied pollen spectra can be subdivided into 7 PZ.
20 PZ-I (ca. 1980–1930 cm) is dominated by *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*,
Cyperaceae, *Poaceae*, and *Ericales* pollen. Presence of *Larix* pollen and high con-
tents of *Sphagnum* and *Lycopodium* spores is also characteristic for the zone. Pollen
concentration is rather low (up to 3500 grains/g). No pollen has been found between
ca. 1930 and 1400 cm and only few pollen grains of *Betula* sect. *Nanae*, *Alnus fruti-
25 cosa*, *Salix*, *Cyperaceae*, *Poaceae* and single spores of *Sphagnum* and *Lycopodium*
have been found in PZ-II (ca. 1400–900 cm). PZ-III (ca. 900–330 cm) is notable for
much higher pollen concentration (up to 101 330 grains/g). The spectra are dominated
by *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, *Cyperaceae*, *Poaceae*, *Ericales* and
spores of *Sphagnum*. Contents of *Sphagnum* as well as pollen concentration reduced

1419

significantly in the upper PZ-IV (ca. 330–250 cm). PZ-V (ca. 250–180 cm) is notable
for the significant increase of *Poaceae* pollen content, while contents of *Betula* sect.
Nanae, *Alnus fruticosa*, *Salix*, *Ericales* and *Sphagnum* are dramatically decreased.
The pollen concentration is the highest in the zone (up to 829 400 grains/g). The con-
5 tents of *Betula* and *Alnus* pollen increased again in PZ-VI (ca. 180–100 cm), which is
also notable for high content of *Artemisia*. The pollen concentration significantly (up to
15 000 grains/g) reduced in this zone. The uppermost PZ-VII (100–0 cm) is dominated
by of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Cyperaceae*, *Poaceae*, and *Ericales*. The
pollen concentration is very high in the zone (up to 770 000 grains/g). Single pollen of
10 long-distance transported *Pinus* and *Picea* are also characteristic for this zone.

The 5011-3 core has been also studied for rhizopods, but no shells were found.

5 Discussion: palaeoenvironmental reconstructions

5.1 MIS 7(?) environment

The oldest pollen spectra are presented in the lower part (1980 to 1930 cm) of the
15 studied section of the core 5011-3 (PZ-I, Fig. 5). The pollen assemblages are domi-
nated by *Alnus fruticosa*, *Betula* sect. *Nanae* and *Poaceae*. However, pollen of *Salix*,
Cyperaceae, *Ericales*, Caryophyllaceae and spores of *Sphagnum*, *Lycopodium* and
Huperzia are also important components of the revealed spectra. They are not dated
but the comparison with lacustrine pollen records shows that spectra of our PZ-I are
20 very similar to those from the zone E14 of the TL-dated lacustrine core LZ1024 (Ma-
trosova, 2009). Basing on the comparison of our record with the LZ1024 one, we may
suggest a MIS 7 age for our PZ-I zone. However, an older age for the revealed inter-
glacial interval cannot be completely excluded.

According to the pollen spectra, shrub alder, dwarf birch, willows and larch grew
25 in the lake catchment that requires the movement of northern boundary of larch for-
est at least 100 km to the north. Our conclusion is also supported by the lacustrine

1420

pollen records (Lozhkin et al., 2007; Matrosova, 2009). However, the cores drilled in the center of the El'gygytgyn Lake do not contain larch pollen at all and show low presence of *Salix*, *Ericales*, Caryophyllaceae pollen and *Sphagnum*, *Lycopodium* and *Huperzia* spores. This difference most likely reflects the larger presence of the local components in the 5011-3 core pointing to the importance of studying of the terrestrial (non-lacustrine) sediments in addition to the lacustrine ones. Taking in consideration all El'gygytgyn pollen records, we assume that open larch forest with shrub alder, dwarf birch and willows dominated the local vegetation during the revealed warm interval. However, grass-sedge dominated communities with other herbs and *Sphagnum* and *Lycopodium* growing in mesic habitats were also common in lake vicinity.

5.2 Lateglacial

Lateglacial sediments are revealed in both radiocarbon dated slope cores (P1 and P2) and in the long permafrost 5011-3 core. Unfortunately, we have a rather good age control only for the lowermost part of the core P1. Taking into consideration the P1 bottom age of $11\,160 \pm 70$ ^{14}C yr BP (12 283–13 424 cal. yr BP), the most reliable P2 age of $10\,450 \pm 60$ ^{14}C yr BP (12 124–12 654 cal. yr BP), and pollen-based correlation with lacustrine pollen records (zone E4 of LZ1024 in Matrosova, 2009) we may assume that our PZ-I of P1 (Fig. 3), PZ-III of P2 (Fig. 4) and PZ-III and PZ-IV of 5011-3 (Fig. 5) accumulated during the Allerød, before 13 cal. kyr.

Sediments attributed to the Allerød are dominated by pollen of *Betula* sect. *Nanae*, *Alnus fruticosa*, *Salix*, *Cyperaceae*, *Poaceae*, *Ericales* and spores of *Sphagnum*. The relatively high pollen concentration is also characteristic for the sediments. However, a number of samples show very low pollen concentrations or do not contain pollen at all. Most likely that reflects a very high accumulation rate during the sedimentation. Such conclusion is in a good agreement with thicknesses of Allerød-attributed deposits of about 2.5 m in the P2 core and at least 6.5 m in the 5011-3 core. Warmer and wetter climate conditions in the Allerød might have intensified erosion and, therefore, higher

1421

influx of terrestrial material. The absence or very low thickness of the underlying Late Pleistocene sediments also might be connected with these erosion processes.

The main pollen taxa in the spectra slightly differ from site to site. For example 5011-3 sediments contain large amounts of *Salix* and *Ericales* pollen and *Sphagnum* spores; P1 and P2 sediments contain numerous pollen of *Cyperaceae*; lacustrine pollen records contain larger amounts of long-distance pollen (including *Betula* and *Alnus*). However, the PG1351 lacustrine record also contains large amounts of *Sphagnum* spores in the late glacial sediments confirming wet habitats in the lake vicinity (Lozhkin et al., 2007). The sphagnobiotic rhizopod, *Heleopera petricola*, found in the Allerød-dated P1 sediments is in a good agreement with numerous *Sphagnum* spores in the pollen records. Such habitats were probably common along the creeks as today.

Our interpretation of the studied sediments is very similar to those from LZ1024 done by Glushkova et al. (2009) and Matrosova (2009). Moreover, they also have reported pollen spectra with dominance of shrub pollen taxa from the not dated terrace sediments (sections GS-10 and GS-12/1 in Fig. 1) attributed to a Late Glacial warm interval. Similar paleoenvironmental records are also known from adjacent regions (e.g. Brubaker et al., 2005; Anderson and Lozhkin, 2006; Shilo et al., 2006, 2007; Kokorowski et al., 2008; Andreev et al., 2009 and references therein). Lozhkin et al. (2007) basing on their PG1351 lacustrine pollen record have suggested that birch was regionally present at about 12 800 yr ^{14}C BP (15 300 cal. yr BP), while alder established in the area only around 10 700 yr ^{14}C BP (12 700 cal. yr BP).

There are plant macrofossil data from the sediments of a section GS-37 (Fig. 1) ^{14}C dated to $12\,215 \pm 40$ yr BP (14 027–14 491 cal. yr BP). The studied sediments do not contain any shrub remains. Glushkova et al. (2009) interpreted it as the absolute absence of any shrub stands in the lake vicinity and very severe climate conditions. Thus, it seems that Allerød pollen and plant macrofossil data are controversial. However, the conclusion about herb dominated tundra vegetation around 14 250 cal. yr BP is based on the single studied sample, which reflect very wet, but not a typical tundra habitat. Moreover, they interpret the sediments containing numerous pebbles and eggs

1422

of *Daphnia* as the lake terrace periodically overflowed by the lake (Glushkova et al., 2009). It is obvious that shrubs cannot survive in such flooded habitats. Therefore, the found plant macrofossils reflect very local flooded habitat environmental condition, which cannot be extrapolated to the whole lake vicinity.

5 Thus, according to the pollen spectra shrub alder, dwarf birch, and willows grew in the lake surrounding during the Allerød interstadial with relatively warm and wet climate. We can reconstruct shrub tundra vegetation with dwarf birch, shrub alder and willow around the lake.

Pollen spectra from PZ-II of P1 (Fig. 3), PZ-IVa of the core P2 (Fig. 4), and PZ-V of 10 5011-3 (Fig. 5) are dominated mostly by *Cyperaceae* and *Poaceae* pollen and reflect disappearance of shrubs from the area pointing to climate deterioration which can be attributed to the Younger Dryas. The most reliable ^{14}C dates from core P2 and P1 (Table 1) well confirm this conclusion. Pollen spectra with significant increase in herbs (mostly *Poaceae*) and *Selaginella rupestris* have been also revealed in the lacustrine 15 sediments (E3 of LZ1024 in Matrosova, 2009) and interpreted as reflected the Younger Dryas event (Glushkova et al., 2009). Thus, grass-herb tundra dominated the area during the Young Dryas cooling. Younger Dryas dated pollen records from the adjacent regions (e.g. Anderson et al., 2002; Kokorowski et al., 2008; Andreev et al., 2009 and references therein) reflect the similar environmental changes.

20 5.3 Holocene

Pollen spectra of the PZ-IVb of P2 (Fig. 4) accumulated before 9640 ± 60 ^{14}C yr BP (11 200–10 780 cal.yr BP) show gradual increase of *Alnus fruticosa* and *Betula* sect. *Nanae* pollen contents reflecting early Holocene climate amelioration. The early Holocene pollen assemblages are also well present in the undated PZ-III of P1 (Fig. 3), 25 where they are dominated mostly by pollen of *Betula* sect. *Nanae*, *Cyperaceae* and *Poaceae* with few *Alnus fruticosa* and *Salix*. Four ^{14}C dates (Table 1) confirm that these sediments were accumulated before 9000 ^{14}C yr BP (10 200 cal.yr BP). Similar pollen assemblages have been revealed in the lowermost pollen zone of the so-called Olga

1423

Creek section (OC on Fig. 1, Shilo et al., 2008; Glushkova et al., 2009). This section is situated ca 100 m from P1 coring site. Their lowermost spectra are also not ^{14}C dated, however, two ^{14}C dates: 9250 ± 90 and 9125 ± 30 yr BP, from overlain sediments confirm that these sediments were accumulated before 9300 ^{14}C yr BP (10 550 cal.yr BP).

5 Similar undated early Holocene pollen assemblages are also reported by Glushkova et al., (2009) from the section GS-12/1 (Fig. 1). Thus, we may assume that shrub tundra with dwarf birches and willows and probably few shrub alder dominated the lake vicinity at the onset of the Holocene. The early Holocene pollen records from adjacent regions (e.g. Anderson et al., 2002; Lozhkin and Anderson, 2002; Kokorowski et al., 10 2008; Andreev et al., 2009 and references therein) have revealed similar environmental changes.

The contents of *Alnus fruticosa* are significantly higher in the PZ-V of the core P2 (Fig. 4) ^{14}C dated to ca 9600 yr BP (11 200–10 780 cal.yr BP) and PZ-IV of the core P1 (Fig. 3) ^{14}C dated around 8900–8800 yr BP (9940–9700 cal.yr BP). Most likely, this 15 increase reflects the further distribution of shrub alder stands in the area at the early Holocene. Pollen spectra of the PZ-V of P2 (Fig. 4) radiocarbon dated to about 7200–7300 cal.yr BP, PZ-VI of 5011-3 (Fig. 5) and bottom spectra from the terrace section GS-8403 (Glushkova et al., 2009) and the section OC in the Enmyvaam River valley (Glushkova and Smirnov, 2007; Shilo et al., 2008; Glushkova et al., 2009) also demonstrate high amounts of *Alnus fruticosa* pollen in the early Holocene sediments. Moreover, the lacustrine sediments (namely E2 of LZ1024 in Matrosova, 2009) accumulated above the sediments attributed to the Younger Dryas also contain very high amounts of *Alnus* (up to 60 %). Large shrub alder trunks and smaller twig fragments ^{14}C dated to 9250 ± 90 and 9125 ± 30 yr BP correspondingly as well as numerous undated alder 25 nuts from the same layers well confirm that shrub alder grew in the lake vicinity at least 10 550 cal.yr BP (Shilo et al., 2008). Thus, it is likely that shrub alder stands were well established in the El'gygytgyn Lake Crater at about 11 200 cal.yr BP or even slightly earlier.

The well-preserved larch seeds (Fig. 6) found in peaty layer in the core P2 prove the local presence of trees directly at the lake crater as early as 11 200–10 780 cal. yr BP. Larch remains were also found in the sediments accumulated shortly before 9300 ^{14}C yr BP (10 550 cal. yr BP) from the OC section (Fig. 1, Shilo et al., 2008; Glushkova et al., 2009), thus, also confirming local presence of the larch trees at the area during the early Holocene. Such forest (tundra-forest) environments are also good habitats for the shrub alder stands. The local presence of *Larix* indicates a treeline shift of about 100 km northward (CAVM-Team, 2003) as result of the early Holocene climate amelioration. Larch needs a mean temperature of the warmest month of at least 10 °C, thus such climate conditions must have existed at the lake crater during the early Holocene.

The studied early Holocene pollen assemblages slightly differ from site to site. For example, the early Holocene 5011-3 spectra (PZ-VI) show high contents of *Artemisia* (up to 25%), while GS-8403 spectra reported by Glushkova et al. (2009) contain up to 23% of *Ericales*. The difference may reflect the mosaic character of the local vegetation cover and/or different age of the revealed pollen assemblages. The lacustrine record (namely E2 of LZ1024 in Matrosova, 2009) accumulated above the sediments attributed to the Younger Dryas shows very high amounts of *Alnus* (up to 60%), which might have been transported from a distance and, thus, reflect the regionally dominated vegetation.

Rhizopod shells of soil-eurybiotic *Centropyxis* and hydrophilic *Diffflugia* taxa (Table 2) are numerous in the P1 early Holocene sediments, however, sphagnobiotic *Arcella*, *Heleopera*, and *Nebela* are also common. The high contents of hydrophilic and sphagnophilic taxa point to wet oligotrophic and mesotrophic soil environment at the core site. Later, after ca 6300 cal. yr BP, the role of soil-eurybiotic species increased reflecting drier soil environment.

Thus, pollen and macrofossil data show that forest and/or forest-tundra communities with larches, shrub alder, dwarf birches, and willows were well distributed the low elevations around the lake during the early Holocene at least between 11 200 and 9100 cal. yr BP. It is most likely that larch and shrub alder grew in the close vicinity

1425

to the lake only before ca. 8200 cal. yr BP. Similar changes in the high Arctic vegetation cover are also characteristic for coastal areas of the Laptev and East Siberian Seas (e.g. Andreev et al., 2009, 2011 and references therein). Found larch remains document that larch grew in approximately 100 km from its modern northern distribution limit. The mean July temperatures were at least 10–12 °C (Lozhkin and Anderson, 1995), ca. 4–5 °C higher than modern July temperatures (Shilo et al., 2008). This is in well agreement with the early Holocene pollen-based paleoclimate reconstruction from the El'gygytgyn lacustrine record (Lozhkin et al., 2007) and other high arctic sites (e.g. Andreev et al., 2009, 2011 and references therein).

A number of ^{14}C dates (Table 1) from P1 (PZ-Va) and P2 (PZ-V) cores confirm that permafrost sediments containing relatively high amounts of *Alnus fruticosa* pollen were accumulated until ca. 3500 cal. yr BP. Therefore we may assume that shrub alder might grow around the lake in more protected habitats or very close to the lake vicinity before this time. This conclusion is in a good agreement with pollen and plant macrofossil data from adjacent regions documenting the presence of shrubs and trees to the north from modern distribution areas (e.g. MacDonald et al., 2000; Andreev et al., 2009, 2011; Binney et al., 2009 and references therein). However, the dated woody remains from the Enmyvaam River valley (Glushkova and Smirnov, 2007; Lozhkin et al., 2011) confirm the presence of high shrubs in the area only until ca. 7400 ^{14}C yr BP (8200 cal. yr BP). The Enmyvaam River terrace deposits also contain the relatively high amounts of *Alnus fruticosa* pollen in the sediments accumulated after 7400 ^{14}C yr BP (Glushkova and Smirnov, 2007; Shilo et al., 2008; Lozhkin et al., 2011), pointing to a possible presence of alder in the vegetation cover, however, the age of the studied spectra is unknown.

Generally, late Holocene pollen spectra from the uppermost sediments (upper spectra of PZ-Vb of the core P1, Fig. 3; PZ-VI of the core P2, Fig. 4) show a decrease in contents of *Alnus fruticosa* and increase of contents of *Salix*, *Pinus*, *Betula*, *Ericales*, and *Cyperaceae*. These changes can be interpreted as disappearance of shrub alder from the lake vicinity. The main components of pollen assemblages slightly change

1426

from site to site reflecting local vegetation cover at coring sites. The higher amounts of *Pinus* s/g *Haploxylon* (most likely *Pinus pumila*) are of long distance origin. Its presence is especially remarkable in lacustrine records (Lozhkin et al., 2007; Matrosova, 2009) reflecting the over-regional vegetation pollen influx. Taking into consideration all pollen records from the study area we may assume that stone pine did not grow around the lake during the Holocene. Late Holocene pollen records from the lake surroundings (sites GS-12/1 and GS-12/2) radiocarbon dated between ca. 900 and 450 cal. yr BP (Glushkova et al., 2009) contain pollen spectra similar to those revealed in this study. They also show lower contents of *Alnus* pollen in many spectra and high fluctuations in *Betula*, *Ericales*, *Thalictrum*, and *Selaginella rupestris* reflecting local environments. Thus, pollen data show that herb tundra communities started to dominate in the lake catchment after ca. 3000 cal. yr BP.

6 Conclusions

New permafrost records document vegetation and climate changes in the El'gygytgyn Lake Crater during the Late Quaternary. The studied records well reflect the local vegetation changes that resulted in better understanding of the possible role of local and regional components in the fossil pollen spectra and in more reliable environmental reconstructions. It is evident that terrestrial records better reflect the local environments than the lacustrine ones there long-distance transported pollen are over-represent the local components.

The oldest pollen spectra of the studied sections of the core 5011-3 are possibly of the MIS 7 age. They document that open larch forest with shrub alder, dwarf birch and willows dominate vegetation suggesting the northern movement of larch forests. Treeless grass-sedge dominated communities with other herbs and *Sphagnum* and *Lycopodium* growing in mesic habitats were also common in lake vicinity.

Lateglacial pollen records show that shrub tundra with dwarf birch, shrub alder and willow dominated in the lake surroundings during the relatively warm Allerød

1427

interstadial. Rather low pollen concentration in many samples of Allerød age reflect very high accumulation rate during the sedimentation.

Younger Dryas pollen records reflect dramatic changes in the vegetation cover. Grass-sedge-herb tundra dominated the area pointing to significant climate deterioration.

Forest-tundra with larches, dwarf birches and willows dominate the lake vicinity at the onset of the Holocene between ca. 11 200 and 9100 cal. yr BP. Shrub alder stands might grow at the low elevations around the lake during the Holocene, between ca. 11 200 and 3500 cal. yr BP. Later, similar-to-modern herb tundra communities dominated the Elgygytgyn Impact Crater.

Acknowledgements. Funding for this research was provided by the ICDP, the U.S. National Science Foundation (NSF), the German Federal Ministry of Education and Research (BMBF), the Deutsche Forschungsgemeinschaft (DFG), the Alfred Wegener Institute for Polar and Marine Research (AWI) and GeoForschungsZentrum Potsdam (GFZ), the Russian Academy of Sciences Far East Branch (RAS FEB), the Russian Foundation for Basic Research (RFBR), and the Austrian Federal Ministry of Science and Research (BMWF).

References

- Anderson, P. M. and Lozhkin, A. V.: Palynological and radiocarbon data from Quaternary deposits of Northeastern Siberia, in: Quaternary vegetation and climate of Siberia and Russian Far East: a Palynological and Radiocarbon Database, edited by: Anderson, P. M. and Lozhkin, A. V., NOAA Paleoclimatology and North-East Science Center, Magadan, 27–34, 2002 (in Russian and English).
- Anderson, P. M., Lozhkin, A. V., and Brubaker, L. B.: Implications of a 24 000-yr palynological record for a Younger Dryas cooling and for boreal forest development in Northeastern Siberia, *Quaternary Res.*, 57, 325–333, 2002.
- Andreev, A. A., Grosse, G., Schirmermeister, L., Kuznetsova, T. V., Kuzmina, S. A., Bobrov, A. A., Tarasov, P. E., Novenko, E. Yu., Meyer, H., Derevyagin, A. Yu., Kienast, F., Bryantseva, A., and Kunitsky, V. V.: Weichselian and Holocene paleoenvironmental history of the Bol'shoy Lyakhovsky Island, New Siberian Archipelago, Arctic Siberia, *Boreas*, 38, 72–110, 2009.

1428

- Belikovich, A. V.: Permanent monitoring in the El'gygytgyn Lake area, IBPS FEB RAS, Magadan, 1988 (in Russian).
- Belikovich, A. V.: Graf method by analyses of habitats in a landscape unit using environment characteristics (El'gygytgyn Lake, Central Chukotka as a model), in: Graf Method in Ecology, edited by: Galanin, A. V., FEB RAS, Vladivostok, 104–111, 1989 (in Russian).
- 5 Belikovich, A. V.: Recreation sources of planned "El'gygytgyn Lake Park", Vestnik FEB RAS, 3, 57–63, 1994 (in Russian).
- Belikovich, A. V. and Galanin, A. V.: "El'gygytgyn Lake" Reservation (Central Chukotka). Vestnik FEB RAS, 4, 22–24, 1994 (in Russian).
- 10 Belyi, F. V.: Impactogenesis and volcanism of El'gygytgyn depression, Petrology, 6, 86–99, 1998.
- Berglund, B. E. and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams, in: Handbook of Holocene Palaeoecology and Palaeohydrology, edited by: Berglund, B. E., Wiley, Chichester, 455–484, 1986.
- 15 Binney, H. A., Willis, K. J., Edwards, M. E., Bhagwat, S. A., Anderson, P., Andreev, A. A., Blaauw, M., Damblon, F., Haesaerts, P., Kienast, F. W., Kremenetski, K. V., Krivonogov, S. K., Lozhkin, A. V., MacDonald, G. M., Novenko, E. Y., Oksanen, P., Sapelko, T. V., Valiranta, M., and Vazhenina, L.: The distribution of late-Quaternary woody taxa in Eurasia: evidence from a new macrofossil database, Quaternary Sci. Rev., 28, 2445–2464, 2009.
- 20 Brigham-Grette, J., Melles, M., Minyuk, P., and Scientific Party: Overview and significance of a 250-ka paleoclimate record from El'gygytgyn Crater Lake, NE Russia, J. Paleolim., 37, 1–16, 2007.
- Blunier, T. and Brook, E. J.: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period, Science, 291, 109–112, 2001.
- 25 Bobrov, A. E., Kupriyanova, L. A., Litvintseva, M. V., and Tarasevich, V. F.: Spores and pollen of gymnosperms from the flora of the European part of the USSR, Nauka, Leningrad, 1983 (in Russian).
- Brubaker, L. B., Anderson, P. M., Edwards, M. E., and Lozhkin, A. V.: Beringia as a glacial refugium for boreal trees and shrubs: new perspectives from mapped pollen data, J. Biogeogr., 32, 833–848, 2005.
- 30 CAVM-Team: Circumpolar Arctic Vegetation Map. Conservation of Arctic Flora and Fauna (CAFF) Map No. 1, US Fish and Wildlife Service, Anchorage, Alaska, 2003.

- Chardez, D.: Ecologie generale des Thecamoebiens (Rhizopoda, Testacea), Bull. Inst. Agr. Stat. Rech. Gembloux, 2, 306–341, 1965.
- Egorova, T. V.: The sedges (*Carex* L.) of Russia and adjacent states, St. Petersburg State Chemical-Pharmaceutical Academy, St. Petersburg, 1999.
- 5 Fradkina, A. F., Alekseev, M. N., Andreev, A. A., and Klimanov, V. A.: East Siberia, Cenozoic climatic and environmental changes in Russia, Geol. Soc. Am. Spec. Paper, 382, 89–103, 2005a.
- Fradkina, A. F., Grinenko, O. V., Laukhin, S. A., Nechaev, V. P., Andreev, A. A., and Klimanov, V. A.: North-Eastern Asia, Cenozoic climatic and environmental changes in Russia, Geol. Soc. Am. Spec. Paper, 382, 105–120, 2005b.
- 10 Glushkova, O. Y. and Smirnov, V. N.: Pliocene to Holocene geomorphic evolution and paleogeography of the El'gygytgyn Lake region, NE Russia, J. Paleolim., 37, 37–47, 2007.
- Glushkova, O. Y., Smirnov, V. N., Matrosova, T. V., Vazhenina, L. N., and Braun, T. A.: Climatic-stratigraphic characteristic and radiocarbon dates of terrace complex in El'gygytgyn Lake basin, Vestnik FEB RAS, 2, 31–43, 2009 (in Russian).
- 15 Grimm, E. C.: TGView, Illinois State Museum, Research and Collections Center, Springfield, 2004.
- Gurov, E. P. and Gurova, E. P.: Stages of shock metamorphism of silica volcanic rocks in the El'gygytgyn meteorite Crater, Chukotka, Dokl. Akad. Nauk SSSR, 249, 121–123, 1979 (in Russian).
- 20 Johnsen, S. J., Dahl-Jensen, D., Dansgaard, W., and Gundestrup, N.: Greenland palaeotemperatures derived from GRIP borehole temperature and ice core isotope profiles, Tellus B, 47, 624–629, 1995.
- Kokorowski, H. D., Anderson, P. M., Sletten, R. S., Lozhkin, A. V., and Brown, T. A.: Late Glacial and early Holocene climatic changes based on a multiproxy lacustrine sediment records from Northeast Siberia, Arct. Antarct. Alp. Res., 40, 497–505, 2008.
- 25 Kozhevnikov, Yu. P.: Vascular plants around El'gygytgyn Lake, in: Natural Depression El'gygytgyn Lake (Problems of Study and Preservation), NEISRI FEB RAS, Magadan, 62–82, 1993 (in Russian).
- 30 Kupriyanova, L. A. and Alyoshina, L. A.: Pollen and Spores of Plants from the Flora of European Part of USSR, Vol. I, Academy of Sciences USSR, Komarov Botanical Institute, Leningrad, 1972 (in Russian).

- Kupriyanova, L. A. and Alyoshina, L. A.: Pollen and Spores of Plants from the Flora of European Part of USSR, Academy of Sciences USSR, Komarov Botanical Institute, Leningrad, 1978 (in Russian).
- 5 Layer, P.: Argon-40/argon-39 age of the Elgygytyn impact event, Chukotka, Russia, *Meteor. Planet. Sci.*, 35, 591–599, 2000.
- Lozhkin, A. V. and Anderson, P. M.: The Last Interglaciation in Northeast Siberia, *Quaternary Res.*, 43, 147–158, 1995.
- Lozhkin, A. V. and Anderson, P. M.: A reconstruction of the climate and vegetation of North-eastern Siberia based on lake sediments, *Paleontol. J.*, 40, 622–628, 2006.
- 10 Lozhkin, A. V., Anderson, P. M., Matrosova, T. V., and Minyuk, P. S.: The pollen record from El'gygytyn Lake: implications for vegetation and climate histories of Northern Chukotka since the late middle Pleistocene, *J. Paleolim.*, 37, 135–153, 2007.
- Lozhkin, A. V., Anderson, P. M., and Vazhenina, L. N.: Younger Dryas and early Holocene peats from Northern Far East Russia, *Quatern. Int.*, 237, 54–64, 2011.
- 15 MacDonald, G. M., Velichko, A. A., Kremenetski, C. V., Borisova, O. K., Goleva, A. A., Andreev, A. A., Cwynar, L. C., Riding, R. T., Forman, S. L., Edwards, T. W. D., Aravena, R., Hammarlund, D., Szeicz, J. M., and Gataullin, V. N.: Holocene treeline history and climate change across Northern Eurasia, *Quaternary Res.*, 53, 302–311, 2000.
- Matrosova, T. V.: Modern spore-pollen spectra of Anadyr Plateau (El'gygytyn Lake), in: *Geology, Geography and Biodiversity of North-East Russia*, Materials of Far-East Regional Conference in Memory of A. P. Vas'kovskiy, edited by: Chereshnev, I. A., NEISRI FEB RAS, Magadan, 159–162, 2006 (in Russian).
- 20 Matrosova, T. V.: Vegetation and climate change in Northern Chukotka during the last 350 ka (basing on lacustrine pollen records from El'gygytyn Lake, *Vestnik FEB RAS*, 2, 23–30, 2009 (in Russian).
- 25 Matrosova, T. V., Anderson, P. M., Lozhkin, A. V., and Minyuk, P. S.: Climate history of Chukotka during the last 300 000 years from the Lake El'gygytyn pollen record, in: *Climate Records from Quaternary Sediments of Beringia*, edited by: Lozhkin, A. V., NEISRI FEB RAS, Magadan, 26–42, 2004 (in Russian).
- 30 Melles, M., Brigham-Grette, J., Glushkova, O., Minyuk, P., Nowaczyk, N. R., and Hubberten, H.-W.: Sedimentary geochemistry of a pilot core from Elgygytyn Lake – a sensitive record of climate variability in the East Siberian Arctic during the past three climate cycles, *J. Paleolim.*, 37, 89–104, 2007.

1431

- Melles, M., Brigham-Grette, J., Minyuk, P., Koeberl, C., Andreev, A., Cook, T., Fedorov, G., Gehhardt, C., Haltia-Hovi, E., Kukkonen, M., Nowaczyk, N., Schwamborn, G., Wennrich, V., and the El'gygytyn Scientific Party: The Lake El'gygytyn Scientific Drilling Project – conquering Arctic challenges in continental drilling, *Sci. Drill.*, 11, 29–40, 2011.
- 5 Nolan, M. and Brigham-Grette, J.: Basic hydrology, limnology, and meteorology of modern El'gygytyn Lake, Siberia, *J. Paleolim.*, 37, 17–35, 2007.
- Reille, M.: Pollen et spores d'Europe et d'Afrique du nord, *Laboratoire de Botanique Historique et Palynologie*, Marseille, 1992.
- Reille, M.: Pollen et spores d'Europe et d'Afrique du nord, supplement 1. *Laboratoire de Botanique Historique et Palynologie*, Marseille, 1995.
- 10 Reille, M.: Pollen et spores d'Europe et d'Afrique du nord, supplement 2. *Laboratoire de Botanique Historique et Palynologie*, Marseille, 1998.
- Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E., Burr, G. S., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., McCormac, F. G., Manning, S. W., Reimer, R. W., Richards, D. A., Southon, J. R., Talamo, S., Turney, C. S. M., van der Plicht, J., and Weyhenmeyer, C. E.: IntCal09 and Marine09 radiocarbon age calibration curves, 0–50,000 years cal BP, *Radiocarbon*, 51, 1111–1150, 2009.
- 15 Schwamborn, G., Meyer, H., Fedorov, G., Schirrmeister, L., and Hubberten, H.-W.: Ground ice and slope sediments archiving Late Quaternary paleoenvironment and paleoclimate signals at the margins of Elgygytyn Impact Crater, NE Siberia, *Quaternary Res.*, 66, 259–272, 2006.
- 20 Schwamborn, G., Förster, A., Diekmann, B., Schirrmeister, L., and Fedorov, G.: Mid to late Quaternary cryogenic weathering conditions in Chukotka, Northeastern Russia: Inference from mineralogical and microtextural properties of the El'gygytyn Crater Lake sediment record, in: *Ninth International Conference on Permafrost*, Institute of Northern Engineering, University of Alaska Fairbanks, 1601–1606, 2008a.
- Schwamborn, G., Fedorov, G., Schirrmeister, L., Meyer, H., and Hubberten, H.-W.: Periglacial sediment variations controlled by late Quaternary climate and lake level change at El'gygytyn Crater, Arctic Siberia, *Boreas*, 37, 55–65, 2008b.
- 25 Shilo, N. A., Lozhkin, A. V., Anderson, P. M., Belaya, B. V., Stetsenko, T. V., Glushkova, O.Yu., Brigham-Grette, J., Melles, M., Minyuk, P. S., Nowaczyk, N., and Forman, S.: First continuous

1432

- pollen record of vegetation and climate change in Beringia during the last 300 thousand years, Dokl. Akad. Nauk, 376, 231–234, 2001 (in Russian).
- Shilo, N. A., Lozhkin, A. V., Anderson, P. M., Brown, T. A., Matrosova, T. V., and Kotov, A. T.: Radiocarbon dates and palynological characteristics of sediments in Lake Melkoe, Anadyr River Basin, Chukotka, Dokl. Akad. Nauk, 407, 235–237, 2006.
- Shilo, N. A., Lozhkin, A. V., Anderson, P. M., Brown, T. A., Pakhomov, A. Yu., and Solomatina, T. B.: Glacial refugium of *Pinus pumila* (Pall.) Regel in Northeastern Siberia, Dokl. Akad. Nauk, 412, 401–403, 2007.
- Shilo, N. A., Lozhkin, A. V., Anderson, P. M., Vazhenina, L. N., Stetsenko, T. V., Glushkova, O. Yu., and Matrosova, T. V.: First data on the expansion of *Larix gmelinii* (Rupr.) Rupr. into arctic regions of Beringia during the early Holocene, Dokl. Akad. Nauk, 423, 680–682, 2008.
- Stockmarr, J.: Tablets with spores used in absolute pollen analysis, Pollen Spores, 13, 614–621, 1971.
- Stuiver, M., Grootes, P. M., and Braziunas, T. F.: The GISP2 $\delta^{18}\text{O}$ climate record of the past 16 500 years and the role of the sun, ocean, and volcanoes, Quaternary Res., 44, 341–354, 1995.
- Van Geel, B.: Non-pollen palynomorphs, in: Tracking Environmental Change Using Lake Sediments. Vol. 3: Terrestrial, Algal and Silicaceous Indicators, edited by: Smol, J. P., Birks, H. J. B., Last, W. M., Bradley, R. S., and Alverson, K., Kluwer, Dordrecht, 99–119, 2001.
- Yurtsev, B. A.: Botanical-geographical zonation and flora regionalization tundra on Chukchi Peninsula, Bot. Zh., 58, 945–964, 1973 (in Russian).

1433

Table 1. Radiocarbon and calibrated ages enclose the two-sigma range of highest probability. The ages have been calibrated using CALIB Rev 6.1.0. (Reimer et al., 2009).

Depth (cm), core	Dated material	^{14}C ages (yr BP)	Calibrated age intervals (cal. yr BP)	Lab. number	Reference
20, P1	Plant remains	3000 ± 30	3078–3268	KIA25979	Schwamborn et al. (2006)
43, P1	Plant remains	3095 ± 45	3209–3403	KIA25980	Schwamborn et al. (2006)
114, P1	Plant remains	3670 ± 30	3906–4087	KIA23976	Schwamborn et al. (2006)
150, P1	Plant remains	3665 ± 35	3890–4090	KIA25981	Schwamborn et al. (2006)
207, P1	Plant remains	8145 ± 45*		KIA28241	Schwamborn et al. (2006)
233, P1	Plant remains	5585 ± 40	4493–6447	KIA23977	Schwamborn et al. (2006)
265, P1	Plant remains	8760 ± 45	9558–9914	KIA23978	Schwamborn et al. (2006)
292, P1	Plant remains	8830 ± 55	9695–10,159	KIA23979	Schwamborn et al. (2006)
314, P1	Plant remains	8885 ± 40	9887–10,182	KIA24865	Schwamborn et al. (2006)
325, P1	Plant remains	8920 ± 110	9660–10,249	KIA28242	Schwamborn et al. (2006)
463, P1	Plant remains	11 160 ± 70	12 801–13 243	KIA23980	Schwamborn et al. (2006)
46, P2	Grass remains	1675 ± 25	1526–1626	KIA24866	Schwamborn et al. (2008)
52, P2	Grass remains	3365 ± 35	3553–3692	KIA27258	Schwamborn et al. (2008)
95, P2	Grass remains	4400 ± 110	4812–5320	KIA27259	Schwamborn et al. (2008)
119, P2	Grass remains	5350 ± 45	5998–6218	KIA27260	Schwamborn et al. (2008)
132 P2	Grass remains	6345 ± 35	7171–7330	KIA24867	Schwamborn et al. (2008)
146–151	<i>Larix</i> seeds	9640 ± 60	10 775–11 193	Poz-42874	This study
170–184	Bulk organic	1890 ± 100*		Poz-42875	This study
205, P2	Grass remains	10 450 ± 60	12 116–12 560	KIA24868	Schwamborn et al. (2008)
210, P2	Grass remains	11 180 ± 147	12 706–13 320	KIA28243	Schwamborn et al. (2008)
226, P2	Grass remains	11 790 ± 242	13 113–14 220	KIA28244	Schwamborn et al. (2008)
0–40, 5011-3	Plant remains	Modern		Poz-33404	This study
40–50, 5011-3	Plant remains	Modern		Poz-33406	This study
50–60, 5011-3	Plant remains	Modern		Poz-33407	This study
60–70, 5011-3	Plant remains	Modern		Poz-33408	This study
70–100, 5011-3	Plant remains	Modern		Poz-33409	This study
100–110, 5011-3	Plant remains	Modern		Poz-33410	This study
173–183, 5011-3	Bulk organic	27 690 ± 200*		Poz-35975	This study
208–230, 5011-3	Bulk organic	20 860 ± 170*		Poz-35977	This study
315–325, 5011-3	Bulk organic	18 800 ± 120*		Poz-35978	This study
395–400, 5011-3	Bulk organic	24 070 ± 320*		Poz-35979	This study
845–852, 5011-3	Bulk organic	24 590 ± 220*		Poz-35980	This study
899–910, 5011-3	Bulk organic	28 440 ± 320*		Poz-35981	This study

The obviously inversed ages were rejected and marked with *.

1434

Table 2. List of testate amoebae species found in the P1 core sediments. Ecological preferences are according Chardez (1965): Sh – *Sphagnum*, m – green moss, s – soils, w – water.

species	depth, cm/ ecology	114-120	187-200	207-212	212-217	217-223	223-227	227-233	233-236	236-242	242-252	252-262	262-268	268-271	271-274	274-277	277-283	283-288	288-292	292-296	296-314	314-325	325-334	463-473
<i>Arcella arenaria</i>	m												1	2	3									
<i>A. arenaria</i> v. <i>compressa</i>	m																							
<i>A. discoides</i>	w						2																	
<i>Bullinularia indica</i>	ms												1											
<i>Centropyxis aculeata</i>	w								2					3	2	3			10		2			
<i>C. aerophila</i>	m	18	4	3		13	19		3	10			14	7	4		5	15	34	19	62	4	2	
<i>C. aerophila</i> v. <i>grandis</i>	w												1					2						
<i>C. aerophila</i> v. <i>sphagnicola</i>	Shm			2									7					2	6					
<i>C. cassis</i>	shSwm						1							2										
<i>C. cassis</i> v. <i>spinifera</i>	wm																							
<i>C. constricta</i>	ws	8	4	3	2	1	8	6	3	3		2	11	25	9	6	13	26	87	11	4	5	1	
<i>C. constricta</i> v. <i>minima</i>	w			5		11						2									1			
<i>C. discoides</i>	w												4											
<i>C. ecornis</i>	w			2		3				10		3	16	2	1	10		3	38	2				
<i>C. ecornis</i> (sensu Ogden, Hedley, 1960)	w																	1						
<i>C. elongata</i>	ms					2																		
<i>C. gibba</i>	wm								2			1				1						1		
<i>C. orbicularis</i>	wShm												2											1
<i>C. plagiostoma</i>	s					7								3				3	9	4				
<i>C. plagiostoma</i> f. <i>A. (longa)</i>	s								2				1	1	1									
<i>C. platystoma</i>	swm		3	3		2	4	2					3	7			3	6	2	44	30		2	
<i>C. platystoma</i>	wShm	54	2			9	10	2	3				16	11			20	13	29	5	31	48	3	
<i>C. sylvatica</i> v. <i>microstoma</i>	s																							
<i>Cylopyxis arceioides</i>	Shwm																							
<i>C. eurystoma</i>	wSh													1	10									
<i>C. eurystoma</i> v. <i>parvula</i>	s								1															
<i>C. kahli</i>	wm		1	3		2				2								7		9	3		1	
<i>Plagiopyxis bathystoma</i>	s		5	2									2											
<i>P. calida</i>	wShms											5											4	
<i>P. declivis</i>	Shs		1																				5	
<i>Heleopera petricola</i>	ws			3										2										
<i>H. petricola</i> v. <i>amethystea</i>	wSh						3							3										2
<i>H. sylvatica</i>	ms													1										
<i>Nebela penardiana</i>	s							1									1							
<i>N. tubulosa</i>	Sh													5										
<i>Diffugia acuminata</i>	w																							
<i>D. globulosa</i>	wSh		7											3			2	3		5				
<i>D. globulosa</i>	wSh		8		7		2	12	8	3			1	5	3	1	32	9	23	19		2	11	
<i>D. lucida</i>	wShm						1							1	1									
<i>D. microstoma</i>	w		1																					
<i>D. minuta</i>	w			2				6														1		
<i>D. penard</i>	w													2										
<i>D. prisis</i>	w																							1
<i>D. mammilaris</i>	w													1										
<i>D. angulostoma</i>	w													3										
<i>D. mica</i>	w																							
<i>D. oblonga</i> v. <i>angusticollis</i>	w																							
<i>D. oblonga</i> v. <i>venusta</i>	w																							
<i>D. gassoweki</i>	w							2																
<i>Lagenodiffugia vas</i>	w														1	1								

1435




Fig. 1. Location map of the study sites and mentioned cores and sections. OC – Olga Creek terrace section from Enmyvaam River valley (Glushkova and Smirnov, 2007; Shilo et al., 2008; Glushkova et al., 2009).

1436




Fig. 2. Lithological, geochronological, grain size and TOC data from P1, P2, and 5011-3 cores.

1437




Fig. 3. Percentage pollen diagram of core P1.

1438




Fig. 4. Percentage pollen diagram of core P2.




Fig. 5. Percentage pollen diagram of core 5011-3.



Fig. 6. Seeds of *Larix* found in core P2.