

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Modelling snow accumulation on Greenland in Eemian, glacial inception and modern climates in a GCM

H. J. Punge¹, H. Gallée², M. Kageyama¹, and G. Krinner²

¹Laboratoire des sciences du climat et de l'environnement (LSCE)/IPSL, CEA-CNRS-UVSQ, UMR8212, 91191 Gif-sur-Yvette, France

²UJF Grenoble 1/CNRS, Laboratoire de Glaciologie et Géophysique de l'Environnement (LGGE) UMR5183, Grenoble, 38041, France

Received: 11 April 2012 – Accepted: 16 April 2012 – Published: 25 April 2012

Correspondence to: H. J. Punge (heinz-jurgen.punge@lsce.ipsl.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Changing climate conditions on Greenland influence the snow accumulation rate and surface mass balance (SMB) on the ice sheet and, ultimately, its shape. This can in turn affect local climate via orography and albedo variations and, potentially, remote areas via changes in ocean circulation triggered by melt water or calving from the ice sheet. Examining these issues in the IPSL global model requires improving the representation of snow at the ice sheet surface. In this paper, we present the new snow scheme implemented in LMDZ, the atmospheric component of the IPSL coupled model. We analyze surface climate and SMB on the Greenland ice sheet under insolation and oceanic boundary conditions for modern, but also for two different past climates, the last glacial inception (115 kyr BP) and the Eemian (126 kyr BP). While being limited by the low resolution of the GCM, present-day SMB is on the same order of magnitude as recent regional model findings. It is affected by a moist bias of the GCM in Western Greenland and a dry bias in the north-east. Under Eemian conditions, the SMB diminishes largely, and melting affects areas with today high surface altitude including recent ice core drilling sites as NEEM. In contrast, glacial inception conditions lead to a higher mass balance overall due to the reduced melting in the colder summer climate. Compared to the widely applied positive degree day (PDD) parameterization of SMB, our direct modelling results suggest a weaker sensitivity of SMB to changing climatic forcing. In addition, significant differences in surface climate and SMB are found between simulations using monthly climatological mean and actual interannually varying monthly mean forcings for the ocean surface temperature and sea ice cover, in particular for the Eemian.

1 Introduction

Among the conditions determining the long term evolution of an ice sheet, the amount of snow and ice accumulated on its top is of primary importance. Together with ablation

CPD

8, 1523–1565, 2012

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 processes at the ice sheet margins and bottom, the surface mass balance (SMB) con-
strains the volume of the ice sheet and thus, on a large scale, impacts the evolution
of the sea level. It is most directly affected by perturbations of the surface climate and
hence the main contributor to the observed ice sheet response to recent anthropogenic
warming (Chen et al., 2006; Mote, 2007; Fettweis et al., 2011). The representations of
snow accumulated on the earth's surface used in numerical models are various and
historically depend on the intended use of the model. For general circulation and re-
gional climate models, a wide range of schemes exists (Slater et al., 2001) from simple
bucket models to sophisticated multi-layer models (e.g. Manabe, 1969; Lynch-Stieglitz,
10 1994; Douville et al., 1995), while models of ice sheet dynamics directed towards long
term ice sheet evolution frequently use simple parameterisations to estimate the SMB
based on mean temperature and precipitation, such as the positive degree-day (PDD)
method (Braithwaite, 1984; Reeh, 1991).

As snow transformation processes can have considerable effects on albedo and thus
15 the radiation budget for all surface types (Brun et al., 1992; Gallée and Duynkerke,
1997; Munneke et al., 2011), more complex models with more accurate representation
of surface processes such as melting and refreezing, albedo variation, layering and
percolation are in development and more and more often used in GCMs. The accu-
rate representation of such processes allows for a more physically based estimation
20 of the SMB on ice sheets. Some specific regional climate models have already inte-
grated such schemes and studied the surface mass balance (SMB) of ice sheets and
its variations in modern climate extensively (Box et al., 2006; Fettweis, 2007; Rignot
et al., 2008; Ettema et al., 2009) to estimate ice mass variations that can be com-
pared to those derived from remote sensing observations (Wouters et al., 2008; Rignot
25 et al., 2008; Velicogna, 2009; van den Broeke et al., 2009; Fettweis et al., 2011). In this
context, regional atmospheric models driven by global circulation data based on obser-
vation (re-)analysis are generally used. The use of GCM forcing recently permitted first
use of such a model under past climate conditions (van de Berg et al., 2011).

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the context of climate model development towards earth system models (ESMs), comprehensive process-based representations of snow physics are beginning to find their way into global climate models to allow for realistic water and energy transformation on ice sheets (e.g. Ridley et al., 2005; Vizcaíno et al., 2010). In parallel, efforts have recently been undertaken to couple an ice sheet model with SMB fields from regional climate model (Helsen et al., 2012). Still, important spatial scale differences between atmospheric and ice sheet models can require complex interpolations and a possible alternative consists in the continued use of parameterisations including the PDD method to estimate SMB.

A weakness of the latter approach is that such parameterisations contain constants which are usually calibrated to yield results that match present day observations, typically for Greenland (e.g. van den Broeke et al., 2010), and are thus a priori valid only for this period. van de Berg et al. (2011) indeed find that PDD factors tuned for modern days fail to describe climate under different insolation accurately. They may also be specific for the shapes and locations of the present day ice sheets. However, ice sheet evolution generally occurs on very long time scales, for which observations are rarely available, and ice sheets take different shapes and appear in different places than at present. With climate changing in function of orbital parameters, phenomena like the daily and seasonal cycle change as well, and hence, possibly, the relationships between temperature and precipitation and the SMB.

Past climate states present an extensive test field to climate and ice sheet models. The Eemian interglacial around 125 kyr BP with its higher boreal summer insolation is an example for a warmer than present climate state (e.g. Huybrechts, 2002). Despite the different origin of the climate change, it is sometimes considered an analogue to anthropogenically caused warmer climate in the future with associated shrinking of the ice sheets (Overpeck et al., 2006; Kopp et al., 2009; van de Berg et al., 2011). The following colder period, the last glacial inception, can serve as an example for a cool climate leading to ice sheet growth (Vettoretti and Peltier, 2003; Calov et al., 2005; Khodri et al., 2005).

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the past, in paleo climate studies focussing on ice sheets, Earth system models of intermediate complexity (EMICs) are generally used. They are characterized by a rather coarse representation of the atmosphere, frequently coupled to finer resolution ice sheet models to study the ice sheets evolution. In these models, the SMB is frequently estimated by the PDD method (e.g. Goosse et al., 2010; Bonelli et al., 2009), while sometimes approaches based on energy balance are used (e.g. Ganopolski et al., 2010; Gallée et al., 1991).

A few recent studies used surface energy balance schemes with albedo parameterizations depending on surface conditions such as snow depth or surface temperature in combination with downscaled EMIC results for coupling of ice sheet and climate models to study the Eemian (Robinson et al., 2011; Fyke et al., 2011). Robinson et al. (2011) obtain constraints on model parameters based on Eemian ice sheet loss, and found the melt rate parameter to be decisive for the shape of the Greenland ice sheet (GIS). Fyke et al. (2011) find large impacts of the use of interactive or constant albedo and meltwater refreezing on the shape of the ice sheet in the Eemian and late Holocene.

In a general circulation model setup with a snow model including interactive albedo, Otto-Bliesner et al. (2006) found significantly lower than present day SMB for the beginning of the Eemian interglacial, important mass losses in an ice sheet model forced by these results, and a sea level rise of 2.2 to 3.4 m. However, sea levels rise estimates from direct observations are significantly higher at 6–9 m (Kopp et al., 2009) – an example for many open questions in the context of past ice sheet evolution.

In a novel approach, van de Berg et al. (2011) use a regional model nested into a GCM and show that the change of melt for the Eemian calculated by a PDD method appears to be 30 % lower than the one in their model, questioning the validity of the PDD approach for past climate, as conclude Ganopolski and Robinson (2011).

We complement these studies by the following approach: a comprehensive snow scheme already in use for regional studies (Gallée et al., 2001; Fettweis, 2007; Fettweis et al., 2011) is integrated into the global atmospheric circulation model LMDZ. This allows to study accumulation and SMB of ice sheets under paleo climate conditions. In

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 addition to the preindustrial climate state, we test our model for two paleo periods with contrasted climates, the Eemian, around 126 kyr BP, and the following glacial inception, around 115 kyr BP. Earth's orbital parameters during these periods leads to increased (decreased) summer insolation in the Northern Hemisphere during the Eemian (last glacial inception) (Braconnot et al., 2008). As this modified insolation is the principal cause of ice sheet melt in the Eemian (van de Berg et al., 2011; Ganopolski and Robin-

10 son, 2011) and growth at inception, reproducing the SMB variations in these climates is a suitable test case for models. Directly modelled SMB fields from GCMs may not be sufficient in terms of resolution to drive ice sheet models, but this study is seen as a first step in the analysis of the impact of this new snow representation on the climates simulated by the GCM. We do not focus on the evolution of the ice sheets, nor their interactions with the oceans, but perform time slice simulations and analyse the sensitivity of the SMB to surface climate conditions with the atmospheric component of the IPSL coupled GCM.

15 After describing the model versions used and the simulations performed in Sect. 2, we evaluate the effect of using an improved versus a simplistic snow scheme in the LMDZ GCM (Sect. 3.1) and compare the SMB obtained by the model to the SMB estimated with a PDD method (Sect. 3.2) for modern and paleo climates, respectively. For all experiments, coupled simulations including the same atmospheric model provide the lower boundary conditions (LBC). This also allows us to investigate the impact of using climatological mean versus variable LBCs in Sect. 3.3 before drawing conclusions in Sect. 4.

2 Tools and methodology

2.1 The atmosphere general circulation model LMDZ and improved snow representation on ice sheets

The LMDZ atmospheric general circulation model is used in its version 4 (Hourdin et al., 2006), as currently used in the coupled atmosphere-ocean model IPSL-CM5 for the IPCC's 5th assessment report. We use a moderate resolution 96×95 grid ($3.75^\circ \times 1.875^\circ$), with 19 layers in the vertical.

The model distinguishes 4 different surface types: "Land", "Land ice", "Ocean" and "Sea ice". At the current stage, the repartition of these types is not adjusted during the run.

The introduction of the snow model just concerns the "Land ice" component. Therefore we limit our analysis essentially to the Greenland ice sheet, where recent ice core drillings (NEEM, NGRIP) promise new insights on the Eemian and past-Eemian time periods and have already produced numerous findings on properties of ice and snow on ice sheets (e.g. Steen-Larsen et al., 2011; Kuramoto et al., 2011). In the standard setup, the model comprises a bulk snow layer for the "Land ice" surface, with melting in the case of surface temperatures above the freezing point, $T_{\text{surf}} > T_0$, at a rate proportional to $T_{\text{surf}} - T_0$. No transformation to ice is foreseen and runoff is immediately transferred to the ocean in the coupled model version. Snow albedo is constant at 0.77 over land ice, and for both of the two wavelength intervals in LMDZ corresponding to visible and infrared light.

The new snow model is based on the CROCUS scheme (Brun et al., 1989, 1992). It is used here in the framework of the snow and ice part of the Soil Ice Snow Vegetation Atmosphere Transfer scheme (SISVAT) (Gallée and Duynkerke, 1997; Lefebvre et al., 2002, 2003) as applied in the regional model MAR (Fettweis, 2007; Lefebvre et al., 2005). The model includes the automatic creation and merging of up to 35 snow layers each with its own properties and with evolving thickness. In particular, the snow type is described by the three variables dendricity, sphericity and grain size. The

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



transformation from fresh snow, which is dendritic, towards more or less compact older snow, occurs as a function of environmental conditions such as temperature and the vertical temperature gradient. It will typically lead to small-grained, non-spheric snow under cold conditions but larger, more spheric grains when temperature approaches the freezing point. Melt and refreezing with ice formation are also represented. Optical properties are adjusted as a function of snow properties, permitting accurate modelling of snow albedo and mass balance (Gallée and Duynkerke, 1997; Lefebvre et al., 2002, 2003). Albedo is calculated for three wavelength bands independently and then interpolated to the two LMDZ bands. We use the same configuration as Fettweis (2007).

At the interface to ice, at the lower boundary of the snow model, we remove one layer when the total snow load in all layers exceeds 5000 kg m^{-2} in order to allow for long simulations and avoid too thick layers, but supply an additional ice layer when the snow load becomes less than 200 kg m^{-2} , assuming the presence of a sufficiently thick ice layer on all land ice points. This is consistent with the assumption of a constant ice sheet topography but will need to be adjusted when changing surface types will be allowed and, ultimately, an ice sheet model is coupled to the ESM. The choice of the threshold is a compromise between the need to avoid the disappearance of all snow model layers or excessive snow or ice load on the one hand and minimum manipulation of the snow models layering scheme.

2.2 Setup of numerical experiments

To examine the impact of different climate conditions on the Greenland SMB, we choose to investigate Eemian and inception climates. These mainly differ from the pre-industrial in their orbital parameters, inducing an amplified seasonal cycle in the Northern Hemisphere for the Eemian but a reduced one for the inception. The same orbital parameters as in the simulations of Braconnot et al. (2008) are prescribed. Because these changes in incoming radiation at the top of the atmosphere have an impact on the atmosphere also via ocean and sea ice feedbacks, we run the atmospheric model with the corresponding sea surface temperatures (SST) and sea ice concentrations (SIC)

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from a 20-yr slice of the coupled atmosphere-ocean-sea ice simulations of Braconnot et al. (2008), interpolated from the slightly different 96×72 grid of their IPSL-CM4 model, as LBC.

We compare simulations for three model setups:

- the original LMDZ4 model (Hourdin et al., 2006) driven by climatological monthly mean LBC computed from the IPSL-CM4 simulations, repeated each year,
- LMDZ+SISVAT driven by the same climatological LBC,
- LMDZ+SISVAT driven by the variable actual monthly mean sea surface temperatures from the IPSL-CM4 simulations.

In the following, we will refer to the first setup as standard LMDZ. The simulations are summarized in Table 1.

To improve the numerical stability of the LMDZ-SISVAT model, it was necessary to improve the vertical resolution near the ground in the simulations with this model. The first of the model's hybrid sigma levels was thereby shifted from $296.5544 \text{ hPa} + 0.9882 p_s$ to $49.8882 \text{ hPa} + 0.9980 p_s$, corresponding approximately to a pressure shift from 1004.3 to 1011.7 hPa at a grid point at sea level, or an altitude shift from 71 to 12 m. In a sense this change anticipates the evolution in GCMs towards higher vertical resolutions near the ground in order to achieve a better accuracy and representation of near-surface processes, following the example of regional models. For LMDZ, a resolution similar to ours has already been used by Krinner et al. (1997), in light of the particular importance in of the boundary layers over ice sheets for turbulent fluxes and katabatic winds. The IPCC-AR5 simulations with the IPSL model are carried out with a lower level at a mean approximate altitude of 33 m. Analysis of results showed that the vertical level change alone leads indeed to significant changes in the modeled atmospheric fields. The additional simulations L_cPIr (standard levels) and LV_cPIr (shifted levels, see Table 1) were thus performed with a slightly more recent version of LMDZ to allow us to separate these changes from the effects of the snow model.

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Present-day surface type repartition and topography are used in all simulations. While this clearly presents a restriction of the model with regard to the accuracy of the results obtained for past periods, our goal here is to test the sensitivity of the new version of this model to past conditions, and equilibration of surface types in the model would happen on time scales much larger than considered in this work. Greenhouse gas concentrations are also the same as at preindustrial times in all simulations.

2.3 Surface mass balance evaluation

LMDZ-SISVAT and standard LMDZ provide total snow amount at the end of each month as output. The monthly SMB in each simulation i , $\Delta h_{\text{SMB, modelled}}^i$ is thus directly accessible as the difference to the previous month.

For the estimation of the annual SMB $\Delta h_{\text{SMB, PDD}}^i$ by the PDD method we use the formulation of Reeh (1991) for the number of N_{PDD}^i of positive degree days and the same parameters as Quiquet et al. (2012):

$$N_{\text{PDD}}^i = \int_{T_0}^{\infty} T \left\{ \int^{t_{\text{Ann}}} \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(T - T^i(t))^2}{(2\sigma)^2}\right) dt \right\} dT \quad (1)$$

where $T^i(t)$ are the climatological monthly mean temperature from January to December near the surface for the respective model and $T_0 = 0^\circ\text{C}$. The constant $\sigma = 5.0^\circ\text{C}$ is supposed to represent the daily cycle and the variability caused by weather in temperature. The SMB is then obtained as

$$\Delta h_{\text{SMB, PDD}}^i = f_{\text{snow}}(T_{\text{Ann}}^i) \cdot P_{\text{Ann}}^i - C_{\text{snow}} \cdot N_{\text{PDD, snow}}^i - C_{\text{ice}} \cdot N_{\text{PDD, ice}}^i \quad (2)$$

where

$$N_{\text{PDD, snow}}^i = \min \left(N_{\text{PDD}}^i, \frac{f_{\text{snow}}(T_{\text{Ann}}^i)P_{\text{Ann}}^i}{C_{\text{snow}}} \right) \quad (3)$$

and

$$N_{\text{PDD, ice}}^i = \max(N_{\text{PDD}}^i - N_{\text{PDD, snow}}^i, 0) \quad (4)$$

5 where T_{Ann}^i and P_{Ann}^i are the annual climatological mean temperature and precipitation, respectively.

The solid fraction of f_{snow} of climatological annual mean precipitation P_{Ann}^i is obtained with the method of Marsiat (1994) as described in Quiquet et al. (2012); $C_{\text{snow}} = 5.0 \text{ mm day}^{-1} \text{ }^\circ\text{C}^{-1}$ and $C_{\text{ice}} = 8.0 \text{ mm day}^{-1} \text{ }^\circ\text{C}^{-1}$ are supposed to represent relation between melt and temperature for snow and ice, respectively. All constants have
10 been chosen by Quiquet et al. (2012) to give a relatively good match to modelled SMB for present day climate.

We follow this approach using the same formulation in order to compare it to the directly modelled SMB.

15 3 Results

3.1 Modelled climate and surface mass balances

3.1.1 Comparison for modern climate

In order to evaluate the mean climate in the simulations, climatologies are computed from 20 yr of the model output fields. An evaluation of modelled temperature in the original LMDZ version and the two simulations with LMDZ-SISVAT for pre-industrial times compared to selected meteorological observations (Cappelen et al., 2011; Steffen and Box, 2001) and present-day simulations of the RACMO2 (Ettema et al., 2010,
20

1958–2007) and MAR (Fettweis et al., 2011, 1980–1999) regional models, which are driven by reanalysis data, is performed in Fig. 1. Data for further stations and details on the comparison are presented in Appendix A. For this model evaluation we assume that the difference resulting from the different time coverage of observation data and among the models, in particular between preindustrial climate and climate at the end of the 20th century is unimportant compared to model biases and inter-model differences.

We find that in almost all cases, the LMDZ-SISVAT simulations are closer to the station data and the regional models than standard LMDZ. The simulations with modified vertical layers have a notable warm bias at Humboldt and the other stations in the northern part of the ice sheet (cf. Fig. A1) during winter, while those with standard levels are biased low at the stations on the ice sheet and the coastal stations in the south. Simulated summer temperature matches better with observations in the run with new snow scheme but not in the run only changing the vertical levels (LVcPI) at all stations in panels 1 and 10 out of 12 in Fig. A1. This improvement can thus be attributed to the change in snow scheme. Overall, we conclude that the simulations with LMDZ-SISVAT are in reasonable agreement with the observed data and the deviations from observations are often of similar magnitude as for the two regional models. The same is generally true for precipitation (cf. Fig. A2).

The main variables determining the SMB are near-surface temperature and precipitation. While the annual mean temperature will affect long term ice sheet dynamics, for SMB mostly the temperature in the summer season (June-July-August, JJA), relatively well represented by LMDZ-SISVAT in the station data, will be decisive, ruling the amount of melt. We thus compare the climatological mean fields of annual mean temperature, JJA temperature and annual mean precipitation in the same five model simulations as above in the form of maps (Fig. 2). Ettema et al. (2009) and Fettweis et al. (2011) performed extensive validation of their models and hence we consider them as references from now on. The results of these two models also reflect the considerable uncertainties that persist, in particular regarding precipitation and SMB (Hanna et al., 2011; Ettema et al., 2009, 2010).

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

On the scale of the Greenland ice sheet, annual mean temperatures in the standard LMDZ version show a distinct cold bias, in particular in the center and at the borders of the southern part of the ice sheet, which disappears in the versions with new snow scheme but also in the one with modified vertical layers. Mean temperatures in the two LMDZ-SISVAT simulations and LVcPIr are similar as in the two regional simulations. All LMDZ simulations show relatively high annual mean temperatures at the northern coast of Greenland.

The picture is similar for boreal summer (JJA) mean temperatures, however the figure confirms that most of the improvement in LScPI and LSiPI compared to LcPI comes from the snow model (LScPI – LVcPI) and not the change of model levels (LVcPI – LcPI), in particular towards the ablation zone. Despite an obvious lack of accuracy near the coasts due to their low spatial resolution, the simulations with LMDZ including the SISVAT snow scheme represent the field quite well.

Resolution-related differences among the annual mean precipitation fields are more pronounced, but the LMDZ simulations globally fall within the range of the regional models. This is confirmed for the total annual mean ice sheet precipitation listed in Table 2. Common to all LMDZ simulations are relatively high precipitation in the north-west of Greenland and low precipitation in the north-east.

Average modelled annual SMB in the simulations with the regional models, and the LMDZ simulations LcPI, LScPI and LSiPI are given in Fig. 3a and can be compared to the ones derived with the PDD method, shown in Fig. 3b. No comparable data is available for LcPIr and LVcPIr, as a snow threshold from the coupled model version is applied in these simulations, above which all snow is melted artificially. While the results obtained by the two methods are very similar for each of the two regional models, clear differences appear in the case of the LMDZ simulations. The latter do not represent adequately the coastal high SMB zone in the south-east, certainly because of the coarse resolution, and generally underestimate SMB in the southern part. The result is more satisfactory for the north. However, the size of the modelled ablation zones with negative SMB, absent in standard LMDZ, is somewhat overestimated in LMDZ-SISVAT,

in particular in the north-east. These factors lead to a rather low estimate of total GIS SMB in LMDZ-SISVAT (Table 2).

The map of SMB estimated by the PDD method, unlike the one of modelled SMB, shows several grid points with values below $-0.7 \text{ m i.e. yr}^{-1}$ at the very margin of the ice sheet. As the grid cells are much larger than the width of the area with this high melt in reality, their contribution to the low estimates of total ice sheet PDD SMB in Table 2 is important.

We conclude that the SMB obtained with LMDZ-SISVAT is on the right order of magnitude, but limited by its coarse resolution, and future higher resolution simulations are required for a meaningful and complete validation, in particular near the borders of the GIS. Direct modelling of the SMB appears to be more accurate than the PDD method in the case of the GCMs, due to resolution-based effects at the coast.

3.1.2 Effect of vertical levels and the new snow scheme

We now compare the standard and improved version simulations with LMDZ for preindustrial climate in more detail in order to evaluate the effects of the model changes. We first evaluate the difference due to the changed model levels in standard LMDZ, LVcPIr – LcPIr, in boreal summer (JJA) and winter (December-January-February, DJF) mean temperatures (Fig. 4a, b, left hand side). These fields then have to be subtracted from the difference between the simulations with new snow scheme and the one with standard LMDZ, LScPI – LcPI, to isolate the effect of the snow scheme (Fig. 4a, b, right hand side).

The change of levels leads to a mean annual warming of about 6 K over the entire ice sheet in winter, and a warming of about 1 K over Greenland, accompanied by a cooling of the same amount over the surrounding oceans, in summer. Similar changes are observed over the other continents and oceans, but a detailed discussion of these global effects is clearly outside the scope of this article. The net effect of the SISVAT snow scheme over Greenland is a cooling of 1–2 K in winter, most pronounced at higher

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



altitudes of the southern half, and a warming of 2–5 K in summer, most pronounced in the north-east.

The modelled surface albedo generally increases with altitude in LMDZ-SISVAT (Fig. 4c, d). The seasonal surface albedo variations can explain the temperature differences noted above: the higher albedo of the fresh snow at the surface in winter (Fig. 4c) leads to a cooling in the south, whereas in the north the effect is absent due to missing sunlight. In contrast, the albedo is lower in summer, except at the highest grid points (Fig. 4d), due to snow transformation and melting, which leads to a higher ratio of absorbed shortwave radiation, and thus a warming. The effect is strongest in the north-east of the ice sheet where precipitation is rare and the age of the surface snow is high.

Cloud cover increases in the model simulations with modified vertical layers, both in winter and summer (Fig. 4e, f), except for the highest altitudes. This has to be attributed to different functioning of the LMDZ cloud physics schemes with the new levels. There is however a distinct impact of the new snow scheme on cloud concentration over Greenland, which roughly correlates with the albedo changes, except for the north-east in summer. Increased cloud cover will tend to limit the increased radiative loss favoured by higher albedo.

As a consequence of the level shift, we find lower annual precipitation in north-west Greenland, and higher precipitation in the south-east and over the Atlantic (Fig. 4g). The net effect on precipitation due to the snow scheme is insignificant over Greenland. The combined effects of level changes and snow model (LScPI – LcPI) on precipitation amount to +33 Gt yr⁻¹ (cf. Table 2). Evaporation (Fig. 4h) increases over the oceans and coastal areas of Greenland as a consequence of the level changes. The introduction of the new snow scheme leads to a decrease of around 0.1 m i.e. yr⁻¹ in the coastal zones, so that the evaporation difference LScPI – LcPI is near zero in the ablation zone, and +16 Gt yr⁻¹ over the entire ice sheet. The precipitation-evaporation (*P-E*) change is hence +17 Gt yr⁻¹, an increase of 3 %, located mostly in the south-east.

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1.3 Combined effect of vertical levels and new snow scheme on Greenland SMB

In order to illustrate the impact of the changes to the LMDZ model on surface mass balance in different climate states, we show the anomaly $LScPI - L_cPI$ for SMB in Fig. 5, as well as the corresponding anomalies for Eemian and glacial inception. Modelled coastal SMB (Fig. 5a) is lower with SISVAT than in standard LMDZ in all climates. The most important contribution to this SMB decrease is the albedo-related warming in the model with new snow scheme. In the south of Greenland, the effect of the warming due to the level changes is of equal magnitude, but at least partially balanced by the increase in precipitation. Despite nearly identical temperature differences of 3–5 K (see Fig. 6) for all climates, the impacts on SMB are much more prominent in Eemian climate and much weaker at glacial inception compared to preindustrial climate: the total ice sheet surface mass balance decreases by 155 Gt for the PI, 578 Gt for the Eemian, and 79 Gt for glacial inception. This reflects the high non-linearity of the temperature-SMB relationship: a roughly similar warming has a much more dramatic effect on an ice sheet in an already warmer climate.

The SMB analyzed by the PDD method (Fig. 5c) shows a similar picture. For the PI climate, the PDD method shows a slightly higher SMB in the south-west, explained by the higher precipitation in $LScPI$, but absent in the modelled SMB. Near the coast, the SMB difference in Eemian climate is more pronounced with the PDD method. The total PDD-estimated ice sheet surface mass balance decreases by 224 Gt for the PI, 774 Gt for the Eemian, and 39 Gt for glacial inception. The difference for inception climate is hence near neutral, despite a similar temperatures increase as in the Eemian simulation. We conclude thus from this subsection that the effect of our model developments on surface mass balance is most pronounced in warm climates and that the spread of these SMB changes among climates estimated with our PDD method is higher than the one obtained by the snow model.

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Climate for Eemian and last inception vs. PI reference

Next, we elaborate the differences of the simulated Eemian and glacial inception climates compared to the preindustrial reference. Figure 6 illustrates the surface climate anomalies for past interglacial and inception simulations compared to the respective PI simulations, in the three model configurations using standard LMDZ and LMDZ-SISVAT with or without interannual variability of the LBC. Differences are shown only where they pass Student's t-test at the 95 % level to assure statistical significance.

The annual mean temperature (Fig. 6a) on Greenland and its surroundings at 126 kyr BP is warmer than in preindustrial times by about 1.8 K in standard LMDZ; the difference is about 1.5 K in the model simulation with updated snow scheme and climatological LBC. Interestingly, the GIS mean warming is only about 1.0 K in the simulation with variable LBC. In the glacial inception climate, while a significant cooling is found on large parts of the GIS in the simulations with climatological LBC, this is not the case in the simulation with variable LBC. We conclude thus that the prescription of climatological LBCs seems to lead to an overestimation of mean near surface temperature variation on Greenland in reaction to past insolation changes.

As the differences in insolation during the two analyzed past epochs concern mostly boreal summer in the Northern Hemisphere (Braconnot et al., 2008), their effect on Greenland temperatures is more pronounced in the JJA seasonal means shown in Fig. 6b. While the tendency in the annual mean is approximately the same over ice sheets and oceans, there is a clear amplification of the temperature differences on the ice sheets in boreal summer, whereas the ocean acts as a heat reservoir, and the increase of its annual cycle amplitude is smaller than over land. Eemian JJA climate is warmer by about 5 K in the center of the GIS in all three model configurations, and inception climate is cooler by about 2.0 K. The difference between simulations with climatological and variable LBC is less pronounced in JJA temperatures than in the annual mean, although the inception cooling is a bit higher in the north in the run with climatological LBC.

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



resulting melt amount variation is much lower, which is most likely again due to the lower temperature.

The SMB variations obtained by the PDD method show some agreement with the directly modelled variations, at least for LMDZ-SISVAT. The PDD method predicts however an increase of SMB at several grid points in the north-west and south-east under Eemian climate due to precipitation increase, absent in LMDZ-SISVAT. Also, near-coastal signals of SMB increase or decrease are amplified by the PDD method.

As a consequence, the total estimated Greenland SMB decrease for the Eemian compared to the PI is relatively high at -229 Gt yr^{-1} for standard LMDZ and -779 Gt yr^{-1} and -673 Gt yr^{-1} for the two LMDZ-SISVAT setups. The modelled total Greenland SMB decrease in these setups are -131 Gt yr^{-1} , -554 Gt yr^{-1} , and -511 Gt yr^{-1} , respectively. The estimates of decrease for LMDZ-SISVAT are still high compared to van de Berg et al. (2011), who find a SMB change of the modelled SMB of -405 Gt yr^{-1} . We conclude that the SMB in LMDZ-SISVAT is likely too sensitive for the Eemian case, in particular when considering that in contrast to this study van de Berg et al. (2011) found lower than modern precipitation in the Eemian, implying an even lower melt increase. Still, the results with LMDZ-SISVAT present an improvement compared to standard LMDZ.

For the inception, the PDD-predicted SMB gains are 81 Gt yr^{-1} , 266 Gt yr^{-1} , and 195 Gt yr^{-1} for the three model versions, respectively, and again much higher than the 36 Gt yr^{-1} , 112 Gt yr^{-1} , and 71 Gt yr^{-1} obtained from the model.

Figure 4 shows the climatological difference between the experiments with LMDZ-SISVAT using climatological and variable sea surface conditions, respectively. Few significant differences are found in temperature and precipitation. However they have visible effects on SMB. In particular, modelled and PDD SMB for 126 kyr BP climate are higher by $0.1\text{--}0.3 \text{ m i.e. yr}^{-1}$ in the south of the ice sheet due to higher precipitation. Slightly higher temperature and lower precipitation in the north-west lead to a slightly lower SMB when using the PDD method. We found thus that the impact of variable LBC on mean climate is not very high in our experiments, but can have significant influence

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on SMB. A more comprehensive study may address whether these signals persist in longer simulations, and if there are similar effects in other models.

4 Conclusions

We evaluated the representation of Greenland surface mass balance in three configurations of the LMDZ model and tested its sensitivity to climate by simulations in paleo climate states. The integration of a physically based snow scheme for ice sheets into a general circulation model improves the modelling of the surface climate and SMB.

A change of the vertical discretization of the model was required to decrease layer thickness near the ground and ensure numerical stability. It lead to additional climatic changes, in particular a warming of the ice sheet in the winter season, and changes in cloud cover and precipitation.

The climatic effects attributed to the new snow scheme include the disappearance of a cold bias of the model in the summer season, which can be explained by the seasonal variations in albedo represented in the new model, and again an alteration of the cloud cover. Overall we find a significant improvement of modelled surface climate compared to the previous model version. In particular the modelled boreal summer near-surface temperature, highly relevant for SMB, is in good agreement with station data.

Interactively modelled albedo also contributes to improving the representation of Greenland climate response to insolation changes in simulations for past climates, as we show for the cases of the Eemian and the last glacial inception. We therefore stress the need to represent physically based snow properties in climate simulations for the past, in particular if insolation conditions deviate significantly from modern day.

The pre-industrial total modelled ice sheet surface mass balance for the Greenland ice sheet is low compared to a range of regional models (see the list in Fettweis, 2007), and in particular in the light of recent studies estimating early 20th century SMB to be higher by 50–200 Gt yr⁻¹ than at the end of the century (Wake et al., 2009; Hanna et al., 2011). This bias is primarily due to low estimates in the south-east, but could

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



possibly be reduced at higher model resolution including better representation of the complex topography of the region; the modelled SMB could then be more suitable for direct use as input to ice sheet models. In the near future resolutions down to 20 km will be achieved in atmospheric GCMs, and hence a representation of physical processes equivalent to present-day regional models can be expected, including feedbacks in the ablation zone such as between katabatic flows and sensible heat flux with potentially high impacts on SMB.

We found that differences in SMB results between PDD estimation and direct modelling mostly concern the margins of the ice sheet, where the PDD modelled SMB changes more drastically with surface climate. The PDD method is more sensitive to the different climatic conditions during the Eemian and the following inception, most prominent in boreal summer temperatures, than direct modelling in terms of total ice sheet SMB. We also note that these climate-induced SMB changes differ most in the south of the ice sheet.

We showed that atmospheric GCM simulations that use a repeated annual cycle of climatological LBC can differ significantly from those using the LBC the climatology is based on in terms of SMB for a time span of 20 yr. These differences are caused by modified mean precipitation and surface temperature differences, linked to the missing representation of extreme temperature or precipitation events in the climatological runs. This effect impacts in particular the reaction of surface climate to insolation change in our simulations. Part of these differences may become insignificant for longer term simulations, as the high variability, especially of precipitation, leads to some uncertainty, but we still strongly recommend the use of variable LBC forcing where available.

Appendix A

Comparison to station data

The model temperatures are interpolated linearly to the station coordinates based on the four nearest grid points. The GC-Net Automatic Weather Stations (AWS) (Steffen et al., 1996) hourly measurements span the period 1995–2008 with differing coverage, and monthly means and standard deviation are given. For the DMI weather stations, we took the average of daily minimum and maximum temperatures since 1877 (Cappelen et al., 2011) to compute the monthly climatology. Note that MAR defines near-surface temperature at the 3 m level while the other models use the 2 m level. Figure A1 shows the temperature comparison as in Fig. 1 for the remaining AWS and near-coastal weather stations. Figure A2 shows the same comparison for precipitation.

Acknowledgements. GC-Net data received from CIRES/University of Colorado, Boulder (K. Steffen). We thank the DMI for providing temperature and precipitation observations for their Greenland weather stations. Janneke Ettema, Jan van Angelen and Michiel van den Broeke (IMAU, Utrecht University) are thanked for providing RACMO2 climate fields. We thank Xavier Fettweis for providing MAR climate data. HJP is supported by the European FP7 project 226520 COMBINE and the ANR project NEEM-France. This work was performed using HPC resources from GENCI-CCRT (Grant t2010012212).



The publication of this article is financed by CNRS-INSU.

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Bonelli, S., Charbit, S., Kageyama, M., Woillez, M.-N., Ramstein, G., Dumas, C., and Quiquet, A.: Investigating the evolution of major Northern Hemisphere ice sheets during the last glacial-interglacial cycle, *Clim. Past*, 5, 329–345, doi:10.5194/cp-5-329-2009, 2009. 1527
- 5 Box, J. E., Bromwich, D. H., Veenhuis, B. A., Bai, L.-S., Stroeve, J. C., Rogers, J. C., Steffen, K., Haran, T., and Wang, S.-H.: Greenland ice sheet surface mass balance variability (1988–2004) from calibrated polar MM5 output, *J. Climate*, 19, 2783–2800, doi:10.1175/JCLI3738.1, 2006. 1525
- 10 Braconnot, P., Marzin, C., Grégoire, L., Mosquet, E., and Marti, O.: Monsoon response to changes in Earth's orbital parameters: comparisons between simulations of the Eemian and of the Holocene, *Clim. Past*, 4, 281–294, doi:10.5194/cp-4-281-2008, 2008. 1528, 1530, 1531, 1539
- Braithwaite, R. J.: Calculation of degree-days for glacier–climate research, *Z. Gletscherkd. Glazialgeol.*, 20, 1–20, 1984. 1525
- 15 Brun, E., Martin, E., Simon, V., Gendre, C., and Coléou, C.: An energy and mass model of snow cover suitable for operational avalanche forecasting, *J. Glaciol.*, 35, 333–342, 1989. 1529
- Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow cover stratigraphy for operational avalanche forecasting, *J. Glaciol.*, 38, 13–22, 1992. 1525, 1529
- 20 Calov, R., Ganopolski, A., Claussen, M., Petoukhov, V., and Greve, R.: Transient simulation of the last glacial inception. Part I: glacial inception as a bifurcation in the climate system, *Clim. Dynam.*, 24, 545–561, doi:10.1007/s00382-005-0007-6, 2005. 1526
- Cappelen, J., Laursen, E. V., Jørgensen, P. V., and Kern-Hansen, C.: DMI Monthly Climate Data Collection 1768–2009, Denmark, The Faroelands and Greenland, DMI Technical Report 10-05, 52 pp., available at: <http://www.dmi.dk/dmi/tr10-05.pdf>, 2010. 1533, 1544
- 25 Chen, J. L., Wilson, C. R., and Tapley, B. D.: Satellite gravity measurements confirm accelerated melting of greenland ice sheet, *Science*, 313, 1958–1960, doi:10.1126/science.1129007, 2006. 1525
- Douville, H., Royer, J. F., and Mahfouf, J. F.: A new snow parameterization for the Météo-France climate model, *Clim. Dynam.*, 12, 21–35, doi:10.1007/BF00208760, 1995. 1525
- 30 Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Bamber, J. L., Box, J. E., and Bales, R. C.: Higher surface mass balance of the Greenland ice

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sheet revealed by high-resolution climate modeling, *Geophys. Res. Lett.*, 36, L12501, doi:10.1029/2009GL038110, 2009. 1525, 1534

Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Box, J. E., and Steffen, K.: Climate of the Greenland ice sheet using a high-resolution climate model – Part 1: Evaluation, *The Cryosphere*, 4, 511–527, doi:10.5194/tc-4-511-2010, 2010. 1533, 1534

Fettweis, X.: Reconstruction of the 1979–2006 Greenland ice sheet surface mass balance using the regional climate model MAR, *The Cryosphere*, 1, 21–40, doi:10.5194/tc-1-21-2007, 2007. 1525, 1527, 1529, 1530, 1542

Fettweis, X., Tedesco, M., van den Broeke, M., and Ettema, J.: Melting trends over the Greenland ice sheet (1958–2009) from spaceborne microwave data and regional climate models, *The Cryosphere*, 5, 359–375, doi:10.5194/tc-5-359-2011, 2011. 1525, 1527, 1534

Fyke, J. G., Weaver, A. J., Pollard, D., Eby, M., Carter, L., and Mackintosh, A.: A new coupled ice sheet/climate model: description and sensitivity to model physics under Eemian, Last Glacial Maximum, late Holocene and modern climate conditions, *Geosci. Model Dev.*, 4, 117–136, doi:10.5194/gmd-4-117-2011, 2011. 1527

Gallée, H. and Duynkerke, P. G.: Air-snow interactions and the surface energy and mass balance over the melting zone of west Greenland during the Greenland Ice Margin Experiment, *J. Geophys. Res.*, 102, 13813–13824, doi:10.1029/96JD03358, 1997. 1525, 1529, 1530

Gallée, H., van Ypersele, J. P., Fichefet, Th., Tricot, Ch., and Berger, A.: Simulation of the last glacial cycle by a coupled, sectorially averaged climate-ice sheet model 1. *The Climate Model*, *J. Geophys. Res.*, 96, 13139–13161, doi:10.1029/91JD00874, 1991. 1527

Gallée, H., Guyomarc'h, G., and Brun, E.: Impact of snow drift on the antarctic ice sheet surface mass balance: possible sensitivity to snow-surface properties, *Bound.-Lay. Meteorol.*, 99, 1–19, doi:10.1023/A:1018776422809, 2001. 1527

Ganopolski, A. and Robinson, A.: Palaeoclimate: the past is not the future, *Nat. Geosci.*, 4, 661–663, doi:10.1038/ngeo1268, 2011. 1527, 1528

Ganopolski, A., Calov, R., and Claussen, M.: Simulation of the last glacial cycle with a coupled climate ice-sheet model of intermediate complexity, *Clim. Past*, 6, 229–244, doi:10.5194/cp-6-229-2010, 2010. 1527

Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A., Seltner, F., Barriat, P.-Y., Campin, J.-M., Deleersnijder, E., Driesschaert, E., Goelzer, H., Janssens, I., Loutre, M.-F., Morales Maqueda, M. A., Opsteegh, T., Mathieu, P.-P., Munhoven, G., Petterson, E. J., Renssen, H., Roche, D. M., Schaeffer, M., Tartinville, B.,

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Timmermann, A., and Weber, S. L.: Description of the Earth system model of intermediate complexity LOVECLIM version 1.2, *Geosci. Model Dev.*, 3, 603–633, doi:10.5194/gmd-3-603-2010, 2010. 1527

Hanna, E., Huybrechts, P., Cappelen, J., Steffen, K., Bales, R. C., Burgess, E., McConnell, J. R., Steffensen, J. P., Van den Broeke, M., Wake, L., Bigg, G., Griffiths, M., and Savas, D.: Greenland ice sheet surface mass balance 1870 to 2010 based on twentieth century reanalysis, and links with global climate forcing, *J. Geophys. Res.*, 116, D24121, doi:10.1029/2011JD016387, 2011. 1534, 1542

Helsen, M. M., van de Wal, R. S. W., van den Broeke, M. R., van de Berg, W. J., and Oerlemans, J.: Coupling of climate models and ice sheet models by surface mass balance gradients: application to the Greenland Ice Sheet, *The Cryosphere*, 6, 255–272, doi:10.5194/tc-6-255-2012, 2012. 1526

Hourdin, F., Musat, I., Bony, S., Braconnot, P., Codron, F., Dufresne, J., Fairhead, L., Filiberti, M., Friedlingstein, P., Grandpeix, J., Krinner, G., Le Van, P., Li, Z.-X., and Lott, F.: The LMDZ4 general circulation model: climate performance and sensitivity to parametrized physics with emphasis on tropical convection, *Clim. Dynam.*, 27, 787–813, doi:10.1007/s00382-006-0158-0, 2006. 1529, 1531

Huybrechts, P.: Sea-level changes at the LGM from ice-dynamic reconstructions of the Greenland and Antarctic ice sheets during the glacial cycles, *Quaternary Sci. Rev.*, 21, 203–231, doi:10.1016/S0277-3791(01)00082-8, 2002. 1526

Khodri, M., Cane, M. A., Kukla, G., Gavin, J., and Braconnot, P.: The impact of precession changes on the Arctic climate during the last interglacial glacial transition, *Earth Planet. Sc. Lett.*, 236, 285–304, doi:10.1016/j.epsl.2005.05.011, 2005. 1526

Kopp, R. E., Simons, F. J., Mitrovica, J. X., Maloof, A. C., and Oppenheimer, M.: Probabilistic assessment of sea level during the last interglacial stage, *Nature*, 462, 863–867, doi:10.1038/nature08686, 2009. 1526, 1527

Krinner, G., Genthon, C., Li, Z.-X., and Le Van, P.: Studies of the Antarctic climate with a stretched-grid general circulation model, *J. Geophys. Res.*, 102, 13731–13745, doi:10.1029/96JD03356, 1997. 1531

Kuramoto, T., Goto-Azuma, K., Hirabayashi, M., Miyake, T., Motoyama, H., Dahl-Jensen, D., and Steffensen, J.: Seasonal variations of snow chemistry at NEEM, Greenland, *Ann. Glaciol.*, 52, 193–200, doi:10.3189/172756411797252365, 2011. 1529

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lefebre, F., Gallée, H., van Ypersele, J., and Huybrechts, P.: Modelling of large-scale melt parameters with a regional climate model in South Greenland during the 1991 melt season, *Ann. Glaciol.*, 35, 391–397, doi:10.3189/172756402781816889, 2002. 1529, 1530
- 5 Lefebre, F., Gallée, H., van Ypersele, J.-P., and Greuell, W.: Modeling of snow and ice melt at ETH Camp (West Greenland): a study of surface albedo, *J. Geophys. Res.*, 108, 4231, doi:10.1029/2001JD001160, 2003. 1529, 1530
- Lefebre, F., Fettweis, X., Gallée, H., van Ypersele, J.-P., Marbaix, P., Greuell, W., and Calanca, P.: Evaluation of a high-resolution regional climate simulation over Greenland, *Clim. Dynam.*, 25, 99–116, doi:10.1007/s00382-005-0005-8, 2005. 1529
- 10 Lynch-Stieglitz, M.: The Development and validation of a simple snow model for the GISS GCM, *J. Climate*, 7, 1842–1855, doi:10.1175/1520-0442(1994)007<1842:TDAVOA>2.0.CO;2, 1994. 1525
- Manabe, S.: Climate and the ocean circulation: I. The atmospheric circulation and the hydrology of the Earth's surface, *Mon. Weather Rev.*, 97, 739–774, 1969. 1525
- 15 Marsiat, I.: Simulation of the Northern Hemisphere continental ice sheets over the last glacial-interglacial cycle: experiments with a latitude-longitude vertically integrated ice sheet model coupled to a zonally averaged climate model, *Paleoclimates*, 1, 59–98, 1994. 1533
- Mote, T. L.: Greenland surface melt trends 1973–2007: evidence of a large increase in 2007, *Geophys. Res. Lett.*, 342, L22507, doi:10.1029/2007GL031976, 2007. 1525
- 20 Munneke, P. K., van den Broeke, M. R., Lenaerts, J. T. M., Flanner, M. G., Gardner, A. S., and van de Berg, W. J.: A new albedo parameterization for use in climate models over the Antarctic ice sheet, *J. Geophys. Res.*, 116, D05114, doi:10.1029/2010JD015113, 2011. 1525
- Otto-Bliesner, B. L., Marshall, S. J., Overpeck, J. T., Miller, G. H., and Hu, A.: Simulating arctic climate warmth and icefield retreat in the last interglaciation, *Science*, 311, 1751–1753, doi:10.1126/science.1120808, 2006. 1527
- 25 Overpeck, J. T., Otto-Bliesner, B. L., Miller, G. H., Muhs, D. R., Alley, R. B., and Kiehl, J. T.: Paleoclimatic evidence for future ice-sheet instability and rapid sea-level rise, *Science*, 311, 1747–1750, doi:10.1126/science.1115159, 2006. 1526
- Quiquet, A., Punge, H. J., Ritz, C., Fettweis, X., Kageyama, M., Krinner, G., Salas y Mélia, D., and Sjolte, J.: Large sensitivity of a Greenland ice sheet model to atmospheric forcing fields, *The Cryosphere Discuss.*, 6, 1037–1083, doi:10.5194/tcd-6-1037-2012, 2012. 1532, 1533
- 30 Reeh, N.: Parameterization of melt rate and surface temperature on the Greenland ice sheet, *Polarforschung*, 59, 113–128, 1991. 1525, 1532

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ridley, J. K., Huybrechts, P., Gregory, J. M., and Lowe, J. A.: Elimination of the Greenland ice sheet in a high CO₂ climate, *J. Climate*, 18, 3409–3427, doi:10.1175/JCLI3482.1, 2005. 1526
- 5 Rignot, E., Box, J. E., Burgess, E., and Hanna, E.: Mass balance of the Greenland ice sheet from 1958 to 2007, *Geophys. Res. Lett.*, 35, L20502, doi:10.1029/2008GL035417, 2008. 1525
- Robinson, A., Calov, R., and Ganopolski, A.: Greenland ice sheet model parameters constrained using simulations of the Eemian Interglacial, *Clim. Past*, 7, 381–396, doi:10.5194/cp-7-381-2011, 2011. 1527
- 10 Slater, A. G., Schlosser, C. A., Desborough, C. E., Pitman, A. J., Henderson-Sellers, A., Robock, A., Vinnikov, K. Y., Mitchell, K., Boone, A., Braden, H., Chen, F., Cox, P. M., de Rosnay, P., Dickinson, R. E., Dai, Y.-J., Duan, Q., Entin, J., Etchevers, P., Gedney, N., Gusev, Y. M., Habets, F., Kim, J., Koren, V., Kowalczyk, E. A., Nasonova, O. N., Noilhan, J., Schaake, S., Shmakin, A. B., Smirnova, T. G., Verseghy, D., Wetzell, P., Xue, Y., Yang, Z.-L., and Zeng, Q.: The Representation of snow in land surface schemes: results from PILPS 2(d), *J. Hydrometeorol.*, 2, 7–25, doi:10.1175/1525-7541(2001)002;0007:TROSIL;2.0.CO;2, 2001. 1525
- 15 Steen-Larsen, H. C., Masson-Delmotte, V., Sjolte, J., Johnsen, S. J., Vinther, B. M., Bréon, F.-M., Clausen, H. B., Dahl-Jensen, D., Falourd, S., Fettweis, X., Gallée, H., Jouzel, J., Kageyama, M., Lerche, H., Minster, B., Picard, G., Punge, H. J., Risi, C., Salas, D., Schwander, J., Steffen, K., Sveinbjörnsdóttir, A. E., Svensson, A., and White, J.: Understanding the climatic signal in the water stable isotope records from the NEEM shallow firn/ice cores in northwest Greenland, *J. Geophys. Res.*, 116, D06108, doi:10.1029/2010JD014311, 2011. 1529
- 20 Steffen, K. and Box, J.: Surface climatology of the Greenland ice sheet: Greenland Climate Network 1995–1999, *J. Geophys. Res.*, 106, 33951–33964, doi:10.1029/2001JD900161, 2001. 1533
- Steffen, K., Box, J., and Abdalati, W.: Greenland Climate Network: GC-Net, in: *CRREL 96–27, Special Report on Glaciers, Ice Sheets and Volcanoes*, tribute to Meier, M. and Colbeck, S. C. (Eds.), 1996. 1544, 1553, 1562
- 30 van de Berg, W. J., van den Broeke, M., Ettema, J., van Meijgaard, E., and Kaspar, F.: Significant contribution of insolation to Eemian melting of the Greenland ice sheet, *Nat. Geosci.*, 4, 679–683, doi:10.1038/ngeo1245, 2011. 1525, 1526, 1527, 1528, 1541

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning recent Greenland mass loss, *Science*, 326, 984–986, doi:10.1126/science.1178176, 2009. 1525
- van den Broeke, M., Bus, C., Ettema, J., and Smeets, P.: Temperature thresholds for degree-day modelling of Greenland ice sheet melt rates, *Geophys. Res. Lett.*, 37, L18501, doi:10.1029/2010GL044123, 2010. 1526
- Velicogna, I.: Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE, *Geophys. Res. Lett.*, 36, L19503, doi:10.1029/2009GL040222, 2009. 1525
- Vettoretti, G. and Peltier, W. R.: Post-Eemian glacial inception. Part I: the impact of summer seasonal temperature bias, *J. Climate*, 16, 889–911, doi:10.1175/1520-0442(2003)016;0889:PEGIPI;2.0.CO;2, 2003. 1526
- Vizcaíno, M., Mikolajewicz, U., Jungclaus, J., and Schurgers, G.: Climate modification by future ice sheet changes and consequences for ice sheet mass balance, *Clim. Dynam.*, 34, 301–324, doi:10.1007/s00382-009-0591-y, 2010. 1526
- Wake, L. M., Huybrechts, P., Box, J. E., Hanna, E., Janssens, I., and Milne, G. A.: Surface mass-balance changes of the Greenland ice sheet since 1866, *Ann. Glaciol.*, 50, 178–184, doi:10.3189/172756409787769636, 2009. 1542
- Wouters, B., Chambers, D., and Schrama, E.: GRACE observes small-scale mass loss in Greenland, *Geophys. Res. Lett.*, 35, L20501, doi:10.1029/2008GL034816, 2008. 1525

Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Table 1. Summary of simulations with the respective forcings.

Simulation	Atm. Model	Land ice model	Approx. 1st layer height (m)	Lower boundary conditions	Period	Eccentricity (°)	Obliquity (°)	Precession (w-180°)	Days vernal to autumnal equinox	Days autumnal to vernal equinox
L_cPI	LMDZ4_v2	std.	71	climatol.	PI	0.016724	23.446	102.04	180	180
LScPI	LMDZ4_v2	SISVAT	12	climatol.	PI	"	"	"	"	"
LSiPI	LMDZ4_v2	SISVAT	12	interann.	PI	"	"	"	"	"
L_cPIr	LMDZ4_v3	std.	71	climatol.	PI	"	"	"	"	"
LVcPIr	LMDZ4_v3	std.	12	climatol.	PI	"	"	"	"	"
L_c126	LMDZ4_v2	std.	71	climatol.	126 kyr	0.039710	23.928	291.24	192	168
LSc126	LMDZ4_v2	SISVAT	12	climatol.	126 kyr	"	"	"	"	"
LSi126	LMDZ4_v2	SISVAT	12	interann.	126 kyr	"	"	"	"	"
L_c115	LMDZ4_v2	std.	71	climatol.	115 kyr	0.041421	22.405	111.01	175	185
LSc115	LMDZ4_v2	SISVAT	12	climatol.	115 kyr	"	"	"	"	"
LSi115	LMDZ4_v2	SISVAT	12	interann.	115 kyr	"	"	"	"	"

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.

Table 2. Summary of simulations, with GIS area and total GIS SMB components. Also given is the average percentage of months with negative SMB over all ice sheet grid points.

Model/ Simulation	Lower boundary type	Time period	Greenland ice area (10^6 km 2)	Total precip (Gt yr $^{-1}$)	Evapo- ration (Gt yr $^{-1}$)	% of months w. SMB \leq 0	Runoff (Gt yr $^{-1}$)	SMB model (Gt yr $^{-1}$)	SMB PDD (Gt yr $^{-1}$)
MAR	observed	1980–1999	1701	662	8	–	265	388	450
RACMO	observed	1958–2007	1711	744	26	–	248	466	570
L_cPI	climatol.	PI (ctrl)	1792	607	38	4.8	153	416	457
LScPI	climatol.	PI (ctrl)	1792	640	54	13.1	325	261	233
LSiPI	interann.	PI (ctrl)	1792	662	54	12.9	328	280	262
L_c126	climatol.	126 kyr	1792	699	45	9.8	369	285	228
LSc126	climatol.	126 kyr	1792	718	59	20.1	952	–293	–546
LSi126	interann.	126 kyr	1792	748	60	19.8	920	–231	–445
L_c115	climatol.	115 kyr	1792	601	34	2.9	115	452	538
LSc115	climatol.	115 kyr	1792	644	60	9.0	211	373	499
LSi115	interann.	115 kyr	1792	626	58	9.8	216	351	457
L_cPIr	climatol.	PI (ctrl)	1792	625	40				445
LVcPIr	climatol.	PI (ctrl)	1792	661	84				486

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 1. Climatological monthly mean 2 m-temperature ($^{\circ}\text{C}$) in Greenland at selected AWS stations of the GC-net (Steffen et al., 1996), compared to interpolated model data. **(a)** Humboldt, **(b)** Summit, **(c)** ETH-Camp, **(d)** South Dome. Vertical bars show interannual standard deviation for the observation data; triangles indicate the seasonal mean for boreal summer (left) and the annual mean (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 2. (a) Climatological annual mean 2 m-temperature (°C) in Greenland, from left to right for the regional models RACMO and MAR, the LMDZ reference simulation (L_cPI), LMDZ with modified levels (LV_cPI), and LMDZ-SISVAT with constant climatological and variable IPSL-CM4 lower boundary conditions (LScPI and LSiPI). (b) as (a), but for climatological June-July-August seasonal mean 2 m-temperature (°C). (c) as (a), but for climatological annual mean precipitation (m i.e. yr⁻¹).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 3. (a) Climatological annual mean surface mass balance modelled on Greenland, from left to right for the regional models RACMO and MAR, the LMDZ reference simulation and LMDZ-SISVAT with constant climatological and variable lower boundary conditions. (b) as (a), but for SMB computed with the PDD method (m i.e. yr⁻¹).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 4. (a) Climatological mean boreal winter (DJF) 2 m-temperature, difference between simulations with modified vertical levels and the standard version (LVcPIr – L_cPIr, left) and difference from introducing the snow scheme (LScPI – L_cPI – LVcPIr + L_cPIr, right). (b) as (a), but for boreal summer (JJA). (c) as (a), but for boreal winter mean surface albedo. (d) as (a), but for boreal summer mean surface albedo.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 4. (e) as (a), but for boreal winter mean total cloud cover. (f) as (a), but for boreal summer mean total cloud cover. (g) as (a), but for annual mean precipitation (m i.e. yr^{-1}). (h) as (a), but for annual mean evaporation (m i.e. yr^{-1}).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 5. (a) Climatological annual mean modelled SMB anomaly (m i.e. yr^{-1}) LMDZ-SISVAT – standard LMDZ in Greenland ($^{\circ}\text{C}$) for preindustrial (left), Eemian (center) and glacial inception (right) simulations. (b) as (a), but for SMB computed with the PDD method (m i.e. yr^{-1}).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 6. (a) Climatological annual mean 2 m-temperature anomaly vs. preindustrial simulation (°C) in Greenland for Eemian (top row) and glacial inception (bottom row) simulations, in standard LMDZ (left) and LMDZ-SISVAT with constant climatological (center) and varying (right) lower boundary conditions. (b) as (a), but for climatological mean boreal summer (JJA) temperature (°C). (c) as (a), but for climatological annual mean precipitation (m i.e. yr⁻¹). (d) as (a), but for annual mean fraction of months with negative modelled surface mass balance.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 7. (a) Climatological annual mean modelled surface mass balance, anomaly vs. preindustrial simulation ($^{\circ}\text{C}$) in Greenland for Eemian (top row), and glacial inception (bottom row) simulations, in standard LMDZ (left) and LMDZ-SISVAT with constant climatological (center) and varying (right) lower boundary conditions. (b) as (a), but for SMB computed with the PDD method (m i.e. yr^{-1}).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. 8. (a) Climatological annual mean 2-m-temperature anomaly variable SST vs. fixed SST (°C) in Greenland for preindustrial (left), Eemian (center) and glacial inception (right) simulations, in LMDZ-SISVAT. (b) as (a), but for mean surface albedo. (c) as (a), but modelled Climatological annual mean surface mass balance (m i.e. yr⁻¹). (d) as (a), but for SMB computed with the PDD method (m i.e. yr⁻¹).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. A1. Climatological monthly mean 2 m-temperature ($^{\circ}\text{C}$) in Greenland as in Fig. 1, but at the GC-net AWS stations (Steffen et al., 1996) TUNU-N, NASA-E, NASA-U, Crawford Point, SADDLE, compared to interpolated model data. Vertical bars show interannual standard deviation for the observation data; triangles indicate the seasonal mean for boreal summer (left) and the annual mean (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. A1. As first part of Fig. A1, but for the DMI weather stations at Danmarkshavn, Upernavik, Ilulissat, Illoqqortoormiut, Tasiilaq, Nuuk and Narsarsuaq. Also shown is the comparison of the mean model temperature for all of Greenland.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. A2. Climatological monthly mean precipitation (m i.e. yr^{-1}) in Greenland at the GC-net station locations Humboldt, TUNU-N, NASA-E, NASA-U, Summit, ETH Camp, Crawford Point, SADDLE, and South Dome, compared to interpolated model data. Vertical bars show inter-annual standard deviation for the observation data; triangles indicate the seasonal mean for boreal summer (left) and the annual mean (right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling snow accumulation on Greenland in a GCM

H. J. Punge et al.




Fig. A2. As first part of Fig. A2, but at the DMI weather stations Danmarkshavn, Upernavik, Ilulissat, Iloqqortoormiut, Tasiilaq, Nuuk, and Narsarsuaq. Also shown is the comparison of the mean model precipitation for all of Greenland.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

