

Abstract

The ability to precisely date climate proxies is central to the reconstruction of past climate variations. To a degree, all climate proxies are affected by age uncertainties, which are seldom quantified. This article proposes a probabilistic age model for proxies based on layer-counted chronologies, and explores its use for annually-banded coral archives. The model considers both missing and doubly-counted growth increments (represented as independent processes), accommodates various assumptions about error rates, and allows to quantify the impact of chronological uncertainties on different diagnostics of variability. In one dimension, we find that time uncertainties primarily affect high-frequency signals but also significantly bias the estimate of decadal signals. We further explore tuning to an independent, tree-ring based chronology as a way to identify an optimal age model. In the multivariate case, a synthetic pseudocoral network is used as testing ground to quantify uncertainties in the estimation of spatiotemporal patterns of variability. Even for small error rates, the amplitude of multidecadal variability is systematically overestimated at the expense of interannual variability (ENSO, in this case), artificially flattening its spectrum at periods longer than 10 yr. An approach to correct chronological errors in coherent multivariate records is presented and validated in idealized cases, though it is found difficult to apply in practice due to the large size of the solution space. We end with a discussion of possible extensions of this model and connections to existing strategies for modeling age uncertainties.

1 Introduction

A peculiar feature of the Earth Sciences is that the independent variable used to peer into Earth's past (the age of the sample) is always uncertain to some degree. In paleoclimatology, time uncertainties depend on the record's structure, which belongs to one of two categories:

6079

1. *Tie-point chronologies*, which include most sedimentary and speleothem archives. Such records are dated via a one-to-one mapping between sample depth and age (that is, an age model). This model typically assumes a piecewise-continuous accumulation rate, constrained by a relatively small number of radiometrically-determined tie points (e.g. ^{14}C dates for sedimentary archives, or U/Th dates for speleothem archives). Alternatively, when radiometric dating proves impossible, records are often dated by "tuning" to a reference chronology, either derived astronomically (orbital tuning, e.g. Bender, 2002; Hays et al., 1976; Lisiecki and Raymo, 2005; Martinson et al., 1982) or derived from a layer-counted chronology believed to be more accurate (e.g. the GISP2 record Grootes et al., 1993, as done in Wang et al., 2001 and many others).
2. *Layer-counted chronologies*, for instance tree-rings, varved sediments, ice cores and annually-banded corals. For such records, a chronology is obtained by equating the layer count with the number of years or seasons of archive accumulation elapsed since the creation of the topmost (most recent) observed layer. Equivalently, layers may be identified through observation of a seasonally or annually varying quantity such as isotopic composition Fairbanks and Dodge (1979). Layer-counted chronologies are assigned ages relative to a modern tie-point, unless supplemented by independent age controls, such as by radiometric dating (e.g. Cobb et al., 2003), and often assume continuous time resolution unless observations indicate otherwise (e.g. St. George et al., 2013).

Although banded records can often reach subannual precision, they are not time-certain and these uncertainties can have a large influence on variability and coherence of the records. Tree-ring chronologies are generally considered the most accurately dated, but they require adjustments (cross-dating) to optimize coherence across nearby sites (e.g. Douglass, 1941; Fritts, 1976; Wigley et al., 1987; Cook and Kairiukstis, 1990; Yamaguchi, 1991; Stokes and Smiley, 1996). Similarly, although the quoted uncertainty on ice core layer counting is typically on the order of 2% (Alley et al.,

1997), this value reflects the reproducibility from one chronologist to the next, not the uncertainty about the true date. Seimon (2003) thus showed how two nearby Andean ice cores (Thompson et al., 1985) differed widely when plotted against their published timescales, but that simple adjustments based on volcanic tephra markers could bring the records into excellent agreement. Finally, Emile-Geay et al. (2013) showed how minor age uncertainties in several high-resolution records (in particular coral chronologies) could mute interannual signals in a multiproxy reconstruction of tropical Pacific sea-surface temperature.

Time uncertainties pose a serious challenge to the analysis of paleoclimate signals, as the mathematical machinery devoted to timeseries analysis (e.g. spectral analysis, singular spectrum analysis, wavelet analysis), or multivariate analysis (e.g. principal component analysis, regression analysis, climate field reconstruction) implicitly assumes a perfectly known time axis. In the non-banded case, this problem was recognized early-on and motivated the use of tuning, though this approach is not without dangers (Blaauw, 2012). Recently, several studies have chosen instead to explicitly take time uncertainties into account (Huybers and Wunsch, 2004; Blaauw, 2010; Haam and Huybers, 2010; Blaauw and Christen, 2011; Rhines and Huybers, 2011; Anchukaitis and Tierney, 2012), which has greatly helped quantifying the uncertainties inherent to the interpretation of climate signals in such records.

To our knowledge, banded records have not spurred similar modeling efforts, with the notable exception of Rhines and Huybers (2011), whose model targeted long ice-core chronologies like that of GISP2 (Alley et al., 1997). Their study used a discrete random walk, which assumes that the probability of missing a band, or counting a band twice, is symmetric. However, these probabilities are likely asymmetric for most proxies. Further, the model forced the authors to make some unphysical choices in order to match published estimates of age uncertainties. The model we propose is formulated more intuitively in terms of the probabilities of miscounting a band, and naturally allows for asymmetries in the probability of missing or doubly-counting a band. Our work also

6081

applies the model to multivariate datasets and explores ways to correct for age errors using information shared amongst similar records.

Since dendroclimatology has dealt with layer-counted chronological errors for decades, it would be tempting to draw from their extensive literature for other proxy classes. Unfortunately, extending dendrochronological techniques to other proxy types spanning large spatial scales is problematic, because cross-dating relies on the existence of a strong common signal shared among series. This assumption is well suited to trees within a restricted biome but may no longer be valid for other proxy types with banded age models (ice cores, annually-banded corals, varved sedimentary records), either because too few (if any) replicates are available at a given site, or because one is interested in synchronizing records across a wide geographical area (e.g. a pan-Pacific coral network), where the common signal may be weaker. Nevertheless, this approach was successfully applied to replicated coral records from the reefs offshore of Amédée Island, New Caledonia (DeLong et al., 2007, 2013) and from the Gulf of Mexico (DeLong et al., 2011).

In this article we propose a new probabilistic age model for layer-counted records (Sect. 2.1). We first use this model to quantify the uncertainty associated with spectral estimates from univariate coral series (Sect. 2.2). We then generalize this framework to the multivariate setting (Sect. 3.1), evaluating the impact of chronological uncertainties on principal component analysis derived from a pan-Pacific coral network (Sect. 3.3). Leveraging the mutual information from these signals, we then propose a method to correct for age uncertainties in banded records (Sect. 4), which we validate on synthetic data. Discussion follows in Sect. 5.

6082

(SSA, Broomhead and King, 1986; Fraedrich, 1986; Vautard and Ghil, 1989) and its missing value counterpart (SSAM) (Schoellhamer, 2001) to reconstruct the time series including miscounted layers. SSAM was shown to preserve the variance for as many as $\frac{M}{2}$ missing points within a sliding window of size M (Schoellhamer, 2001). In specific applications, other methods would be relevant (e.g. the Lomb-Scargle periodogram Mudelsee et al., 2009, for spectral analysis with missing data) but interpolating with SSAM allows us to apply other analysis methods without further modification, which will prove useful in multivariate applications (Sect. 3).

2.2 Uncertainty quantification in coral records

The above model is now applied to two coral datasets. The first one is the Havannah island fluorescence record (Isdale et al., 1998), chosen for its particularly long, continuous time-span. Because the record already contains dating inaccuracies, we attempt to correct for age errors as explained in (ii). The age uncertainty in coral records is not known, because many of them are not site-replicated. However, DeLong et al. (2013) suggested an error rate of $-2.1/+3.7\%$ in unreplicated samples. We thus assume a conservative error rate of 0.05 for both missing and doubly-counted annual bands (i.e. $\theta_1 = \theta_2 = 0.05$) and generate an ensemble of 1000 plausible age-corrected realizations. The 95 % confidence interval of the realization ensemble is shown along with the original time series in Fig. 1a.

The corresponding spectral densities were estimated in Fig. 1b using the multi-taper method (MTM Thomson, 1982) for each age model realization with a conservative time-bandwidth product of $7/2$ (results are insensitive to this choice).

It is generally presumed that dating errors should affect the estimation of interannual variance most strongly, but only negligibly affect decadal signals. Figure 1b shows this expectation to be approximately correct, though the impact on decadal variability is substantial. Indeed, the width of the confidence bands roughly triples from periods of 50 to 5 yr.

6085

It is however noteworthy that the spectral uncertainties due to the age perturbations (shaded area) are markedly narrower than the uncertainties in the spectral estimates themselves (dashed lines, obtained via standard χ^2 intervals), even though a widening of the ensemble spread can be observed as the frequency increases. This result persists with the robust method of Chave et al. (1987) and its associated jackknife intervals.

To check that those results are not an artifact of the choice of coral timeseries, we repeat this analysis on a $\delta^{18}\text{O}$ record from Palmyra Island (Cobb et al., 2003). Figure 1c shows the cold-season average (December-January-February, DJF) of a continuous splice (1636–1703) of this record, together with the width of the age-perturbed ensemble using $\theta_1 = \theta_2 = 0.05$. Figure 1d shows the impact of age uncertainties on the estimated spectrum, and confirms the previous results: spectral energy of age-corrupt datasets leaks from the interannual band to lower frequencies, but chronological uncertainties generally produce narrower intervals than those due to the spectral estimation process.

2.3 Age model identification

We have shown how the model described in Sect. 2 can be used to generate ensembles of plausible chronologies for any banded record given some a priori estimate of chronological errors. Learning from other dataserries with more replicated and accurate chronologies can then be used to choose an optimal chronology from among an ensemble of plausible chronologies. For example, Li et al. (2011) adjusted age assignments of U/Th-dated fossil Palmyra island corals, observing that the correlation between the nominally-dated modern record, and the first principal component of the North American Drought Atlas (NADA PC1), interpreted as a proxy for ENSO-related hydroclimatic variability, was highly significant over the 1891–1994 time period. They then used this property to align the older sequences, before the onset of the instrumental period, to NADA PC1 within the range of U/Th dating error. To identify an optimal chronology, we similarly choose the realization that maximizes the correlation between

6086

consider a “pseudocoral” network of 22 corals, whose location is derived from an existing network described in Table 1, and derived from simulated climate fields. Pseudocoral locations were chosen to mimic the position of publicly available¹ coral $\delta^{18}\text{O}$ observations with a resolution better than 2 months (Fig. 3).

5 The pseudocoral observations $\mathbf{C} := \{\zeta, \tau\}$ were simulated according to the forward model of Thompson et al. (2011), a simple representation of the effects of temperature and sea water isotopic variation on the oxygen isotope composition of coral aragonite:

$$\zeta_{ij} = a_1 \text{SST}_{ij} + a_2 \text{SSS}_{ij} + \sigma \cdot \epsilon_{ij} \quad (6)$$

where a_1 and a_2 describe the dependence of oxygen isotopic equilibrium on the temperature of carbonate formation and regional regression coefficients of SSS onto sea water $\delta^{18}\text{O}$ respectively, $\epsilon \sim \mathcal{N}(0, 1)$ is an uncorrelated standard Gaussian white noise process, and SST and SSS stand for sea surface temperature and sea surface salinity, respectively. The SST and SSS fields were extracted from the CMIP3 “H1” simulation of the Geophysical Fluid Dynamics Laboratory (GFDL) Coupled Model 2.0 (Knutson et al., 2006). The GFDL CM2.0 model is a state-of-the art coupled general circulation model whose performance at simulating tropical Pacific climate is extensively described (Wittenberg et al., 2006; Capotondi et al., 2006; Vecchi et al., 2006; Kug et al., 2010; Xie et al., 2010). Briefly, the model successfully captures the observed annual-mean trade winds and precipitation, sea surface temperature, surface heat fluxes, surface currents, equatorial undercurrent, and subsurface thermal structure. CM2.0 displays a robust El Niño Southern Oscillation (ENSO) with multidecadal fluctuations in amplitude, an irregular period between 2 and 5 yr, and a distribution of SST anomalies that is skewed toward warm events as observed. Although the model is limited by biases common to other coupled GCMs, it provides a virtual laboratory of adequate realism to explore the impact of pseudocoral sampling with chronological errors. The data cover the interval [1860, 2000]; the monthly output was averaged over the DJF months to

¹<http://www.ncdc.noaa.gov/paleo/corals.html>

sample of the period of most intense ENSO teleconnections. Our pseudocoral coral matrix thus contains $n = 140$ samples at $p = 22$ sites.

In the remainder of the paper, we will use C as a multivariate testing ground for our method. For this study, we limit ourselves to the noiseless case where $\sigma = 0$, though in principle one would have to account for observational noise. Since our interest lies primarily in assessing uncertainties arising from chronological errors, a noise-free matrix is an appropriate first step. We generate an ensemble of age-perturbed realizations of the pseudocoral dataset by disturbing the chronology τ according to the model of Sect. 3.1 using again $\theta_1 = \theta_2 = 0.05$.

10 3.3 Uncertainty quantification

Inspired by the work of Anchukaitis and Tierney (2012), we first seek to quantify the effect of age uncertainties on the estimation of spatiotemporal modes of variability. Since the CM2.0 model simulates a vigorous ENSO, we seek to answer the following question: were tropical climate variability sampled according to the network of Table 1, how faithfully would ENSO variability be captured in the presence of chronological uncertainties? Figure 3 shows the result of a principal component analysis applied to the error-free network, together with an identical analysis applied to the time-perturbed ensemble. The visualization is analogous to that of Tierney et al. (2013, their Fig. 2).

The unperturbed mode accounts for 25% of the total variance, and is dominated by the contributions of central and eastern-Pacific pseudocorals. Its principal component (PC) is characterized by a concentration of variance in the interannual band (Fig. 3c, red curves), and its spatial structure (light colored dots with white outline) closely matches the SST pattern obtained by regressing the principal component onto the SST field, meaning that SST dominates SSS in Eq. (6), at least on interannual timescales. The same map presents results from the time-uncertain ensemble, showing that in the presence of time uncertainties the vast majority of positive loadings (dark-colored disks, circled in black) tend to shrink compared to their unperturbed counterparts (light-colored disks, circled in white). The picture is less clear for neg-

ative loadings, but those exert a lesser control on the overall structure. In the age-perturbed ensemble, the amount of variance explained by the ENSO mode ranges between 6 and 10% (median value: 8%) down from 25% for the error-free data. In such cases, the ENSO mode thus ceases to be the leading mode, as most of its energy is transferred to a trend mode similar to the one identified by Thompson et al. (2011). In our case, the importance of this mode is mainly an artifact of dating errors. A spectral analysis of the age-perturbed PC (Fig. 3c, blue colors) shows a systematic increase in multidecadal variability (10–50 yr periods) at the expense of interannual variability. The transfer of power from high to low-frequencies may be seen by computing the spectral slope β , where we fit the model $S(f) \propto f^\beta$ to the continuum of each spectrum ($f \leq 1/15$). The spectrum of the unperturbed ENSO PC has a slope $\beta_o = -0.72$, while the age-perturbed ensemble shows values centered around a median value $\beta_m = -0.22$ with an inter-quartile range of $[-0.41, 0]$. The loss of coherency thus has the effect of flattening the spectrum of the ENSO mode. Even when the error rate is 50 times smaller ($\theta = 0.001$), one still observes that age uncertainties tend to exaggerate multidecadal (20 yr and longer) variations at the expense of interannual variations (Fig. 5). This may explain why Ault et al. (2009) found, using a similar PCA approach, that their coral network exhibited very energetic variability at such scales, with an amplitude that grew back in time. The presence of small age uncertainties could explain part of the multidecadal power enhancement they observed in the late nineteenth century, since fewer instrumental observations were available to provide chronological anchors to coral timeseries at that time than in the twentieth century. Our analysis suggests that the accumulation of age model uncertainty is one possible – non-climatic – explanation for this apparent enhancement of low-frequency variability in an all-coral network, though a more detailed investigation (incorporating knowledge of error rates in each of their records) would be required to evaluate this quantitatively.

6091

4 Coherence-based identification of age models

As in Sect. 2.3, one would like to choose an optimal chronology among the ensemble of competing age models. To do so, we apply an optimization principle based on the idea that all pseudocorals in our network share some information about a common signal (ENSO in our pseudocoral benchmark). Indeed, coral networks have been shown to record spatiotemporally coherent climate modes (Evans et al., 1998, 2000, 2002), a property shared by our pseudocoral network benchmark (Fig. 3). Moreover, we showed in Sect. 3.3 that chronological errors lead to a loss of coherency among records; that is, we expect age errors to lower interannual coherency among records compared to what it might otherwise be. One way to correct for this effect is to pick the age model that maximizes interannual coherency between measurements.

This exercise is however complicated by the large dimensionality of the space of possible solutions. The dimensionality may be estimated in several ways: assuming that age-perturbations follow a Bernoulli process, an $n \times p$ data matrix admits up to 3^{np} age perturbation matrices. This number quickly becomes astronomical, even for moderate n and p (an NP-hard problem). A more relevant measure, however, is the number of solutions for a given error rate. Given a symmetric error rate $\theta_1 = \theta_2 = \theta$, one expects on average, $\binom{np}{\tilde{x}}$ (where \tilde{x} denotes the nearest integer of x) plausible solutions, which induces a serious computational challenge. This is illustrated in Table 2, which charts the number of solutions for various values of np and θ .

In practice, we generate an ensemble of $L = 10^5$ time perturbations $\mathbf{\Delta} = \{\mathbf{\Delta}^{(1)}, \mathbf{\Delta}^{(2)}, \dots, \mathbf{\Delta}^{(L)}\}$ according to Eq. (4), and choose an optimal chronology amongst those realizations. The optimization principle is described next.

4.1 Optimization scheme

Because age errors tend to lower coherency amongst records (Fig. 4), the quantity we seek to maximize is the generalized magnitude squared coherency (GMSC,

6092

high dimensionality of the solution space can be readily observed here, noticing that the realization maximizing the GMSC barely surpasses the reference coherence spectrum in addition to its distance to the original chronology exceeding the distance between the reference and true data. The issue may be alleviated if there is some prior information about the timing of age errors, for instance an observed mismatch between how two series portray an event – like a historic volcanic eruption – that should be synchronous.

5 Discussion

In this article, we put forth a probabilistic model for layer-counted chronologies. The model was designed for univariate as well as multivariate datasets, and is flexible enough to accommodate location-specific and asymmetric error rates. We used this model to generate ensembles of time-uncertain datasets, which we analyzed using spectral methods and principal component analysis.

In the univariate case, we showed that age-perturbations tend to blur variability between interannual and multidecadal scales. In the multivariate case, we showed that even small age offsets between records may significantly lower coherency between records, producing an apparent amplitude modulation of high-frequency signals. These short-term phase interferences thus led to long-term effects, exaggerating low-frequency variability at the expense of interannual signals. This suggests that at least part of the reason for the observed nineteenth century increase in the amplitude of low-frequency variability (Ault et al., 2009) could be the presence of age errors.

We also attempted to correct for age errors using an optimization principle based on the global coherence of a multivariate record. The large dimensionality of the age model space, however, made the random sampling scheme an inefficient method of exploration. There is no easy way around this problem. Nonetheless, we showed that our optimization strategy was able to correct dating errors when sufficient restrictions were placed on the solution space, provided the mutual information was sufficiently strong.

6097

There are important limitations to our study. First and foremost, our framework assumes that the error rates are known. Ideally, one would estimate the parameters (θ_1, θ_2) from the data, but this is only possible in the presence of independent information (e.g. replicated records, or absolute chronostratigraphic markers like historically-dated events DeLong et al., 2007, 2013). We also restricted our analysis to symmetric error rates, which is an unlikely scenario according to DeLong et al. (2013) who emphasized that double counting is typically more common than missing years. Considering that the events perturbing the chronologies are hidden states of the process and are not being directly measured, a maximum likelihood principle would have to be employed to evaluate the error rate, which substantially complicates the problem of designing a likelihood function. However, when the common signal driving the record is strong enough, a coherency-based Bayesian approach could potentially learn from the data. In this case, one would relax the assumption of known rate parameters (θ_1, θ_2) in favor of prior distributions on those parameters, which would incorporate as much scientific information as possible. In this regard, one should note that most geochronologies are developed with the purpose of producing the single best representation of past changes. We suggest that there would be merit in also providing raw chronologies so that the error rate may be estimated via comparing it with other, absolutely dated records (for instance, instrumental observations at a nearby site, if available). In other words, we hope to encourage field scientists to report these error rates rather than exclusively seeking to reduce them, as this would greatly aid the quantification of uncertainties associated with each record.

Second, we assumed that time perturbations follow a homogeneous Poisson process, with a uniform rate for all pseudocorals considered and assumed that they are symmetric, which is unlikely. In nature, each record will display a different rate, and in situations of uneven dating conditions (e.g. annual bands being more or less visible in a given time interval), an inhomogeneous Poisson process would constitute a more suitable model within a single record. The error rate could be informed by local expertise and change with the location of the data as well as with time and account for known

6098

climatic events that could have affected the coral or breaks, boring and growth structures that are present in the core. Assessments of where and when corals might have over or under-grown impose restrictions on the age model space, which would greatly improve the ability of our optimization principle to identify the correct chronology.

5 Third, we assumed that age perturbations taking place at different locations are independent. Nonetheless, one could easily conceive of a scenario in which a regional phenomenon would simultaneously affect the growth of all corals located in one area. The model would therefore have to include regional random time perturbations, which could be done by specifying parameters conditional on the climate.

10 Fourth, we assumed that all records considered had annual resolution, while in reality it could range from a month to several years. Adapting the method to a network of variable resolution would require the conditioning of the rate parameter θ on the resolution of the record's chronology, as for instance, monthly data tends to present a lower error rate than yearly data, but would add subannual errors from selecting incorrect tie points in the seasonal cycle particularly in regions where the seasonal cycle and annual density banding are weak (e.g., key ENSO sensitive regions of the central Pacific).

15 Fifth, we assumed that the latest time t_i is known exactly (except in Sect. 2.3), which is only true of records that grew continuously until that point (e.g. living corals, living trees). We could refine our fossil corals framework with a mixed approach, where the age of tie points would be allowed to vary according to their distribution (e.g. a Gaussian), and the layer-count model would be extended on either side.

20 Finally, our optimization scheme was shown to be effective only in extremely restricted situations (error-free measurements of $\delta^{18}\text{O}$, age model ensemble spanning the true age model). It remains to be seen whether these conditions could ever be met in practice. One potential problem is the fact that the "clock" used to synchronize the records does not beat periodically: ENSO is only a quasi-periodic phenomenon, even in the GFDL CM2.0 model. Perhaps a more regular cycle (annual, semi-annual) would provide stronger synchronization constraints (but it would limit the analysis to records

6099

that have seasonal or better resolution). Alternatively, one could restrict the optimization process to small, regional ensembles of proxies that share a stronger common signal, much like dendrochronological approaches.

5 With these caveats in mind, several points deserve mention. First, the simple example of Fig. 4 demonstrates how miscounting errors may produce an apparent amplitude modulation of otherwise stationary signals, producing spurious trends in the average derived from them. Because climate reconstruction methods rely in some fashion on a weighted average of observations assumed to be contemporaneous, this may partially explain why many ENSO reconstructions (e.g. Mann et al., 2000; Wilson et al., 10 2010; McGregor et al., 2010; Li et al., 2011; Emile-Geay et al., 2013) display increasing trends in ENSO variance over time (McGregor et al., 2013). As pointed out in the latter study, computing ENSO variance on individual records prior to averaging them together is more robust to chronological errors. In the meantime, interpreting trends in variance obtained from the average of age-uncertain records as reflecting a response to external forcing may be perilous. At the very least, a probabilistic model such as the 15 one proposed here can provide a null hypothesis against which to assess such trends.

More generally, a model such as this one may be used to evaluate the plausible impact of dating uncertainties on all manner of inferences derived from time-uncertain records. Just as proxy system models are now becoming a standard tool to interpret 20 environmental controls in paleoclimate proxies, chronological models should become an integral component of proxy system models for age-uncertain paleodata, so that they may be used to improve interpretation, reconstruction, uncertainty estimation, and data/model comparison studies (Evans et al., 2013). Because chronological errors might themselves depend on climatic events (see introduction), a complete proxy system model should in fact allow environmental controls to influence chronological 25 information.

We hope this study stimulates a discussion on how to best represent chronological uncertainty in layer-counted proxy archives, and leads to a better characterization of these uncertainties.

6100

1. its mean is equal to $\lambda_1 - \lambda_2$, its variance to $\lambda_1 + \lambda_2$.
2. for $\lambda_1 = \lambda_2$, the distribution is symmetric, centered at zero.
3. for $\lambda_1 = \lambda_2$ large enough, it is well approximated by a normal distribution $\mathcal{N}(0, \lambda_1 + \lambda_2)$, thus the difference between the Poisson model and the binomial model (which also converges to a normal distribution) becomes negligible.

Acknowledgements. M.C., J.E.G., M.N.E., N.M. and D.M.T. acknowledge funding from NOAA award NA10OAR4310115. Matlab code implementing the age model ensembles is freely available on the Common Climate Google Code repository (<https://code.google.com/p/common-climate/>) (will be released after manuscript acceptance).

10 References

- Abram, N. J., Gagan, M. K., Cole, J. E., Hantoro, W. S., and Mudelsee, M.: Recent intensification of tropical climate variability in the Indian Ocean, *Nat. Geosci.*, 1, 849–853, doi:10.1038/ngeo357, 2008. 6111
- Abramowitz, M. and Stegun, I. A.: *Handbook of Mathematical Functions*, National Bureau of Standards, Applied Math #55, Dover Publications, New York, 1965. 6102
- Alley, R. B., Shuman, C. A., Meese, D. A., Gow, A. J., Taylor, K. C., Cuffey, K. M., Fitzpatrick, J. J., Grootes, P. M., Zielinski, G. A., Ram, M., Spinelli, G., and Elder, B.: Visual stratigraphic dating of the GISP2 ice core: basis, reproducibility, and application, *J. Geophys. Res.-Atmos.*, 102, 26367–26381, doi:10.1029/96JC03837, 1997. 6080, 6081
- Anchukaitis, K. J. and Tierney, J. E.: Identifying coherent spatiotemporal modes in time-uncertain proxy paleoclimate records, *Clim. Dynam.*, 41, 1–16, doi:10.1007/s00382-012-1483-0, 2012. 6081, 6090
- Asami, R., Yamada, T., Iryu, Y., Quinn, T. M., Meyer, C. P., and Pauley, G.: Interannual and decadal variability of the western Pacific sea surface condition for the years 1787–2000: reconstruction based on stable isotope record from a Guam coral, *J. Geophys. Res.*, 110, C05018, doi:10.1029/2004JC002555, 2005. 6111

6103

- Ault, T. R., Cole, J. E., Evans, M. N., Barnett, H., Abram, N. J., Tudhope, A. W., and Linsley, B. K.: Intensified decadal variability in tropical climate during the late 19th century, *Geophys. Res. Lett.*, 36, L08602, doi:10.1029/2008GL036924, 2009. 6091, 6097
- Bender, M. L.: Orbital tuning chronology for the Vostok climate record supported by trapped gas composition, *Earth Planet. Sc. Lett.*, 204, 275–289, doi:10.1016/S0012-821X(02)00980-9, 2002. 6080
- Blaauw, M.: Methods and code for “classical” age-modelling of radiocarbon sequences, *Quat. Geochronol.*, 5, 512–518, doi:10.1016/j.quageo.2010.01.002, 2010. 6081
- Blaauw, M.: Out of tune: the dangers of aligning proxy archives, *Quaternary Sci. Rev.*, 36, 38–49, doi:10.1016/j.quascirev.2010.11.012, 2012. 6081
- Blaauw, M. and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process, *Bayesian Anal.*, 6, 457–474, 2011. 6081
- Boiseau, M., Juillet-Leclerc, A., Yiou, P., Salvat, B., Isdale, P., and Guillaume, M.: Atmospheric and oceanic evidences of El Niño-Southern Oscillation events in the south central Pacific Ocean from coral stable isotopic records over the last 137 yr, *Paleoceanography*, 13, 671–685, doi:10.1029/98PA02502, 1998. 6111
- Broomhead, D. and King, G. P.: Extracting qualitative dynamics from experimental data, *Physica D*, 20, 217–236, doi:10.1016/0167-2789(86)90031-X, 1986. 6085
- Capotondi, A., Wittenberg, A., and Masina, S.: Spatial and temporal structure of tropical Pacific interannual variability in 20th century coupled simulations, *Ocean Model.*, 15, 274–298, 2006. 6089
- Chakrabarty, K., Sitharama Iyengar, S., Qi, H., and Cho, E.: Grid coverage for surveillance and target location in distributed sensor networks, *IEEE T. Comput.*, 51, 1448–1453, 2002. 6093
- Charles, C. D., Hunter, D. E., and Fairbanks, R. G.: Interaction between the ENSO and the Asian Monsoon in a coral record of tropical climate, *Science*, 277, 925–928, doi:10.1126/science.277.5328.925, 1997. 6111
- Charles, C. D., Cobb, K., Moore, M. D., and Fairbanks, R. G.: Monsoon-tropical ocean interaction in a network of coral records spanning the 20th century, *Mar. Geol.*, 201, 207–222, doi:10.1016/S0025-3227(03)00217-2, 2003. 6111
- Chave, A. D., Thomson, D. J., and Ander, M. E.: On the robust estimation of power spectra, coherences, and transfer functions, *J. Geophys. Res.*, 92, 633–648, doi:10.1029/JB092iB01p00633, 1987. 6086

6104

- Cobb, K. M., Charles, C. D., Cheng, H., and Edwards, R. L.: El Niño/Southern Oscillation and tropical Pacific climate during the last millennium, *Nature*, 424, 271–276, 2003. 6080, 6086, 6114, 6116
- Cobb, K. M., Charles, C. D., and Hunter, D. E.: A central tropical Pacific coral demonstrates Pacific, Indian, and Atlantic decadal climate connections, *Geophys. Res. Lett.*, 28, 2209–2212, doi:10.1029/2001GL012919, 2001. 6111
- Cook, E. R. and Kairiukstis, L. A.: *Methods of Dendrochronology: Applications in the Environmental Sciences*, Kluwer Academic Publishers, Dordrecht, 1990. 6080
- Damassa, T. D., Cole, J. E., Barnett, H. R., Ault, T. R., and McClanahan, T. R.: Enhanced multidecadal climate variability in the seventeenth century from coral isotope records in the western Indian Ocean, *Paleoceanography*, 21, PA2016, doi:10.1029/2005PA001217, 2006. 6111
- DeLong, K. L., Quinn, T. M., and Taylor, F. W.: Reconstructing twentieth-century sea surface temperature variability in the southwest Pacific: a replication study using multiple coral Sr/Ca records from New Caledonia, *Paleoceanography*, 22, PA4212, doi:10.1029/2007PA001444, 2007. 6082, 6098
- DeLong, K. L., Flannery, J. A., Maupin, C. R., Poore, R. Z., and Quinn, T. M.: A coral Sr/Ca calibration and replication study of two massive corals from the Gulf of Mexico, *Palaeogeogr. Palaeoclimatol.*, 307, 117–128, 2011. 6082
- DeLong, K. L., Quinn, T. M., Taylor, F. W., Shen, C.-C., and Lin, K.: Improving coral-base paleoclimate reconstructions by replicating 350 yr of coral Sr/Ca variations, *Palaeogeogr. Palaeoclimatol.*, 373, 6–24, 2013. 6082, 6083, 6085, 6098
- Douglass, A. E.: Crossdating in dendrochronology, *J. Forestry*, 39, 825–831, 1941. 6080
- Emile-Geay, J. and Eshleman, J. A.: Toward a semantic web of paleoclimatology, *Geochem. Geophys. Geosyst.*, 14, 457–469, doi:10.1002/ggge.20067, 2013. 6111
- Emile-Geay, J., Cobb, K., Mann, M., and Wittenberg, A. T.: Estimating central equatorial Pacific SST variability over the past millennium, Part 2: Reconstructions and implications, *J. Climate*, 26, 2329–2352, doi:10.1175/JCLI-D-11-00511.1, 2013. 6081, 6100
- Evans, M. N., Kaplan, A., and Cane, M. A.: Optimal sites for coral-based reconstruction of global sea surface temperature, *Paleoceanography*, 13, 502–516, doi:10.1029/98PA02132, 1998. 6092

6105

- Evans, M. N., Kaplan, A., and Cane, M. A.: Intercomparison of coral oxygen isotope data and historical sea surface temperature (SST): potential for coral-based SST field reconstructions, *Paleoceanography*, 15, 551–562, 2000. 6092
- Evans, M. N., Kaplan, A., and Cane, M. A.: Pacific sea surface temperature field reconstruction from coral $\delta^{18}\text{O}$ data using reduced space objective analysis, *Paleoceanography*, 17, 7–1, doi:10.1029/2000PA000590, 2002. 6092
- Evans, M., Tolwinski-Ward, S., Thompson, D., and Anchukaitis, K.: Applications of proxy system modeling in high resolution paleoclimatology, *Quaternary Sci. Rev.*, 76, 16–28, 2013. 6100
- Fairbanks, R. G. and Dodge, R. E.: Annual periodicity of the $^{18}\text{O}/^{16}\text{O}$ and $^{13}\text{C}/^{12}\text{C}$ ratios in the coral *Montastrea annularis*, *Geochim. Cosmochim. Acta*, 43, 1009–1020, 1979. 6080
- Felis, T., Pätzold, J., Loya, Y., Fine, M., Nawar, A. H., and Wefer, G.: A coral oxygen isotope record from the northern Red Sea documenting NAO, ENSO, and North Pacific teleconnections on Middle East climate variability since the year 1750, *Paleoceanography*, 15, 679–694, doi:10.1029/1999PA000477, 2000. 6111
- Fraedrich, K.: Estimating the dimensions of weather and climate attractors, *J. Atmos. Sci.*, 43, 419–432, doi:10.1175/1520-0469(1986)043<0419:ETDOWA>2.0.CO;2, 1986. 6085
- Fritts, H. C.: *Tree Rings and Climate*, Academic Press, 567 pp., London, 1976. 6080
- Grootes, P. M., Stuiver, M., White, J. W. C., Johnsen, S., and Jouzel, J.: Comparison of oxygen isotope records from the GISP2 and GRIP Greenland ice cores, *Nature*, 366, 552–554, doi:10.1038/366552a0, 1993. 6080
- Guttman, I.: *Statistical Tolerance Regions: Classical and Bayesian*, vol. 26, Griffin's statistical monographs series, Publisher: Hafner Pub. Co. the University of Michigan, 1970. 6095
- Haam, E. and Huybers, P.: A test for the presence of covariance between time-uncertain series of data with application to the Dongge Cave speleothem and atmospheric radiocarbon records, *Paleoceanography*, 25, PA2209, doi:10.1029/2008PA001713, 2010. 6081
- Hamming, R. W.: Error detecting and error correcting codes, *Bell Syst. Tech. J.*, 29, 147–160, 1950. 6095
- Hays, J. D., Imbrie, J., and Shackleton, N. J.: Variations in the Earth's orbit: pacemaker of the ice ages, *Science*, 194, 1121–1132, 1976. 6080
- Huybers, P. and Wunsch, C.: A depth-derived Pleistocene age model: Uncertainty estimates, sedimentation variability, and nonlinear climate change, *Paleoceanography*, 19, PA1028, doi:10.1029/2002PA000857, 2004. 6081

6106

- Isdale, P. J., Stewart, B. J., Tickle, K. S., and Lough, J. M.: Palaeohydrological variation in a tropical river catchment: a reconstruction using fluorescent bands in corals of the Great Barrier Reef, Australia, *The Holocene*, 8, 1–8, doi:10.1191/095968398670905088, 1998. 6085, 6114
- 5 Knutson, T. R., Delworth, T. L., Dixon, K. W., Held, I. M., Lu, J., Ramaswamy, V., Schwarzkopf, M. D., Stenchikov, G., and Stouffer, R. J.: Assessment of twentieth-century regional surface temperature trends using the GFDL CM2 coupled models, *J. Climate*, 19, 1624–1651, doi:10.1175/JCLI3709.1, 2006. 6089
- Kug, J.-S., Choi, J., An, S.-I., Jin, F.-F., and Wittenberg, A. T.: Warm pool and cold tongue El Niño events as simulated by the GFDL CM2.1 coupled GCM, *J. Climate*, 23, 1226–1239, doi:10.1175/2009JCLI3293.1, 2010. 6089
- 10 Kuhnert, H., Pätzold, J., Hatcher, B., Wyrwoll, K. H., Eisenhauer, A., Collins, L. B., Zhu, Z. R., and Wefer, G.: A 200 yr coral stable oxygen isotope record from a high-latitude reef off Western Australia, *Coral Reefs*, 18, 1–12, doi:10.1007/s003380050147, 1999. 6111
- Kuhnert, H., Pätzold, J., Wyrwoll, K.-H., and Wefer, G.: Monitoring climate variability over the past 116 yr in coral oxygen isotopes from Ningaloo Reef, Western Australia, *Int. J. Earth Sci.*, 15, 88, 725–732, doi:10.1007/s005310050300, 2000. 6111
- Li, J., Xie, S.-P., Cook, E. R., Huang, G., D'Arrigo, R., Liu, F., Ma, J., and Zheng, X.-T.: Interdecadal modulation of El Niño amplitude during the past millennium, *Nat. Clim. Change*, 1, 114–118, 2011. 6086, 6087, 6100
- 20 Linsley, B. K., Dunbar, R. B., Wellington, G. M., and Mucciarone, D. A.: A coral-based reconstruction of intertropical convergence zone variability over Central America since 1707, *J. Geophys. Res.*, 99, 9977–9994, doi:10.1029/94JC00360, 1994. 6111
- Linsley, B. K., Ren, L., Dunbar, R. B., and Howe, S. S.: El Niño Southern Oscillation (ENSO) and decadal-scale climate variability at 10° N in the eastern Pacific from 1893 to 1994: a coral-based reconstruction from Clipperton Atoll, *Paleoceanography*, 15, 322–335, doi:10.1029/1999PA000428, 2000. 6111
- Linsley, B. K., Kaplan, A., Gouriou, Y., Salinger, J., deMenocal, P. B., Wellington, G. M., and Howe, S. S.: Tracking the extent of the South Pacific Convergence Zone since the early 1600s, *Geochem. Geophys. Geos.*, 7, Q05003, 2006. 6111
- 30 Linsley, B. K., Zhang, P., Kaplan, A., Howe, S. S., and Wellington, G. M.: Interdecadal-decadal climate variability from multicoral oxygen isotope records in the South Pacific Convergence Zone region since 1650 AD, *Paleoceanography*, 23, PA2219, doi:10.1029/2007PA001539, 2008. 6111

6107

- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, PA1003, doi:10.1029/2004PA001071, 2005. 6080
- Mann, M., Gilleb, E., Bradley, R., Hughes, M., Overpeck, J., Keimigc, F., and Gross, W.: Global temperature patterns in past centuries: an interactive presentation, *Earth Interact.*, 4, 1–29, 2000. 6100
- 5 Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global temperature for the past 11 300 yr, *Science*, 339, 1198–1201, doi:10.1126/science.1228026, 2013. 6087
- Martinson, D. G., Menke, W., and Stofa, P.: An inverse approach to signal correlation, *J. Geophys. Res.*, 87, 4807–4818, doi:10.1029/JB087iB06p04807, 1982. 6080
- 10 McGregor, S., Timmermann, A., and Timm, O.: A unified proxy for ENSO and PDO variability since 1650, *Clim. Past*, 6, 1–17, doi:10.5194/cp-6-1-2010, 2010. 6100
- McGregor, S., Timmermann, A., England, M. H., Elison Timm, O., and Wittenberg, A. T.: Inferred changes in El Niño–Southern Oscillation variance over the past six centuries, *Clim. Past*, 9, 2269–2284, doi:10.5194/cp-9-2269-2013, 2013. 6100
- 15 Mudelsee, M., Scholz, D., Röthlisberger, R., Fleitmann, D., Mangini, A., and Wolff, E. W.: Climate spectrum estimation in the presence of timescale errors, *Nonlinear Proc. Geoph.*, 16, 43–56, doi:10.5194/npg-16-43-2009, 2009. 6085
- PAGES2K Consortium: Continental-scale temperature variability during the past two millennia, *Nat. Geosci.*, 6, 339–346, doi:10.1038/ngeo1797, 2013. 6087
- 20 Pfeiffer, M., Timm, O., Dullo, W.-C., and Podlech, S.: Oceanic forcing of interannual and multidecadal climate variability in the southwestern Indian Ocean: evidence from a 160 yr coral isotopic record (La Réunion, 55° E, 21° S), *Paleoceanography*, 19, PA4006, doi:10.1029/2003PA000964, 2004. 6111
- 25 Quinn, T. M., Crowley, T. J., Taylor, F. W., Henin, C., Joannot, P., and Join, Y.: A multicentury stable isotope record from a New Caledonia coral: interannual and decadal sea surface temperature variability in the southwest Pacific since 1657 AD, *Paleoceanography*, 13, 412–426, doi:10.1029/98PA00401, 1998. 6111
- Quinn, T. M., Taylor, F. W., and Crowley, T. J.: Coral-based climate variability in the Western Pacific Warm Pool since 1867, *J. Geophys. Res.-Oceans*, 111, C11006, doi:10.1029/2005JC003243, 2006. 6111
- 30 Ramirez, D., Via, J., and Santamaria, I.: A generalization of the magnitude squared coherence spectrum for more than two signals: definition, properties and estimation, in: *IEEE Interna-*

6108




Fig. 1. Continued.

6115




Fig. 2. Palmyra relict coral $\delta^{18}\text{O}$ records (top): published data SB3 in red, SB8 in orange and SB13 in brown (Cobb et al., 2003), the shaded area is the 95 % confidence interval of the ensemble of realizations with final time shifted $t_i \pm 10$ yr and $\theta_1 = \theta_2 = 0.05$ and the blue chronology corresponds to the realizations that maximize the correlation with the first principal component of the NADA record (bottom) over each continuous section. The correlation coefficients with NADA PC1 are also shown, wherein ρ is computed using the aligned Palmyra chronologies, and ρ_0 using the original chronologies.

6116




Fig. 3. Spatiotemporal uncertainty quantification on a pseudocoral network. **(a)** EOF loadings (circles) corresponding to the ENSO mode of an ensemble of age-perturbed pseudocoral records with miscounting rate $\theta = 0.05$. EOF loadings for error-free data are shown in light colors circled in white, while the median and 95 % quantile are shown by dark disks and black-circled disks, respectively. Contours depict the SST field associated with the mode's principal component PC **(b)**, whose power spectrum is shown in **(c)**. Results for the time-uncertain ensemble are shown in blue: median (solid line), 95 % confidence interval (light-filled area) and interquartile range [25–75 %] (dark-filled area). Results for the original (error-free) dataset are depicted by solid red lines. Dashed red lines denote χ^2 error estimates for the MTM spectrum of the error-free dataset.

6117




Fig. 4. Illustration of the phase distortion of coherent signals due to miscounted annual bands ($\theta_1 = \theta_2 = 0.05$). (top) Five hypothetical harmonic signals (middle) the same signals after introduction of age perturbations (bottom) the same age-perturbed signals, after making the (erroneous) assumption that 1 layer amounts to 1 yr. Note the loss of coherency reflected in the ensemble average, which is performed as part of composites or most climate reconstructions.

6118




Fig. 7. Coherence spectra **(a)** and on the next page Hamming distance distributions to the original chronology **(b)** for ensembles of chronologies sampled around the reference perturbation matrix (with $\theta = 0.05$), in restricted neighborhood of sizes $2 \cdot N_v + 1$ for N_v increasing from 1 to 10. The thick red curve represents the coherence spectrum for the error-free data, the results related to the reference perturbed dataset are shown in green and plotted in black are the results for the ensemble realization (sampled across the entire solution space) that maximizes the GMSC (as described in Sect. 4.1). Finally the results for the neighborhood ensembles (shaded areas): the 95 % CIs and Hamming distance distributions, and corresponding realizations maximizing the GMSC (solid lines) are shown in dark blue, blue, pink and light brown for $N_v = 1, 4, 7$ and 10, respectively.

6121




Fig. 7. Continued.

6122




Fig. A1. Skellam probability mass functions for different values of its parameters λ_1 and λ_2 .