

Dear Dr. Winguth,

Please find our revised manuscript of “Climate trends in northern Ontario and Quebec from borehole temperature profiles” (cp-2016-55). We thank the anonymous reviewer and A.E.Taylor for their reviews. We appreciate the useful comments by R.Way. In response to their suggestions and questions, we have made several changes to the manuscript and added some information to clarify some points about data collection and processing. A detailed list and explanation of the changes follows. The changes have been highlighted on the attached copy of the revised manuscript.

Response to Comments

In response to Anonymous Reviewer

1) Anonymous Reviewer: Borehole sites: In adequately determining if borehole sites are appropriate for use in climate reconstructions, several criteria are required. While the authors have addressed several of the sites and determined they were unsuitable (as presented in Table 2 of the manuscript), information regarding the other sites is not included that would aid a reader in understanding the conditions at the boreholes. For example, no discussion of slope, topography, vegetation or surface material is given, although the authors do reference previous studies. A discussion of vegetation and ground cover at the sites would be extremely useful, however, especially considering that the argument that one site (Thierry Mine) may have additional warming due to the removal of vegetation was put forth. Further, some sites are said to be "too shallow" or on

the side of steep hills. What exactly is too shallow and steep? Can a quantitative discussion replace the qualitative explanation? Also, are all of the boreholes vertical? At least one site was excluded because it was plunging under a lake. It should be clear.

We agree with the reviewer. The dip of the boreholes varies between 40° and 90°. This has been added to Table 1 (P21). Furthermore, a more detailed description of each site will be provided in an appendix.

In the original manuscript, we qualitatively explained why some boreholes were rejected (too shallow, proximity to a lake, slope, etc.). This means that holes less than 300 m were rejected for being too shallow. Holes were rejected for being near a lake when the mean distance was less than the depth of the hole, or less than 300 m. Holes were deemed too steep and rejected if they had slope of 5% or more over distance comparable to depth. This explanation has been added to the revised manuscript (P5L33, P6L1-2).

2) Anonymous Reviewer: ... if the results of the removal of the steady state gradient as shown in Figures 2 and 3 are different if the length (100 m) is modified?

In their heat flow studies, Lévy et al. (2010) and Jaupart et al. (2014) have determined the heat flow for the majority of the boreholes presented in this manuscript and estimated how heat flux varies with the depth interval where it is estimated. They found that, below 300 m, the standard deviation on the estimates of the heat flux is less than 5% of the mean suggesting that the reference temperature pro-

file does not vary much with depth. This has been clarified in the revised manuscript (P6L11-12).

3) Anonymous Reviewer: The authors state that only one site has a ground surface temperature (GST) that was affected by the LIA. However, based on the temperature anomalies shown in Figures 2 and 3, it would seem that other sites exhibit cooling at the same depths as Otokwin. Musselwhite, TM0608, and CC0713 all have temperature anomalies that indicate cooling at the same general depth. Is this not a LIA signature

The reviewer is correct that the temperature anomalies of Musselwhite, TM0608, and CC0713 indicate cooling. This has been clarified in the revised manuscript (P6L25, P7L2-3).

The profiles and anomalies at Musselwhite and TM0608 are noisy, which could mask a clear LIA signal when inverting the profiles. (Mareschal and Beltrami, 1992) showed that resolution decreases when noise and errors must be filtered. Large singular values are required when dealing with noisy data since they reduce the impact of noise but retain the gross features of the solution. Furthermore, a test was run with a 1 K cooling between 1600 and 1800 and varying the singular value cutoff. This showed that for noise-free synthetic data, a 1 K cooling cannot be resolved with less than 5 singular values. The CC0713 temperature anomaly is less noisy and shows a mild cooling of ≤ 0.2 K. Our test shows that a cooling of this magnitude cannot be resolved in our reconstructions. This was further confirmed when we inverted the site individually and observed no LIA signal. This has been added in the revised manuscript (P7L29-

33, P8L1-2).

4) Anonymous Reviewer: Also, the anomaly shown in CC0712 (Figure 3, top left) has a very interesting profile. What is the cause?

The reviewer is correct. These discontinuities are also observed in the gradient between 100 and 300 m, which could be due to small water flows. This has been added to the revised manuscript (P7L3-5).

5) Anonymous Reviewer: Other questions I have about the results that don't have any explanation or that aren't adequately explained include the assertion that the Thierry Mine signal may be amplified by the clearing of vegetation between 1934 and 1950. However, most of the GST histories show a large increase in temperature at this same time, indicating it may not be vegetation alone. Have the authors done any modeling or do they have any surface temperatures to support this hypothesis

The warming at Thierry Mine is greater than the other sites (at least 0.7 K greater than any other site). During sampling, a large clearing near all three sites was noted. This was further confirmed by examining a satellite image of the region, which located the boreholes ~500 m from a large clearing. In the original manuscript, we hypothesize this warming to be the result of vegetation clearing Lewis and Wang (1998); Lewis (1998).

However, the satellite image also locates the three boreholes ~300 m

from a lake. Lakes disturb a profile if they are at distance less than the depth of the boreholes (Lewis and Wang, 1992). The Thierry Mine boreholes are 530 m or deeper. Therefore, they could be influenced by the presence of the lake. We hypothesize that the greater warming signal is related to the change in vegetation cover and the presence of the nearby lake (Lewis and Wang, 1992, 1998; Lewis, 1998). This has been added to and clarified in the revised manuscript (P7L13-14).

6) Anonymous Reviewer: Lastly, one site (Eleonore) has warming that began considerably earlier than the other sites. Why might this be?

We do not believe Eleonore shows earlier warming. There is a clear recent warming signal (Figure 6).

7) Anonymous Reviewer: ...Corvet, which is located on the side of a 30 m hill. However, what is the slope? How much of an effect does this have? It is still being used, so the authors must think it isn't significant.

Errors were made in the coordinates of Corvet in the manuscript. Corvet is located at 53°19.072'N and 73°55.760'W. Using the elevation of Google Earth images of the site, we see that the topography is less than 5m. This has been corrected in the manuscript.

8) Anonymous Reviewer: The authors discuss what the LIA surface

signal should be for the region, but do not see a ground signal. Perhaps a simple forward model of driving into the ground a surface temperature time series with the appropriate LIA signal and making a comparison to the boreholes would be appropriate? Then, the authors could argue whether the signal is strong enough to actually be observed, or whether it is not seen due to snow or something else. This is similar to the arm waving argument used to interpret a possible ground warming due to longer/deeper snow cover in the region, but it seems that other authors have performed some analysis that may provide quantitative support to their arguments (perhaps Bartlett et al., 2005?)

An appropriate LIA signal for the region is unknown. Pollen data have reconstructed a $\sim 0.3^\circ$ LIA signal Gajewski (1988); Viau and Gajewski (2009); Viau (2012). We ran with a 1 K cooling between 1600 and 1800 and varying singular values. At least five singular values were required to resolve the signal. Furthermore, noisy data requires larger cutoffs to decrease the impact of noise on the solution (Mareschal and Beltrami, 1992). This illustrates that a weak LIA signal cannot be resolved in a noisy environment. This has been clarified in the revised manuscript (P7L29-34, P8L1-2).

9) Anonymous Reviewer: I did notice that on page 11 in the reference section that Jaupart and Mareschal, 2011 was published in Cambridge, not Cabridge; also, the next two references following the previous reference are of Jaupart et al., 2014 and are a duplicate.

This has been fixed in the revised manuscript.

In response to comments by A.E.Taylor

10) A.E.Taylor: My main suggestion is that at least a further brief description should be made to the immediate borehole surface character and subsurface geology, even though this might be more fully covered in the referenced work on these boreholes. Indicate the predominant rock type, possibly with a profile of thermal conductivity that might explain some variations seen in the temperatures.

We agree with the reviewer that a more detailed description of each site is useful. We have added the dip of the boreholes to Table 1 (P21). Furthermore, a detailed description of each site will be provided in an appendix.

11) A.E.Taylor: Equations in sec. 2 indicate that conductivity profiles are part of the program input data with the temperature residuals. I assume there is not significant layering except at Nielson where the thermal depth is used to reduce the effect of conductivity variations.

The reviewer is correct. Nielsen Island is the only site to present significant layering, which is accounted for by using the thermal depth to account for the conductivity variations. This is noted on P5, L31-32 of the revised manuscript.

12) A.E.Taylor: The authors do mention other physical elements or

processes that may affect the results (and conclusions?): snow cover or lack of, borehole area vegetation/forest cover and peat, drainage, surface overburden (any?). These might be augmented by the authors visual observations of the immediate borehole surrounds during logging visits.

Visual observations of the immediate borehole surroundings are routinely noted in our log books when logging the boreholes. Furthermore, we examine the borehole locations on satellite images and topographic maps. This will be included in an appendix with a detailed description of each site.

13) A.E.Taylor: For Camp Coulon, the 3 temperature profiles do appear to be distinctly different (Fig. 3). The authors combine such closely spaced holes for an ensemble inversion, a usual practice. But one might wonder if separate inversions would give very different reconstructed temperature histories that might suggest other factors at play (thermal conductivity? Water flow?). Particularly the odd profile at CC1012, compared to nearby holes and considering such precise temperature measurements.

The three Camp Coulon sites (0712, 0713, 0714) were inverted individually. Similar trends are observed throughout the three GST reconstructions. A warming signal of $\sim 1-1.5$ K occurs at ~ 100 years BP. The maximum temperature at 0712, however, is reached 20 years before that of 0713 and 0714. Differences between the reconstructions could be the result of noise in the profiles. Therefore, we chose to simultaneously invert the profiles to increase the signal to noise ratio.

14) ...Presumably (?) the holes are uncased.

The holes are cased for the upper ~10-15 m, usually until bedrock is reached. This has been clarified in the revised manuscript (P5L17-18).

15) Could the consistent offsets in temperatures at N1015 (~300 m), N1013 (~240 m), and N1012 (~160 m) possibly result from a sub-horizontal water flow (Fig. 2, Ontario) (e.g., geophysical models of Broedehoeft and others). These holes seem spatially very close (how much?) while N1012 is further south, so perhaps justifying the simultaneous inversion, but also questioning the reason for the differing temperature profiles. Similar offset feature at Eleonore (~350 m, Fig. 3, Quebec).

The reviewer is correct. The consistent offsets in temperatures at N1015, N1013, and N1012 could be the result of a sub-horizontal water flow. We examined our log book for any signs of oscillation or instability during the temperature measurements, often a sign of water flow. Several unstable measurements were noted at N1013. However, no instability was noted at the other two sites (N1012 and N1015) but the temperature gradient did reverse. This could be attributed to water flow.

The holes are spatially very close. There was an error made in the coordinates of N1015, which are 52°44'25.5"N and 86°18'11.9"W. This has been fixed in the revised manuscript (Table 1, P21).

16) Maybe better here for the authors to conclude that their GST reconstructions show the potential for permafrost across the region is minimal to absent over the past 5 centuries, with its occurrence highly dependent on surface character and snow cover effects. Suggest add a citation for nearby locations, if any, where permafrost has been documented.

We agree with the reviewer. We have better qualified our conclusions by stating that the GST reconstructions show a minimal potential for permafrost across the region for the past five centuries in the revised manuscript (P1L7-8, P8L32-33, P9L1).

Thibault and Payette (2009) document permafrost in the James Bay peatland bogs near the Northern Quebec sites of Lagrande, Eleonore, and Eastmain. This citation has been added to the manuscript (P8L29-30).

17) ...the temperature scale ranges vary through Fig. 2 and 3 but the sensitivities are consistent; it would be useful to indicate that in the caption as it makes for inter-comparison; same for Fig. 4-6. A few editorial suggestions are on an annotated copy of the PDF manuscript

We appreciated the suggestions and have incorporated them into the manuscript.

We have also considered the comments from yourself and R.Way when revising the manuscript.

In response to R.Way

16) The authors incorrectly state that their sample sites are all located within the sporadic (10-50% of the land surface) to extensive discontinuous zones (50-90% of the land surface). In nearly every case the sampled sites are located within the isolated patches permafrost zone (<10% of land surface) according to maps produced by Heginbottom et al (1995), Payette (2001) and recent spatial numerical modelling of permafrost distribution for Labrador-Ungava (Way and Lewkowicz, 2016). Considering the more realistic permafrost extent, there is no discrepancy between the borehole observations and existing permafrost maps.

We are thankful to R.Way for pointing out that the map we were using is out of date. According to the map from the geological survey, our boreholes lie in a region of discontinuous permafrost. In Quebec, we cover a region classified as having only isolated patches of permafrost. In northern Ontario, Noront (4 holes) is near the southern edge of a region with extensive discontinuous permafrost (50-90%). This has been fixed in the revised manuscript (P3L9-14, P5L26-30, P8L34).

For regions of discontinuous isolated patches of permafrost, where we did not find permafrost at any site, our statistics are not sufficient to draw any conclusion. However, for Noront, near the southern boundary of extensive discontinuous permafrost, using the low frequency of 50%, the probability that not one of 4 holes meets per-

mafrost is low: 1/16 or 6.25%. Furthermore, in Manitoba, we had sampled a region of sporadic discontinuous permafrost (10- 50%) between the towns of Flin Flon, Thompson, and Lynn Lake, but we found that only 1 hole in more than 60 was blocked, as described and discussed by Guillou-Frottier et al. (1998). Using the lowest value of 10% permafrost coverage, the probability that only one hole out of 60 encounters permafrost is only $0.1 \times 10^{-59} \times 60$ or 1.20%; for the 50% frequency, the probability is only 5.2×10^{-17} . This demonstrates a possible discrepancy between borehole observations and existing permafrost maps. We do, however, believe that we must better qualify our conclusions by stating that our reconstructions indicate that a minimal potential for permafrost across the region for the past five centuries. This has been clarified in the manuscript (P1L7-8, P8L32-33, P9L1).

17) R.Way: A further point on a similar subject is that the temperature sampling methodology here is at too coarse a resolution (depth) to detect thin permafrost bodies if they were to exist. In the southern end of the discontinuous zone you would be more likely to find thinner permafrost bodies therefore this is a serious limitation of the study.

We disagree. To avoid repetition of the measurement procedure, we did not sufficiently emphasize the obvious fact that the temperature measurements in boreholes must be performed a long time after drilling when the ground has returned to thermal equilibrium. Should permafrost be present, the hole would be frozen and we would not be able to lower the probe in it to measure temperature. There is no permafrost in the holes that we measured regardless of

the depth sampling interval. A second point is that thin permafrost bodies are very unlikely to be present for simple reasons of thermodynamics. The second law of thermodynamics implies that any temperature oscillation decays with time, and it follows that in steady state the vertical temperature profile can have no maximum or minimum, implying that the existence of a thin frozen layer is impossible in steady state. A frozen underground layer could exist but only as a transient. The life time of such a layer depends on its thickness. For standard values of thermal diffusivity, a 1m thick layer would last less than 1 year.

18) The article also does not present any indication of the land cover types encountered in the study area and correspondingly, does not consider how permafrost is distributed across the landscape (e.g. Shur and Jorgenson, 2007; Jorgenson et al., 2010). In northern Ontario and Quebec, permafrost is largely absent from forested areas at the southern end of the discontinuous zones where snow accumulates while concurrently being present on wind exposed mountaintops (Brown, 1979; Ives, 1979; Allard and Séguin, 1987; Granberg, 1989; Ou et al. 2016a,b; Way and Lewkowitz, 2016). Ignoring these critical variables makes it untenable to draw large-scale conclusions on permafrost from the provided data.

R.Way is correct that we did not consider the landscape in these studies. The main reason being that we use holes of opportunity that were drilled for mining exploration in any surface environment. They represent thus a random unbiased sampling of the landscape. Outside Quebec, the study area is in lowlands without “mountaintops”. The topography near our sites on the Quebec side is more

marked but remains low. Most likely, disagreements arise from the inconsistent and poor quality of the permafrost data because of insufficient sampling and the extreme spatial variability of the land surface.

19) In general, I believe that the discussion of permafrost in this article should be removed in its entirety as the methodology, discussion and interpretations presented are not appropriate for the analysis of permafrost distribution and history. Finally, the lack of consideration of the literature on permafrost in western Quebec and northern Ontario must be addressed.

We appreciate the comment but must disagree. The methodology is sound, and our reconstruction of past temperature histories is based on very simple physics. As much as we understand the need for considering the literature on permafrost, we believe that permafrost studies would gain to be confronted with physical models. We do, however, believe that we must better qualify our conclusions by stating that our reconstructions indicate that a minimal potential for permafrost across the region for the past five centuries. This has been done in the revised manuscript (P1L7-8, P8L32-33, P9L1).

In response to Editor: A.Winguth

20) I noticed that the temperature change (“1-2 K”) and the time period (“for the last 150 years”) in the abstract and discussion and conclusion are vague. Please replace the temperature range with mean and standard deviation and specify the time period (e.g “for

the period 1850 to 2000”).

This has been fixed in the revised manuscript (P1L4, P7,L10-11, P8L25).

We trust that we have addressed all the comments and that our revisions have resulted in improving significantly the manuscript.

Sincerely yours,

Carolyne Pickler
Hugo Beltrami
Jean-Claude Mareschal

References

- Gajewski, K.: Late Holocene climate changes in eastern North America estimated from pollen data, *Quaternary Research*, 29, 255–262, doi:10.1016/0033-5894(88)90034-8, 1988.
- Guillou-Frottier, L., Mareschal, J. C., and Musset, J.: Ground surface temperature history in central Canada inferred from 10 selected borehole temperature profiles, *Journal of Geophysical Research*, 103, 7385–7397, doi:10.1029/98JB00021, 1998.
- Jaupart, C., Mareschal, J., Bouquerel, H., and Phaneuf, C.: The building and stabilization of an Archean Craton in the Superior Province, Canada, from a heat flow perspective, *Journal of Geophysical Research: Solid Earth*, 119, 9130–9155, doi:10.1002/2014JB011018, 2014.
- Lévy, F., Jaupart, C., Mareschal, J.-C., Bienfait, G., and Limare, A.: Low heat flux and large variations of lithospheric thickness in the Canadian Shield, *Journal of Geophysical Research: Solid Earth* (1978–2012), 115, doi:10.1029/2009JB006470, 2010.
- Lewis, T.: The effect of deforestation on ground surface temperatures, *Global and Planetary Change*, 18, 1–13, doi:10.1016/S0921-8181(97)00011-8, 1998.
- Lewis, T. J. and Wang, K.: Influence of terrain on bedrock temperatures, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 98, 87–100, doi:10.1016/0031-0182(92)90190-G, 1992.
- Lewis, T. J. and Wang, K.: Geothermal evidence for deforestation induced warming: Implications for the climatic impact of land development, *Geophysical Research Letters*, 25, 535–538, doi:10.1029/98GL00181, 1998.

- Mareschal, J.-C. and Beltrami, H.: Evidence for recent warming from perturbed geothermal gradients: examples from eastern Canada, *Climate Dynamics*, 6, 135–143, doi:10.1007/BF00193525, 1992.
- Thibault, S. and Payette, S.: Recent permafrost degradation in bogs of the James Bay area, northern Quebec, Canada, *Permafrost and Periglacial Processes*, 20, 383–389, doi:10.1002/ppp.660, 2009.
- Viau, A. and Gajewski, K.: Reconstructing millennial-scale, regional paleoclimates of boreal Canada during the Holocene, *Journal of Climate*, 22, 316–330, doi:10.1175/2008JCLI2342.1, 2009.
- Viau, A. E.: Climate of the last 2,000 years in North America reconstructed from pollen records, *Quaternary International*, 279, 520, doi:10.1016/j.quaint.2012.08.1800, 2012.

Climate trends in northern Ontario and Quebec from borehole temperature profiles

C. Pickler¹, H. Beltrami², and J.-C. Mareschal¹

¹GEOTOP, Centre de Recherche en Géochimie et en Géodynamique, Université du Québec à Montréal, Canada

²Climate & Atmospheric Sciences Institute and Department of Earth Sciences, St. Francis Xavier University, Antigonish, Nova Scotia, Canada

Correspondence to: H.Beltrami (hugo@stfx.ca)

Abstract. The ground surface temperature histories of the past 500 years were reconstructed at 10 sites containing 18 boreholes in northeastern Canada. The boreholes, between 400 and 800 m deep, are located north of 51°N, and west and east of James Bay in northern Ontario and Quebec. We find that both sides of James Bay have experienced similar ground surface temperature histories with a warming of $\sim 1-21.51 \pm 0.76$ K during the last 150 years period of 1850 to 2000, similar to borehole reconstructions for the southern portion of the Superior Province and in agreement with available proxy data. A cooling period corresponding to the little ice age was found at only one site. Despite permafrost maps locating the sites in a region of discontinuous permafrost, the ground surface temperature histories suggest that the potential for permafrost was minimal to absent over across the region the entire region is and was free of permafrost for the past 500 years. This could be the result of air surface temperature interpolation used in permafrost models being unsuitable to represent account for the spatial variability of ground temperatures along with an offset between ground and air surface temperatures due to the snow cover.

1 Introduction

Earth's subsurface thermal regime is governed by the outflow of heat from the interior and by temporal variations in ground surface temperature (GST). The heat flux from the interior of the Earth varies on time scales of the order of a few Myears in active tectonic regions and several 100 Myears in stable continents. It can be considered as steady state relative to the time scale of climatic surface temperature variations. To determine the heat flow from the Earth's interior, temperature-depth profiles are measured in boreholes. In homogeneous rocks with no heat production, the steady state temperature profile linearly increases with depth. Persistent temporal changes in the ground surface energy balance cause variations of the ground temperature that diffuse downwards and are recorded as temperature anomalies superimposed on the linear steady-state geotherm (e.g., Hotchkiss and Ingersoll, 1934; Birch, 1948; Beck, 1977). The extent to which the ground surface temperature changes are recorded is proportional to their duration and amplitude and inversely proportional to the time when they occurred. For periodic oscillations of the surface temperature, the temperature is propagated downward as a damped wave. The amplitude of the wave decreases exponentially with depth over a length scale δ (skin depth) proportional to the square root of the period ($\delta = \sqrt{\kappa T/\pi}$), where κ is the thermal diffusivity of the rock, $\approx 10^{-6} \text{ m}^2 \text{ s}^{-1}$ or $\approx 31.5 \text{ m}^2 \text{ yr}^{-1}$. This damping removes the

high-frequency variability that is present in meteorological records and allows for the preservation of the long-term climatic trends in the ground temperature signal (e.g., Beltrami and Mareschal, 1995).

From the interpretation of temperature-depth profiles, it is possible to infer centennial trends in Earth's surface temperature variations. The first attempt to infer climate history from temperature-depth profiles was the study by Hotchkiss and Ingersoll (1934) who estimated the timing of the ice retreat at the end of the last glaciation. It was, however, not until the 1970s that systematic studies were undertaken to infer past climate from such profiles (e.g., Cermak, 1971; Sass et al., 1971; Vasseur et al., 1983). In the 1980s, with increasing concern over global warming, use of borehole temperature-depth profiles to estimate recent (<300 years) climate change became widespread following the study of Lachenbruch and Marshall (1986). This has lead to many local, regional, and global studies (e.g., Huang et al., 2000; Harris and Chapman, 2001; Gosselin and Mareschal, 2003; Beltrami and Bourlon, 2004; Pollack and Smerdon, 2004; Chouinard et al., 2007; Pickler et al., 2016).

Because of the availability of suitable temperature-depth profiles in the Canadian Shield, many studies have been undertaken in central and eastern Canada . The majority of these studies have used temperature-depth profiles from the southern portion of the Superior Province of the Canadian Shield ($\sim 45^{\circ}$ - 50° N), where many mining exploration holes are readily available and the crystalline rocks are less likely to be affected by groundwater flow than sedimentary rocks. These studies have shown a warming signal of ~ 1 - 2 K over the last ~ 150 - 200 years following a period of cooling about 200-500 years BP associated with the little ice age (LIA) (e.g., Beltrami and Mareschal, 1992; Wang et al., 1992; Guillou-Frottier et al., 1998; Gosselin and Mareschal, 2003; Chouinard and Mareschal, 2007) .

For logistical reasons, mining exploration has been restricted to the southernmost part of the Shield and the few holes that have been drilled in northern regions cannot be measured because they are blocked by permafrost. Nevertheless, a few studies were conducted at higher latitudes. Majorowicz et al. (2004) reconstructed the GST history for 61 temperature-depth profiles between 60° N and 82° N in northern Canada. They found strong evidence that GST warming started in the late 18th century and continued until present. Simultaneous inversion of their data yielded a warming of ~ 2 K for the last 500 years. Studies in Ellesmere Island (above 60° N) have shown varying trends, confirming that temperatures do not increase uniformly over Arctic regions. Taylor et al. (2006) reconstructed the 500 year GST history from three boreholes and found a 3 K warming since the LIA minimum, ~ 200 yrs BP, which is consistent with Beltrami and Taylor (1995) results and the oxygen isotopes studies on ice cores from the region (Fisher and Koerner, 1994). Chouinard et al. (2007) used three temperature-depth profiles in a region with continuous permafrost at the northernmost tip of Quebec to infer the GST history. They found a very strong and recent warming of ~ 2.5 K, with the largest part of this warming occurring in the preceeding 15 yrs, i.e. much later than in Ellesmere Island. Because of lack of adequate borehole temperature depth profiles in eastern Canada between 51° N and 60° N, the large region between the Canadian Arctic and the southern part of the Canadian Shield has not been studied and the climate trends of the last 500 years for this region remain unclear except for boreholes at Voisey Bay, at 56° N on the east coast of Labrador, which show almost no climate signal (Mareschal et al., 2000).

The first motivation of this study is to reduce the gap in data between the Arctic and southeastern Canada. We shall examine 18 temperature-depth profiles measured at 10 sites from eastern Canada to reconstruct the GST histories for the last 500 years. The sites are located in the poorly sampled region north of 51°N, west and east of James Bay in northern Ontario and Quebec. They are to the north of the previous eastern Canada studies and south of the Arctic ones, in a part of the Superior Province where heat flux is extremely low ($< 30\text{mW m}^{-2}$) (Jaupart et al., 2014).

The second motivation of the study is to assess whether borehole temperature profiles can be used to retrace the evolution of permafrost in northern Ontario and Quebec. Permafrost maps locate the Ontario sites in a region classified as with extensive discontinuous permafrost, i.e. where permafrost affects between 50-90% of the ground, while the sites in Quebec are in a region described as with sporadic discontinuous permafrost, i.e. where less than 50% of the ground is frozen. **the boreholes in a region of discontinuous permafrost (Brown et al., 2002). All sites, excluding Noront, lie in discontinuous isolated patches of permafrost, i.e. where less than %10 of the ground is frozen. Noront lies near the southern boundary of extensive discontinuous permafrost, i.e. where permafrost affects between 50-90% of the ground.** In regions with an absence of ground temperature measurements, such as northern Ontario and Quebec, permafrost maps are estimated from surface air temperature and their contour lines (Heginbottom, 2002). The -2.5°C mean annual surface air temperature (SAT) contour line for the period 1950-1980 crosses the southern part of our study region in Ontario, and most of the Quebec sites are located between -2.5 and -5°C SAT contour lines (Phillips, 2002). However, permafrost was not encountered during sampling of the Quebec or Ontario boreholes. It is also worth pointing out that the ground is covered by thick snow cover during several months (from mid December to late April) in the regions above 50°N . Studies demonstrated that the ground surface temperatures are strongly affected by the duration of the snow cover and are offset from SAT (Bartlett et al., 2005; Zhang, 2005; González-Rouco et al., 2006, 2009; García-García et al., 2016). In these regions with extensive snow cover, the borehole temperature profiles are affected by changes in both SAT and snow cover. Meteorological and proxy data indicate that there is more snowfall and longer snow cover on the ground in the Quebec region than in Ontario (Bégin, 2000; Brown and Mote, 2009; Brown, 2010; Environment Canada, 2010; Nicault et al., 2014). This points to possibly warmer present ground surface temperatures and smaller permafrost extent in northern Quebec than in Ontario, and the prospect for different ground surface temperature histories between the regions.

2 Theory

Assuming Earth is a half-space where physical properties only vary with depth, the temperature at depth z , $T(z)$, can be written as (Jaupart and Mareschal, 2011):

$$T(z) = T_o + Q_o R(z) - \int_0^z \frac{dz'}{\lambda(z')} \int_0^{z'} H(z'') dz'' + T_t(z) \quad (1)$$

where T_o is the reference surface temperature, Q_o the reference surface heat flux, the integral accounts for the vertical distribution of heat producing elements $H(z)$, and $T_t(z)$ is the temperature perturbation at depth z due to time-varying changes to the surface boundary condition. The thermal depth $R(z)$ is defined as:

$$R(z) = \int_0^z \frac{dz'}{\lambda(z')} \quad (2)$$

5 where λ is the thermal conductivity.

The temperature perturbation can be calculated by the following equation (Carslaw and Jaeger, 1959):

$$T_t(z) = \int_0^\infty \frac{z}{2\sqrt{\pi\kappa t^3}} \exp\left(-\frac{z^2}{4\kappa t}\right) T_o(t) dt \quad (3)$$

where κ is thermal diffusivity and $T_o(t)$ is the surface temperature at time t before present. For a step change in surface temperature, ΔT , at time t before present, the temperature perturbation $T_t(z)$ is given by Carslaw and Jaeger (1959):

$$10 \quad T_t(z) = \Delta T \operatorname{erfc}\left(\frac{z}{2\sqrt{\kappa t}}\right) \quad (4)$$

where erfc is the complementary error function. If the GST perturbations are approximated by their mean values ΔT_k during K intervals (t_{k-1}, t_k) , the temperature perturbation is written as follows:

$$T_t(z) = \sum_{k=1}^K \Delta T_k \left(\operatorname{erfc}\frac{z}{2\sqrt{\kappa t_k}} - \operatorname{erfc}\frac{z}{2\sqrt{\kappa t_{k-1}}} \right) \quad (5)$$

ΔT_k is the average difference between the ground surface temperature during the time interval (t_{k-1}, t_k) and the reference
15 surface temperature T_0 .

2.1 Inversion

To reconstruct the GST history for each temperature-depth profile, we must invert equation 5. The inversion involves solving for the parameters T_o , Q_o , and ΔT_k of the temperature-depth profile. Equation 5 yields a system of linear equations in the unknown parameters for each depth where temperature has been measured. If N temperature measurements were made in the
20 borehole, a system of N linear equations with $K + 2$ unknowns, T_o , Q_o , and the K values of ΔT_k is obtained. However this system of equations is ill-conditioned and its solution is unstable to small perturbations in the temperature data, i.e., a small error in the data results in a very large error in the solution (Lanczos, 1961). Different inversion methods are available to stabilize (regularize) the solution of ill posed problems (Backus-Gilbert method, Tikhonov regularization algorithm, Bayesian methods, singular value decomposition, Monte-Carlo methods). All these inversion techniques have been applied to reconstruct the GST
25 history (e.g., Vasseur et al., 1983; Nielsen and Beck, 1989; Shen and Beck, 1991; Mareschal and Beltrami, 1992; Clauser and

Mareschal, 1995; Mareschal et al., 1999). In this paper, we have used the singular value decomposition because it is a very simple method to reduce the impact of noise and errors on the solution (Lanczos, 1961). This technique is well documented for geophysical studies (Jackson, 1972; Menke, 1989) and its application for inversion of the ground temperature history is straightforward (Mareschal and Beltrami, 1992).

5

For sites including several boreholes with similar surface conditions, the data are inverted simultaneously because it is assumed that they have experienced the same surface temperature variations and therefore consistent subsurface temperature anomalies. It was expected that consistent trends in the temperature profiles would reinforce each other while errors and random noise would cancel each other. However, the resulting improvement in the signal to noise ratio remains marginal unless a sufficiently large number of profiles with the same GST history are available, which is almost never the case. Simultaneous inversion is described in detail and discussed by Beltrami and Mareschal (1992), Clauser and Mareschal (1995), and Beltrami et al. (1997), among others.

3 Description of data

Figure 1 shows the locations of the thirteen sites including twenty-five boreholes across northern Ontario and Quebec. The heat flow of these sites has previously been studied and a detailed description of the measurement techniques and sites can be found in the heat flow publications (Jessop, 1968; Jessop and Lewis, 1978; Lévy et al., 2010; Jaupart et al., 2014). All the sites are located north of 51°N, west and east of James Bay, and the boreholes range in depth between 400 and 800 m. All the holes were cased in the upper ~10-15 m and, excluding ~~except~~ Otoskwin and Nielsen Island, were drilled for mining exploration purposes. The temperature was measured at 10 m intervals using a calibrated thermistor. The sampling rate is higher at Otoskwin with temperature measurements every 1 m, while Nielsen Island was measured every 30 m. The overall accuracy is estimated on the order of 0.02 K with a precision of greater than 0.005 K. Thermal conductivity was measured on core samples by the method of divided bars (Misener and Beck, 1960). Radiogenic heat production measurements were also made on core samples but are not needed for corrections because the holes are not deep and the heat production rate is low.

Only eighteen holes proved suitable for inversion of the ground surface temperature. Their location and depth can be found in Table 1. ~~Four~~ **Three northern Ontario** sites are located in ~~in northern Ontario~~ in a region of ~~extensive~~ discontinuous **isolated patches of** permafrost: Musselwhite, Thierry Mine, and Otoskwin, ~~and Noront~~. **One site, Noront, lies near the southern edge of a region with extensive discontinuous permafrost.** Thierry Mine (0605, 0606, 0608) and Noront (1012, 1013, 1014, 1015) include several boreholes. Six sites are located in northern Quebec in a region of ~~sporadic~~ discontinuous **isolated patches of** permafrost (Nielsen Island, LaGrande, Eastmain, Eleonore, Corvet, Camp Coulon), with Eastmain (0803, 0804) and Camp Coulon (0712, 0713, 0714) having multiple boreholes. Systematic variations in thermal conductivity observed at Nielsen Island were corrected by using the thermal depth (Bullard, 1939). Some measurements were not used for this study for different reasons (Table 2). **Boreholes less than 300 m deep were rejected for being too shallow. When the mean distance to a lake was**

less than the depth of the hole, or less than 300 m, they were rejected. Furthermore, boreholes were deemed too steep and rejected if they had a slope of 5% or more over a distance comparable to the depth. The profile at Miminiska Lake (ONT) is too shallow to be inverted; the boreholes at Clearwater (QC) are plunging under a lake that affects the temperature profiles; the borehole at Poste Lemoyne is on the side of a very steep hill and the profile is seriously perturbed by the topography. We also discarded one of the temperature profiles at the Eleonore site because the borehole was plunging under a recently filled water reservoir and one of the profiles at the LaGrande site because it was a few m away from the edge of a 30 m cliff. The borehole temperature-depth profiles at sites with multiple boreholes were truncated at the depth of the shallowest borehole to ensure that the same period of time was being studied (Thierry Mine at 530 m, Noront at 400 m, Eastmain at 400 m, and Camp Coulon at 400 m) (Beltrami et al., 2011). The temperature anomaly for each site was calculated by subtracting from the data the estimated steady-state temperature obtained by least-square fitting of a linear function to the bottom 100 m of the profile (Figures 2-3). Tests were made to show that, below 300 m, the heat flux does not vary with the selected depth interval (Lévy et al., 2010; Jaupart et al., 2014) and that the reference temperature profile is stable.

4 Results

The temperature-depth profiles from the ten sites were inverted to reconstruct the GST histories for the last 500 years divided in intervals of 20 years (Figures 4-6). Simultaneous inversion was used at sites with multiple boreholes, Thierry Mine, Noront, Eastmain, and Camp Coulon. The cutoff value or number of eigenvalues determines which part of the solution is eliminated to reduce the impact of noise. A lower cutoff value results in higher resolution in the reconstruction of the GST but at the expense of stability (Mareschal and Beltrami, 1992). Three eigenvalues (0.2 cutoff) were retained for all the sites except Ootskwin and Corvet, where four eigenvalues (0.08 cutoff) were retained. The results of the inversions are summarized in Table 3. Although the ground surface temperature histories differ in their details, they consistently show a trend of warming relative to the reference temperature (i.e. temperature 500 yrs before logging). Only one site shows indications that the GST was affected by the LIA, a cold period that occurred between 200-500 yrs BP.

In northern Ontario, the trends of the inferred GST differ between sites. Ootskwin is the only site to show a LIA signal, with a cooling of ~ 0.5 K with respect to the reference temperature (500 yrs BP). Evidence for this cooling can be found in the temperature anomaly at ~ 200 m but is also observed at Musselwhite and TM0608, where no LIA is reconstructed (Figure 2). Moreover, there is a noticeable change in the Ootskwin temperature gradient at ~ 200 m, which cannot be correlated to variations in thermal conductivity measured on 80 samples from the borehole. The recent warming at Musselwhite and Ootskwin occurred around the same time but it is observed earlier (~ 250 -300 yrs BP) at Thierry Mine and Noront (Figure 4). The total amplitude of warming differs greatly between the sites: 0.50 K with respect to reference temperature at Ootskwin, 0.88 K at Musselwhite, 1.85 K at Noront and 2.85 K at Thierry Mine. It is likely that the Thierry Mine signal was amplified by the clearing of vegetation that took place during the operation of the mine between 1934 and 1950.

Unlike for northern Ontario, a LIA signal was not found for any of the northern Quebec sites (Figures 5-6). A LIA signal was expected because pollen data have suggested that the cooling during the LIA (up to -0.3°C for North America) was strongest in northern Quebec (Gajewski, 1988; Viau and Gajewski, 2009; Viau, 2012). However, a cooling signal at ~ 200 m, which could be associated with the LIA, is observed in the temperature anomaly of CC0713 (Figure 3). Discontinuities are observed in the temperature anomaly of CC0712 between 100 and 300 m. These are also observed in the temperature gradient and could be due to small water flows. The onset of the recent climate warming is the same for all the sites (~ 100 -150 yrs BP), except Eleonore, where it began ~ 200 -300 yrs BP. The amplitude of the warming varies between 0.5-2 K (Figures 5-6) with the largest warming occurring at Corvet (2.18 K).

5 Discussion and Conclusions

Borehole temperature profiles in northern Ontario and Quebec consistently show a ground surface temperature increase of $\sim 1-2.151 \pm 0.76$ K above the reference temperature. Most of this increase took place during the last 150 yrs for the period of 1850 to 2000. Two sites, Thierry Mine and Corvet appear to have large has a shows larger than average warming signals, 2.85 K and 2.18 K, respectively. The area around the Thierry Mine boreholes (0605, 0606, 0608) was cleared in the 1940s after the first opening of the mine and a satellite image locates all three boreholes ~ 300 m from a lake. Lakes disturb a profile if they are at a distance less than the depth of the boreholes (Lewis and Wang, 1992). The proximity to the lake along with the change in vegetation cover could explain the enhanced warming signal (Lewis and Wang, 1998; Lewis, 1998). Corvet is located on the side of a 30 m hill. It has been shown that topography distorts the temperature isotherms: a positive topography leads to a reduced temperature gradient and an increased apparent warming signal. These examples further This illustrates the significant influence of non-climatic effects on ground surface temperature reconstructions from borehole temperature-depth profiles.

A cooling period corresponding to the LIA was found for only one site, Otokwin (ON), which exhibits marked perturbations of the temperature profile. While spatial and temporal variation in the LIA have been noted (Matthews and Briffa, 2005), the absence of a consistent LIA signal in northern Ontario and Quebec deserves some discussion. The LIA cooling period has been inferred from different proxies and selected borehole temperature-depth profiles in eastern Canada (e.g., Archambault and Bergeron, 1992; Beltrami and Mareschal, 1992; Wang and Lewis, 1992; Chouinard et al., 2007; Bunbury et al., 2012). For example, pollen data indicate a pronounced LIA cooling in Quebec (Viau and Gajewski, 2009; Gajewski, 1988). The lack of LIA signal in the majority of the borehole inversions could be related to a combination of several factors. One is the limited resolution of the inversion of borehole temperature profiles. In the presence of noise, a period of weak cooling between 500 and 200 yrs B.P. followed by strong warming is difficult to resolve. Resolution at Otokwin is better because the singular value cutoff was lowest at this site. Furthermore, Mareschal and Beltrami (1992) showed that resolution decreases when noise and errors must be filtered and a higher singular value cutoff required to reduce the impact of noise but retain the gross features of the solution. Profiles and anomalies of Musselwhite and TM0608 are noisy, this could explain the absence of LIA signal in the reconstructions. However, the CC0713 temperature anomaly is less noisy and shows a mild cooling of $\leq 0.2\text{K}$. A test run with

1 K cooling between 1600 and 1800 and varying singular value cutoff showed that, in noise-free synthetic data, a 1 K cooling cannot be resolved with less than 5 singular values. This explains why a mild cooling, such as that observed at CC0713 (≤ 0.2 K) and in pollen data (≤ 0.3 K) (Gajewski, 1988; Viau and Gajewski, 2009; Viau, 2012), could not be resolved. Also, Chouinard and Mareschal (2007) suggested that the LIA could have started ~ 100 yrs earlier in northern Quebec than in southern Canada.

5 Resolving the LIA would require a borehole deeper than ~ 400 m, which is not the case of all the holes. Let us also point out that the sampling resolution at Nielsen Island is low, with measurements only every 30 m, which is not sufficient to resolve a LIA signal. While the absence of LIA signal in Quebec was unexpected, its absence in northern Ontario confirms the findings of Gosselin and Mareschal (2003), who found only 2 sites with a LIA signal among 33 temperature-depth profiles from north-western Ontario. They hypothesized that the lack of LIA signal could be due to the influence of Lake Superior because the

10 two sites with LIA signals were above 50°N and the furthest from Lake Superior. This is not supported by the present study because the 4 Ontario sites are several hundreds of kilometres away from Lake Superior. It is also possible that the LIA signal is masked by other physical effects, such as an advance and retreat of permafrost or a change in the precipitation regime and the duration of the ground snow cover during the LIA.

15 No geographic trends in the GST histories were observed, despite different SAT conditions. Meteorological data from the NOAA weekly dataset and 8 general circulation models (GCMs) for the period of 1970-1999 display a longer snow cover duration in northern Quebec than in northern Ontario (Brown and Mote, 2009). The higher precipitation is confirmed by proxy reconstructions of lake levels and tree forms (Bégin, 2000; Lavoie and Payette, 1992). Because of the greater snowfall and longer snow cover, the present ground surface relative to air surface temperatures in Quebec are warmer than in Ontario.

20 However these dissimilar conditions have not resulted in noticeable discordance between the GST histories between northern Ontario and Quebec, suggesting that the same differences in precipitation persisted throughout the period reconstructed.

The magnitude of the recent warming is about the same as the $\sim 1\text{-}2$ K warming for the ~~past 150 years~~ **period of 1850 to 2000** inferred from several studies in the southern portion of the Superior Province (Beltrami and Mareschal, 1992; Shen and Beck, 1992; Chouinard and Mareschal, 2007) and less than the very pronounced warming in the eastern Canadian Arctic (Beltrami and Taylor, 1995; Taylor et al., 2006; Chouinard et al., 2007).

The sites are located in a region described as discontinuous permafrost, where ground temperatures are slightly below freezing, at least according to the Canadian and world permafrost maps. No sign of permafrost was found at any of the measured

30 sites nor at the sites that were excluded (see Table 2), **however, permafrost has been reported in the James Bay lowlands near Eleonore, Eastmain, and Lagrande (Thibault and Payette, 2009)**. Not only are the present average ground surface temperatures well above the freezing point of water, but also, except for Nielsen Island, the ground surface temperature histories retrieved from inversion reveal that the temperature has remained well above the freezing point for the last 500 years, **indicating that the potential for permafrost was minimal to absent. We have also noted that** during logging of more than 100 holes in the

35 regions of Manitoba, Saskatchewan, and northern Ontario classified as ~~extensive~~ discontinuous permafrost, permafrost has been

encountered at only one hole, north of the town of Lynn Lake in northern Manitoba (Guillou-Frottier et al., 1998). Clearly, the spatial distribution of permafrost outlined in the available permafrost maps is ~~inaccurate~~ **questionable** and thus ~~unreliable~~ possibly because they are not based on sufficiently deep ground temperature measurements but estimated from interpolated sparse records of SAT (Heginbottom, 2002; Gruber, 2012). The discrepancy between permafrost maps and direct field observations reveal that SAT interpolations are unsuitable to estimate the spatial variations of ground temperatures. This is likely because the maximum thickness of snow exceeds 1 m at the end of the winter remaining on the ground from mid December to mid April, resulting in a large offset between the GST and SAT (Zhang, 2005; Grosse et al., 2016). Our study suggests that borehole temperature profiles could be used in the future to assess the reality of the permafrost retreat assumed to have occurred after the LIA (Halsey et al., 1995; Schuur et al., 2008). Furthermore, borehole temperature profiles might be a better means for determining the southern extent of areas of past and present permafrost than current permafrost maps and a useful tool for validation of climate models.

15 **Appendix A: Detailed Site Description**

Ten sites, including eighteen boreholes, were utilized to reconstruct the ground surface temperature history of northern Ontario and Quebec for the past 500 years. A detailed description of the rock type and geological unit of each site can be found in Table A1 and heat flow studies of the region (Jessop, 1968; Jessop and Lewis, 1978; Lévy et al., 2010; Jaupart et al., 2014).

20 Four sites (Musselwhite, Thierry Mine, Otokwin, Noront) comprising nine boreholes are located in northern Ontario. The Musselwhite (0601) site is located in a clearing of ~60 m in diameter with a lake ~330 m to the west and ~180 m to the east. Three boreholes (0605, 0606, 0608) are located at Thierry Mine, ~500 m away from a large clearing. This clearing is associated with the development of the nearby mine in 1934-1950. Furthermore, all three sites are ~300 m from a lake. 0605 and 0608 are found in a clearing due to drilling of ~80 m in diameter. The Otokwin borehole was measured in 1985 and is
25 located ~180 m from the Otokwin River. The final four Ontario boreholes (1012, 1013, 1014, 1015) are found in Noront. The sites are fairly flat, swampy and muddy. They are in the McFaulds Lake project, ~300 km north of the town of Geraldton and within a region referred to as the Ring of Fire of the James Bay Lowlands. N1012 and N1013 are ~40 m apart, ~500 from N1015 and ~400 m from N1014. The complex lithological column has led to noisy temperature-depth profiles (Jaupart et al., 2014).

30

Northern Quebec is home to the remaining nine boreholes, found in six distinct sites. Nielsen Island is the northern most site of this study. It was logged in 1977 and is located on an island in the Hudson Bay. The LaGrande borehole (0405) is in a large clearing for drilling, in fairly flat and swampy region. It is located ~400 m from a main Hydro Quebec power line

and power line clearing. Eleonore (0502) is ~ 200 m from the Opinaco reservoir and is dipping towards the reservoir. Two boreholes (0803, 0804) are in Eastmain, ~ 220 km from the mining camp of Matagami. 0803 is ~ 50 m south of a road but is dipping away from the road. Corvet (0716) is located in a fairly flat area. The final three boreholes (0712, 0713, 0714) are in Camp Coulon. 0713 is ~ 20 km north of the hydroelectric station of Laforge Deux. 0712 and 0714 are ~ 450 m apart and
5 located on top of a small relief, with 0712 found in a clearing of ~ 30 m diameter.

Acknowledgements. The authors are grateful to Christian Chouinard, Claude Jaupart and our colleagues at the Institut de Physique du Globe de Paris who helped with field measurements and core sample analysis. We appreciate the useful and constructive comments from two reviewers, an anonymous reviewer and A.E.Taylor, R.Way and Editor A.Winguth. This work was supported by grants from the Natural Sciences and Engineering Research Council of Canada Discovery Grant (NSERC DG 140576948) and the Canada Research Program (CRC
10 230687) to H. Beltrami. Computational facilities provided by the Atlantic Computational Excellence Network (ACENnet-Compute Canada) with support from the Canadian Foundation for Innovation. H. Beltrami holds a Canada Research Chair in Climate Dynamics. C.Pickler is funded by graduate fellowships from a NSERC CREATE Training Program in Climate Sciences based at St.Francis Xavier University.

References

- Archambault, S. and Bergeron, Y.: An 802-year tree-ring chronology from the Quebec boreal forest, *Canadian Journal of Forest Research*, 22, 674–682, doi:10.1139/x92-090, 1992.
- Bartlett, M. G., Chapman, D. S., and Harris, R. N.: Snow effect on North American ground temperatures, 1950–2002, *Journal of Geophysical Research: Earth Surface* (2003–2012), 110, doi:10.1029/2005JF000293, 2005.
- Beck, A.: Climatically perturbed temperature gradient and their effect on regional and continental heat-flow means, *Tectonophysics*, 41, 17–39, doi:10.1016/0040-1951(77)90178-0, 1977.
- Bégin, Y.: Reconstruction of subarctic lake levels over past centuries using tree rings, *Journal of Cold Regions Engineering*, 14, 192–212, doi:10.1061/(ASCE)0887-381X(2000)14:4(192), 2000.
- 10 Beltrami, H. and Bourlon, E.: Ground warming patterns in the Northern Hemisphere during the last five centuries, *Earth and Planetary Science Letters*, 227, 169–177, doi:10.1016/j.epsl.2004.09.014, 2004.
- Beltrami, H. and Mareschal, J.-C.: Ground temperature histories for central and eastern Canada from geothermal measurements: Little Ice Age signature, *Geophysical research letters*, 19, 689–692, doi:10.1029/92GL00671, 1992.
- Beltrami, H. and Mareschal, J.-C.: Resolution of ground temperature histories inverted from borehole temperature data, *Global and Planetary Change*, 11, 57–70, doi:10.1016/0921-8181(95)00002-9, 1995.
- 15 Beltrami, H. and Taylor, A. E.: Records of climatic change in the Canadian Arctic: towards calibrating oxygen isotope data with geothermal data, *Global and Planetary Change*, 11, 127–138, doi:10.1016/0921-8181(95)00006-2, 1995.
- Beltrami, H., Cheng, L., and Mareschal, J. C.: Simultaneous inversion of borehole temperature data for determination of ground surface temperature history, *Geophysical Journal International*, 129, 311–318, doi:10.1111/j.1365-246X.1997.tb01584.x, 1997.
- 20 Beltrami, H., Smerdon, J., G.S.Matharoo, and N.Nickerson: Impact of maximum borehole depths on inverted temperature histories in borehole paleoclimatology, *Climate of the Past*, 7, 715–748, doi:10.5194/cp-7-745-2011, 2011.
- Birch, A. F.: The effects of Pleistocene climatic variations upon geothermal gradients, *American Journal of Science*, 246, 729–760, doi:10.2475/ajs.246.12.729, 1948.
- Brown, J., Ferrians, O., Heginbottom, J. A., and Melnikov, E.: Circum-Arctic Map of Permafrost and Ground-Ice Conditions, Version 2., 25 Map, National Snow and Ice Data Center, Boulder (CO), 2002.
- Brown, R. D.: Analysis of snow cover variability and change in Québec, 1948–2005, *Hydrological Processes*, 24, 1929–1954, doi:10.1002/hyp.7565, 2010.
- Brown, R. D. and Mote, P. W.: The response of Northern Hemisphere snow cover to a changing climate, *Journal of Climate*, 22, 2124–2145, doi:10.1175/2008JCLI2665.1, 2009.
- 30 Bullard, E.: Heat Flow in South Africa, *Proceedings of the Royal Society London A*, 173, 474–502, 1939.
- Bunbury, J., Finkelstein, S. A., and Bollmann, J.: Holocene hydro-climatic change and effects on carbon accumulation inferred from a peat bog in the Attawapiskat River watershed, Hudson Bay Lowlands, Canada, *Quaternary Research*, 78, 275–284, doi:10.1016/j.yqres.2012.05.013, 2012.
- Carslaw, H. and Jaeger, J.: *Conduction of Heat in Solids*, Oxford Science Publications, New York, 1959.
- 35 Cermak, V.: Underground temperature and inferred climatic temperature of the past millennium, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 10, 1–19, doi:10.1016/0031-0182(71)90043-5, 1971.

- Chouinard, C. and Mareschal, J.-C.: Selection of borehole temperature depth profiles for regional climate reconstructions, *Climate of the Past*, 3, 297–313, doi:10.5194/cp-3-297-2007, 2007.
- Chouinard, C., Fortier, R., and Mareschal, J.-C.: Recent climate variations in the subarctic inferred from three borehole temperature profiles in northern Quebec, Canada, *Earth and Planetary Science Letters*, 263, 355–369, doi:10.1016/j.epsl.2007.09.017, 2007.
- 5 Clauser, C. and Mareschal, J.-C.: Ground temperature history in central Europe from borehole temperature data, *Geophysical Journal International*, 121, 805–817, doi:10.1111/j.1365-246X.1995.tb06440.x, 1995.
- Environment Canada: Canadian climate normals or averages 1981-2010, National Climate Data and Information Archive., 2010.
- Fisher, D. A. and Koerner, R. M.: Signal and noise in four ice-core records from the Agassiz Ice Cap, Ellesmere Island, Canada: details of the last millennium for stable isotopes, melt and solid conductivity, *The Holocene*, 4, 113–120, doi:10.1177/095968369400400201, 1994.
- 10 Gajewski, K.: Late Holocene climate changes in eastern North America estimated from pollen data, *Quaternary Research*, 29, 255–262, doi:10.1016/0033-5894(88)90034-8, 1988.
- García-García, A., Cuesta-Valero, F., Beltrami, H., and Smerdon, J.: Simulation of Air and Ground Temperatures in PIMIP3/CMIP5 Last Millenium Simulation: Implications for Climate Reconstructions from Borehole Temperature Profiles, *Environmental Research Letters*, 11, doi:10.1088/1748-9326/11/4/044022, 2016.
- 15 González-Rouco, J., Beltrami, H., Zorita, E., and Von Storch, H.: Simulation and inversion of borehole temperature profiles in surrogate climates: Spatial distribution and surface coupling, *Geophysical Research Letters*, 33, doi:10.1029/2005GL024693, 2006.
- González-Rouco, J., Beltrami, H., Zorita, E., and Stevens, M.: Borehole climatology: a discussion based on contributions from climate modeling, *Climate of the Past*, 5, 97–127, doi:10.5194/cp-5-97-2009, 2009.
- Gosselin, C. and Mareschal, J.-C.: Recent warming in northwestern Ontario inferred from borehole temperature profiles, *Journal of Geophysical Research: Solid Earth*, 108, doi:10.1029/2003JB002447, 2003.
- 20 Grosse, G., Goetz, S., McGuire, A. D., Romanovsky, V. E., and Schuur, E. A.: Changing permafrost in a warming world and feedbacks to the Earth system, *Environmental Research Letters*, 11, 040201, doi:10.1088/1748-9326/11/4/040201, 2016.
- Gruber, S.: Derivation and analysis of a high-resolution estimate of global permafrost zonation, *The Cryosphere*, 6, 221–233, doi:10.5194/tc-6-221-2012, 2012.
- 25 Guillou-Frottier, L., Mareschal, J. C., and Musset, J.: Ground surface temperature history in central Canada inferred from 10 selected borehole temperature profiles, *Journal of Geophysical Research*, 103, 7385–7397, doi:10.1029/98JB00021, 1998.
- Halsey, L. A., Vitt, D. H., and Zoltai, S. C.: Disequilibrium response of permafrost in boreal continental western Canada to climate change, *Climatic Change*, 30, 57–73, doi:10.1007/BF01093225, 1995.
- Harris, R. N. and Chapman, D. S.: Mid-latitude (30–60 N) climatic warming inferred by combining borehole temperatures with surface air temperatures, *Geophysical Research Letters*, 28, 747–750, doi:10.1029/2000GL012348, 2001.
- 30 Heginbottom, J. A.: Permafrost mapping: a review, *Progress in Physical Geography*, 26, 623–642, doi:10.1191/0309133302pp355ra, 2002.
- Hotchkiss, W. and Ingersoll, L.: Post-glacial time calculations from recent geothermal measurements in the Calumet Copper Mines, *Journal of Geology*, 42, 113–142, 1934.
- Huang, S., Pollack, H. N., and Shen, P.-Y.: Temperature trends over the past five centuries reconstructed from borehole temperatures, *Nature*, 403, 756–758, doi:10.1038/35001556, 2000.
- 35 Jackson, D.: Interpretation of inaccurate, insufficient, and inconsistent data, *Geophysical Journal International*, 28, 97–109, doi:10.1111/j.1365-246X.1972.tb06115.x, 1972.

- Jaupart, C. and Mareschal, J.-C.: Heat generation and transport in the Earth, Cambridge University Press, Cambridge, United Kingdom, 2011.
- Jaupart, C., Mareschal, J., Bouquerel, H., and Phaneuf, C.: The building and stabilization of an Archean Craton in the Superior Province, Canada, from a heat flow perspective, *Journal of Geophysical Research: Solid Earth*, 119, 9130–9155, doi:10.1002/2014JB011018, 2014.
- 5 Jessop, A. and Lewis, T.: Heat flow and heat generation in the Superior province of the Canadian Shield, *Tectonophysics*, 50, 55–57, doi:10.1016/0040-1951(78)90199-3, 1978.
- Jessop, A. M.: Three measurements of heat flow in eastern Canada, *Canadian Journal of Earth Sciences*, 5, 61–68, doi:10.1139/e68-006, 1968.
- Lachenbruch, A. and Marshall, B.: Changing climate: Geothermal evidence from permafrost in the Alaskan Arctic, *Science*, 234, 689–696, doi:10.1126/science.234.4777.689, 1986.
- 10 Lanczos, C.: *Linear Differential Operators*, D. Van Nostrand, Princeton, N.J., 1961.
- Lavoie, C. and Payette, S.: Black spruce growth forms as a record of a changing winter environment at treeline, Quebec, Canada, *Arctic and Alpine Research*, 24, 40–49, doi:10.2307/1551318, 1992.
- Lévy, F., Jaupart, C., Mareschal, J.-C., Bienfait, G., and Limare, A.: Low heat flux and large variations of lithospheric thickness in the Canadian Shield, *Journal of Geophysical Research: Solid Earth (1978–2012)*, 115, doi:10.1029/2009JB006470, 2010.
- 15 Lewis, T.: The effect of deforestation on ground surface temperatures, *Global and Planetary Change*, 18, 1–13, doi:10.1016/S0921-8181(97)00011-8, 1998.
- Lewis, T. J. and Wang, K.: Influence of terrain on bedrock temperatures, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 98, 87–100, doi:10.1016/0031-0182(92)90190-G, 1992.
- 20 Lewis, T. J. and Wang, K.: Geothermal evidence for deforestation induced warming: Implications for the climatic impact of land development, *Geophysical Research Letters*, 25, 535–538, doi:10.1029/98GL00181, 1998.
- Majorowicz, J. A., Skinner, W. R., and Safanda, J.: Large ground warming in the Canadian Arctic inferred from inversions of temperature logs, *Earth and Planetary Science Letters*, 221, 15–25, doi:10.1016/S0012-821X(04)00106-2, 2004.
- Mareschal, J., Poirier, A., Rolandone, F., Bienfait, G., Gariépy, C., Lapointe, R., and Jaupart, C.: Low mantle heat flow at the edge of the North American continent, Voisey Bay, Labrador, *Geophysical Research Letters*, 27, 823–826, doi:10.1029/1999GL011069, 2000.
- 25 Mareschal, J.-C. and Beltrami, H.: Evidence for recent warming from perturbed geothermal gradients: examples from eastern Canada, *Climate Dynamics*, 6, 135–143, doi:10.1007/BF00193525, 1992.
- Mareschal, J.-C., Rolandone, F., and Bienfait, G.: Heat flow variations in a deep borehole near Sept-Iles, Québec, Canada: Paleoclimatic interpretation and implications for regional heat flow estimates, *Geophysical Research Letters*, 26, 2049–2052, doi:10.1029/1999GL900489, 1999.
- 30 Matthews, J. A. and Briffa, K. R.: The ‘Little Ice Age’: Re-evaluation of an evolving concept, *Geografiska Annaler: Series A, Physical Geography*, 87, 17–36, doi:10.1111/j.0435-3676.2005.00242.x, 2005.
- Menke, W.: *Geophysical Data Analysis: Discrete Inverse Theory*, vol. 4, Academic Press, San Diego, 1989.
- Misener, A. and Beck, A.: *The measurement of heat flow over land*, Interscience, New York, 1960.
- 35 Nicault, A., Boucher, E., Tapsoba, D., Arseneault, D., Berninger, F., Bégin, C., DesGranges, J., Guiot, J., Marion, J., Wicha, S., et al.: Spatial analysis of black spruce (*Picea mariana* (Mill.) BSP) radial growth response to climate in northern Québec–Labrador Peninsula, Canada, *Canadian Journal of Forest Research*, 45, 343–352, doi:10.1139/cjfr-2014-0080, 2014.

- Nielsen, S. and Beck, A.: Heat flow density values and paleoclimate determined from stochastic inversion of four temperature-depth profiles from the Superior Province of the Canadian Shield, *Tectonophysics*, 164, 345 – 359, doi:10.1016/0040-1951(89)90026-7, 1989.
- Phillips, D.: Climatic Atlas of Canada series 1, Tech. rep., Canadian Climate Center, Ottawa (ON), 2002.
- Pickler, C., Beltrami, H., and Mareschal, J.-C.: Laurentide Ice Sheet basal temperatures during the last glacial cycle as inferred from borehole data, *Climate of the Past*, 12, 115–127, doi:10.5194/cp-12-115-2016, 2016.
- 5 Pollack, H. N. and Smerdon, J. E.: Borehole climate reconstructions: Spatial structure and hemispheric averages, *Journal of Geophysical Research: Atmospheres*, 109, doi:10.1029/2004JD005056, 2004.
- Sass, J. H., Lachenbruch, A. H., and Jessop, A. M.: Uniform heat flow in a deep hole in the Canadian Shield and its paleoclimatic implications, *Journal of Geophysical Research*, 76, 8586–8596, doi:10.1029/JB076i035p08586, 1971.
- 10 Schuur, E. A., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V., Hagemann, S., Kuhry, P., Lafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A., Romanovsky, V., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G., and Zimov, S. A.: Vulnerability of permafrost carbon to climate change: implications for the global carbon cycle, *BioScience*, 58, 701–714, doi:10.1641/B580807, 2008.
- Shen, P. and Beck, A.: Paleoclimate change and heat flow density inferred from temperature data in the Superior Province of the Canadian Shield, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 98, 143–165, doi:10.1016/0031-0182(92)90194-A, 1992.
- 15 Shen, P. Y. and Beck, A. E.: Least squares inversion of borehole temperature measurements in functional space, *Journal of Geophysical Research: Solid Earth*, 96, 19965–19979, doi:10.1029/91JB01883, 1991.
- Taylor, A., Wang, K., Smith, S., Burgess, M., and Judge, A.: Canadian Arctic Permafrost Observatories: detecting contemporary climate change through inversion of subsurface temperature time series, *Journal of Geophysical Research*, 111, doi:10.1029/2004JB003208, 2006.
- 20 Thibault, S. and Payette, S.: Recent permafrost degradation in bogs of the James Bay area, northern Quebec, Canada, *Permafrost and Periglacial Processes*, 20, 383–389, doi:10.1002/ppp.660, 2009.
- Vasseur, G., Bernard, P., de Meulebrouck, J. V., Kast, Y., and Jolivet, J.: Holocene paleotemperatures deduced from geothermal measurements, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 43, 237 – 259, doi:10.1016/0031-0182(83)90013-5, 1983.
- Viau, A. and Gajewski, K.: Reconstructing millennial-scale, regional paleoclimates of boreal Canada during the Holocene, *Journal of Climate*, 22, 316–330, doi:10.1175/2008JCLI2342.1, 2009.
- 25 Viau, A. E.: Climate of the last 2,000 years in North America reconstructed from pollen records, *Quaternary International*, 279, 520, doi:10.1016/j.quaint.2012.08.1800, 2012.
- Wang, K. and Lewis, T. J.: Geothermal evidence from Canada for a cold period before recent climatic warming, *Science*, 256, 1003–1005, doi:10.1126/science.256.5059.1003, 1992.
- 30 Wang, K., Lewis, T. J., and Jessop, A. M.: Climatic changes in central and eastern Canada inferred from deep borehole temperature data, *Global and planetary change*, 6, 129–141, doi:10.1016/0921-8181(92)90031-5, 1992.
- Zhang, T.: Influence of the seasonal snow cover on the ground thermal regime: An overview, *Reviews of Geophysics*, 43, doi:10.1029/2004RG000157, 2005.




Figure 1. Map of Ontario and western Quebec showing the location of sites (red dots). For sites with several boreholes (Camp Coulon , Eastmain, Thierry Mine, and Noront), the number of profiles available is enclosed in parenthesis. Black diamonds show the locations of sites that were discarded.




Figure 2. Temperature anomalies for the northern Ontario boreholes. Holes TM0605, TM0606, and TM0608 are from the Thierry Mine site; holes N1012, N1013, N1014, and N1015 belong to the Noront site. The anomaly is obtained by subtracting the estimated steady-state geotherm obtained by the least-square fit of a straight line to the bottom 100 m of the borehole temperature-depth profile. The black line represents the best linear fit, while the pink and blue lines are the upper and lower bounds, respectively, of the 2σ confidence intervals. For N1014, TM0606, and Otokwin, the upper and lower bounds of the confidence interval are not visible due to the temperature scale. The temperature anomaly at Musselwhite was cut at 650 m.




Figure 3. Temperature anomalies for the northern Quebec boreholes. CC0712, CC0713, CC0714 are the boreholes from Camp Coulon; Ea0803 and Ea0804 are the boreholes from Eastmain . The anomaly is obtained by subtracting the estimated steady-state geotherm obtained by the least-square fit of a straight line to the bottom 100 m of the borehole temperature-depth profile. The black line represents the best linear fit, while the pink and blue lines represent the upper and lower bounds, respectively, of the 2σ confidence intervals. For N1014, TM0606, and Otokwin, the upper and lower bounds of the confidence interval are not visible due to the temperature scale. The temperature anomaly at Nielsen Island was cut at 600m.




Figure 4. GST histories for the northern Ontario sites determined by inversion of the anomalies. For multiple holes at a given site (Thierry mine and Noront), simultaneous inversion was used. The pink and blue lines represent the inversions of the upper and lower bounds of the anomaly. For Otokwin, the three lines are superposed.




Figure 5. GST histories for the northern Quebec sites. Simultaneous inversion was used for Eastmain, which includes two holes. The pink and blue lines represent the inversions of the upper and lower bounds of the anomaly.




Figure 6. GST histories for the northern Quebec sites. Simultaneous inversion was used for Camp Coulon, which includes more than one hole. The pink and blue lines represent the inversions of the upper and lower bounds of the anomaly.

Table 1. Location and technical information concerning the boreholes used in this study, where **true depth** is the depth corrected for the dip of the borehole, λ is the thermal conductivity, Q is the heat flux, and Q_{corr} is the heat flux corrected for post glacial warming

Site	Log ID	Latitude	Longitude	Dip (°)	True depth (m)	λ (W m ⁻¹ K ⁻¹)	Q (mW m ⁻²)	Q_{corr} (mW m ⁻²)	Reference
Musselwhite Thierry Mine	0601	52°37'28.3"	90°23'32.7"	76	740	2.73	30.7	33	Lévy et al. (2010)
							28		
	0605	51°30'24"	90°21'11"	74	802	2.91	26.8	29.7	
	0606	51°30'22"	90°21'11"	70	530	2.61	24.4	27.0	
Otoskwin	0608	51°30'24.3"	90°21'10.7"	62	737	2.60	24.6	27.1	Lévy et al. (2010)
							25		
	n/a	51°49.5'	89°35.9'	90	602	2.79	20	25	Jessop and Lewis (1978)
Noront							35		
	1012	52°44'23.9"	86°17'42.1"	68	761	3.2	34.1	37.6	Jaupart et al. (2014)
	1013	52°44'25"	86°17'42.1"	55	389	3.1	28.3	34.1	Jaupart et al. (2014)
	1014	52°44'29.9"	86°17'59.9"	77	762	3.1	31.7	35.6	Jaupart et al. (2014)
	1015	52°44'25.5"N	86°18'11.9"W	75	806	3.1	29.5	33.4	Jaupart et al. (2014)
Nielsen Island							26		
	0-77	55°23.7'	77°41.0'	90	1408	5.5 (\leq 400m) 2.4 ($>$ 400 m)	-	26.8	Jessop (1968)
LaGrande							22		
	0405	53°31'45"	76°33'15"	60	600	2.89	19.2	21.9	Lévy et al. (2010)
Eastmain							34		
	0803	52°10'16"	76°27'66"	60	398	2.9	28.8	34.1	Jaupart et al. (2014)
	0804	52°10'16"	76°27'66"	60	390	2.9	26.7	32.9	Jaupart et al. (2014)
Eleonore							31		
	0502	52°42'05"	76°04'46"	56	527	2.47	30.7	32.8	Lévy et al. (2010)
Corvet							27		
	0716	53°19.072'N	73°55.760'W	67	479	2.80	24.0	26.8	Lévy et al. (2010)
Camp Coulon							28		
	0712	54°47'43"	71°17'09"	54	561	3.69	26.4	29.0	Lévy et al. (2010)
	0713	54°47'95"	71°17'20"	40	460	3.73	24.5	27.5	Lévy et al. (2010)
	0714	54°47'43"	71°17'34"	56	384	-	-	-	Lévy et al. (2010)

Table 2. Location and technical information concerning boreholes not suitable for this study, where T_o is the reference surface temperature and Q_o is the reference heat flux

Site	Log ID	Latitude	Longitude	T_o (°C)	Q_o (mW m ⁻²)	Remark	Reference
Miminiska	0603	51°34'51"	88°31'09"	3.33±0.02	25.8±1.5	Too shallow	(Lévy et al., 2010)
LaGrande	0406	53°31'42"	76°33'49"	2.67±0.005	14.1±0.3	Topography	(Lévy et al., 2010)
Eleonore	0503	52°42'00"	76°04'45"	2.56±0.01	27.7±0.6	Reservoir	(Lévy et al., 2010)
Clearwater	0505	52°12'33"	75°48'38"	2.33±0.02	30.6±0.4	Lake	(Lévy et al., 2010)
	0506	52°12'31"	75°48'23"	2.23±0.01	31.3±0.2	Lake	(Lévy et al., 2010)
	0507	52°12'39"	75°48'23"	2.73±0.004	30.5±0.3	Lake	(Lévy et al., 2010)
Poste Lemoyne	0715	53°27'37"	75°12'21"	1.83±0.004	24.5±0.2	Topography	(Lévy et al., 2010)

Table 3. Summary GST History Results where T_o is the reference surface temperature, Q_o is the reference heat flux, ΔT is the difference between the maximal temperature and the reference temperature 500 yrs before logging.

Site	Log ID	year	T_o (°C)	Q_o (mW m ⁻²)	ΔT (K)	LIA
Musselwhite Thierry Mine	0601	2006	2.56±0.01	30.0±0.5	0.88	no
					2.85	no
	0605	2006	2.63±0.01	25.9±0.5		
	0606	2006	2.55±0.01	23.5±0.7		
	0608	2006	2.62±0.01	23.9±0.3		
Otoskwin	n/a	1985	2.81±0.001	19.5±0.05	0.50	yes
Noront					1.85	no
	1012	2010	1.16±0.01	35.5±0.9		
	1013	2010	1.51±0.02	31.3±1.5		
	1014	2010	1.78±0.004	32.2±0.3		
	1015	2010	2.35±0.02	27.6±1.4		
Nielsen Island	0-77	1977	-0.43±0.02	23.7±0.4	1.35	no
LaGrande	0405	2004	2.18±0.004	19.4±0.2	0.50	no
Eastmain					2.15	no
	0803	2008	2.98±0.01	27.3±0.4		
	0804	2008	3.01±0.003	27.0±0.2		
Eleonore	0502	2005	2.04±0.01	30.9±0.5	1.46	no
Corvet	0716	2007	2.45±0.01	24.9±0.3	2.18	no
Camp Coulon					1.34	no
	0712	2007	1.46±0.01	26.6±0.4		
	0713	2007	1.83±0.01	24.2±0.6		
	0714	2007	1.33±0.004	-		

Table A1. Geological unit and rock type concerning the boreholes used in this study

Site	Log ID	Rock Type	Geological Unit	Reference
Musselwhite	0601	gneiss	Sachigo subprovince	Lévy et al. (2010)
Thierry Mine	0605	granite	Uchi belt	Lévy et al. (2010)
	0606			Lévy et al. (2010)
	0608			Lévy et al. (2010)
Otoskwin	n/a	-	-	Jessop and Lewis (1978)
Noront	1012	-	North Caribou terrane	Jaupart et al. (2014)
	1013			Jaupart et al. (2014)
	1014			Jaupart et al. (2014)
	1015			Jaupart et al. (2014)
Nielsen Island	0-77	-	-	Jessop (1968)
LaGrande	0405	granodiorite	LaGrande volcano-plutonic belt	Lévy et al. (2010)
Eastmain	0803	metasedimentary and volcanics	LaGrande volcano-plutonic belt	Jaupart et al. (2014)
	0804			Jaupart et al. (2014)
Eleonore	0502	wacke	LaGrande volcano-plutonic belt	Lévy et al. (2010)
Corvet	0716	intermediate volcanics	LaGrande volcano-plutonic belt	Lévy et al. (2010)
Camp Coulon	0712	rhyolite	LaGrande volcano-plutonic belt	Lévy et al. (2010)
	0713			Lévy et al. (2010)
	0714			Lévy et al. (2010)