

Actions on comments of reviewer 1-3 for the manuscript ‘North Atlantic Oscillation controls on oxygen and hydrogen isotope gradients in winter precipitation across Europe; implications for palaeoclimate studies’

Reviewer

Authors response and/or action

Actions to the common comments of all reviewers

All three reviewer commented on the stationarity of the NAO through the Holocene.

- **Reviewer 1 states:** I would suggest that the assumption of time-stationarity back to the early Holocene of the wNAOi and d18Opw is possibly correct but currently unsupported. The two sentences around Line 460 do not do this point justice. The manuscript would be stronger by either supporting the assertion with additional evidence that, say, the NAO existed more or less as we know it know during the early Holocene NAO, when the presence of upstream ice sheets and different insolation and vegetation regimes were present, or by providing stronger caveats for extending the NAO discussion prior to the late Holocene.

Revised. We have included recent studies that investigate the stationarity of the NAO during the Holocene and account their results in the discussion and interpretation of the findings of this study. Furthermore, we have extended the discussion on the stationarity of the NAO during the Holocene in general and highlight potential caveats.

Perhaps using the term “NAO-like” instead of the NAO, while emphasizing that early-Holocene climate had distinctly different forcings and boundary conditions than the late Holocene, would be advisable.

Revised.

- **Reviewer 2 states:** I wonder if a spatially stationary gradient can satisfactorily capture the complexity of the NAO. In particular, I suggest that the authors consider the ‘Augmented NAO Index’ of Wang et al., 2012, GRL within the context of their own work. The presumed stationarity of the pressure systems defining the NAO is an issue with most NAO studies, but because this study focuses on a spatial gradient perhaps it should focus more attention on this issue.

We have included the findings of (Wang et al., 2012) in the discussion section of this manuscript and highlight potential caveats that are related to the spatial stationarity.

Would the longitudinal gradient not change orientation through time as pressure centres shift location?

If this would be the case we would expect to see that any reconstructed longitudinal gradient based on speleothem $\delta^{18}\text{O}$ records has a positive slope (towards the east). However, this is not the case as McDermott et al. (2011) have demonstrated for the last 12 ka: instead the reconstruction of Holocene slopes reveals that the slopes of the longitudinal speleothem $\delta^{18}\text{O}$ gradients are steeper in the early Holocene and become progressively shallower until about 5 ka. This result demonstrates that if the pressure centres shift their location, as suggested from 9 ka to 8 ka by the recent study of Wassenburg et al. (2016), that this has only a minor influence on the orientation of the longitudinal gradient.

Particularly in the Early Holocene, where more extensive sea ice would have relocated the pressure centers?

Considering the results of McDermott et al. (2011) and Wassenburg et al. (2016) the shift of the location of the pressure centres is likely to be more gradual during the Holocene until stable environmental boundary conditions are reached. Taking account of the modelling results of Wassenburg et al. (2016) the retreat of the Laurentide ice sheet and a changing meltwater flux into the North Atlantic are responsible for the changes of the NAO (or as stated in Wassenburg et al. (2016) the ‘reorganisation’ of the NAO) and the shift of the pressure systems particularly of the Icelandic low: during the course of the Holocene the simulations would suggest that the Icelandic low system is shifted towards north-east.

- **Reviewer 3 states:** Finally, as the two other reviewers question the stationarity of the NAO and the gradient in the past and, in particular, its dependence on the location of the pressure systems during

the Early Holocene, it may be interesting to read and discuss the recent paper by Wassenburg et al. (2016).

Revised. We have included the aforementioned study of Wassenburg et al. (2016) and have extended the discussion on the stationarity of the NAO during the Holocene.

Actions to the comments of Reviewer 1

General Comments:

I would suggest that the assumption of time-stationarity back to the early Holocene of the wNAOi and d18Opw is possibly correct but currently unsupported. The two sentences around Line 460 do not do this point justice. The manuscript would be stronger by either supporting the assertion with additional evidence that, say, the NAO existed more or less as we know it know during the early Holocene NAO, when the presence of upstream ice sheets and different insolation and vegetation regimes were present, or by providing stronger caveats for extending the NAO discussion prior to the late Holocene. Perhaps using the term “NAO-like” instead of the NAO, while emphasizing that early-Holocene climate had distinctly different forcings and boundary conditions than the late Holocene, would be advisable.

Revised. This is an issue raised by all reviewers and its detailed comment can be found at the beginning of this document.

One other conclusion is pretty easily testable but doesn't seem to have been evaluated rigorously: that precipitable water is less during negative wNAOi states. The current study would have much stronger standing with readers if estimates of precipitable water (say from the NCEP database) and wNAOi were compared directly.

Revised. We have included NCEP/NCER reanalysis data of precipitable water. We evaluated the dependence of the precipitable water on the wNAOi over Europe and have added a new figure in the manuscript.

Specific Comments:

Line 122: change to “more strongly negative”.

Revised.

Please give the altitudes of the >350m non-alpine stations. If they don't differ much from the <350m stations (line 134), then why separate them out? Would it be better to include them with the <350 m stations because of similar response to the NAO?

Revised. The response of the non-Alpine stations is indeed similar to the NAO as for the continental stations: this is indicated by the sensitivity of the proxy to the NAO as well as by the calculated slopes. The slopes of the continental gradient in response to the NAO calculated for all continental stations (including the non-Alpine stations) are similar compared to one shown in the manuscript and are within the range of uncertainties.

Line 296: replace “confirmed” with “supported”. From a semantics point of view, observations can “confirm,” but models, being not real, can only support.

Revised.

Line 337: change exceptions to exception, or provide another example.

Revised.

Lines 411-417: the point about modern relationships maybe not being representative of past conditions is important and requires some more emphasis.

Revised (see revision for general comments).

Line 443: would read better as “situated in the Swiss Jura mountains approximately xxx km from the alpine divide. . .”

Revised.

Line 446: equilibrium typo; line 446 “net” not “nett”

Revised.

Line 457: There is no Figure 8 in the manuscript. Supplemental Figure? I would like to such a figure in the main text, as it is a crucial test of the current manuscript's hypothesis.

Revised.

Line 458-460: this is where I would suggest the assumption of stationarity of the wNAOi and d18Opw is not supported. Certainly not for the “entire Holocene”, but probably true for the past few millennia or so after ice sheets had decayed and land vegetation was established. One way around this problem is, for the pre-late Holocene, to refer to “NAO-like” behavior.

Revised. See also comment in the Section General Comments.

Actions to the comments of Reviewer 2

General Comments:

It is not clear to me that the ECHAM5-wiso simulations add anything substantial to the manuscript. They seem to me to interfere with the flow and the communication of the main points of the research. The research does not provide a thorough test of the model, or a comparison with other existing models (both are beyond the scope of the manuscript). Because the manuscript uses measured data (e.g., NAO data, GNIP data), there exists little uncertainty regarding the quality of the data or the synoptic conditions associated with the data. **So why include the model results, which are considerably less certain than the actual data, and in fact are sometimes inconsistent with the other results, and presumably wrong? I would suggest either i) remove all mentions to the model or ii) move the model and associated discussion to the supplemental material.** The strongest reason for keeping the model in is the independent analysis of the amount of precipitable water in the atmosphere - so maybe Figure S2 could be moved to the main text, with the 13 main GNIP sites marked on it, and the rest of the model discussion moved to the supplemental material. I think that a reference to the Supplemental Material would suffice as an explanation of how the new figure (the old S2) was constructed.

Revised. We have moved the ECHAM5-wiso evaluation to the Supplementary Material and it is discussed there in a new section and mention the results in the manuscript when it fits the discussion. We have additionally included NCEP/NCER reanalysis data of the amount of precipitable water to independently constrain the dependence of the amount of precipitable water on the wNAOi. The evaluations show that the dependence of the amount of precipitable water between the NCEP/NCER reanalysis data and the ECHAM5-wiso simulation are similar.

I wonder if a spatially stationary gradient can satisfactorily capture the complexity of the NAO. In particular, I suggest that the authors consider the ‘Augmented NAO Index’ of Wang et al., 2012, GRL within the context of their own work. The presumed stationarity of the pressure systems defining the NAO is an issue with most NAO studies, but because this study focuses on a spatial gradient perhaps it should focus more attention on this issue. Would the longitudinal gradient not change orientation through time as pressure centres shift location? Particularly in the Early Holocene, where more extensive sea ice would have relocated the pressure centers?

Revised. This is an issue raised by all reviewers and its detailed comment can be found at the beginning of this document.

How dependent are the gradients calculated in Figure 4 on Valentia and Wallingford? Would the gradients be uniformly flat if these two sites were omitted? Even if this is the case, it does not imply a problem with the conclusions, though I suggest the authors investigate whether or not the gradients are driven by these two sites.

Revised. We have included an additional figure in the supplementary material. The results emphasise that the maritime stations have only a minor influence on temperature gradients. The effect is stronger for the absolute values of the precipitation gradients but the NAO dependence is only slightly modified. Therefore, conclusions drawn from the original dataset are not hampered by the including the maritime stations.

Also, should these be referred to as ‘continental sites’ or as ‘maritime sites’ particularly since they do seem to behave very differently than the other sites?

Revised.

Specific Comments:

Line 23: ‘analyzed extensively’ instead of ‘with great effort’, which suggests the analyses themselves were difficult.

Revised.

Line 82: how was it determined that these stations had no Mediterranean influence?

We did not determine whether these stations have a physical influence of Mediterranean moisture, but labelled continental stations as non-Mediterranean influenced if they have a distance to the Mediterranean coastline >100km. We have included this information in the manuscript.

Line 447: ‘nett’

Revised.

Line 468: ‘Central Europe’ or ‘central Europe’ – be consistent

Revised.

Actions to the comments of Reviewer 3

General and Specific Comments:

Line 40 ff.: The second paragraph of the introduction is relatively long and mainly summarises what will be shown and discussed in the paper. This section could be shortened substantially to make the paper more concise.

Revised. We shortened it slightly.

Line 46: “...to better evaluate the NAO-dependence on isotope longitudinal gradients.” I thought the gradient depends on the state of the NAO and not the other way round? This should be clarified here and throughout the paper.

Revised.

Line 56: 37 stations have been analysed. 28 are GNIP stations, 6 are ANIP stations. What about the remaining 3?

Revised.

Line 126 ff.: “Comparison of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients derived from the ECHAM5-wiso with those from the station-based data show that slopes from the ECHAM5-wiso data reproduce the observed station-based slopes quite well (Figure 4).” I do not agree with this statement. All model data sets show a curve (i.e., the most negative slopes are shown for the 3rd and 4th NAO class) rather than a linear relationship with the NAO classes. It is, thus, misleading to state that the model data reproduce the station data “quite well”. **It would be good to see the fit statistics (slope, r^2 and p-value) not only for the station data, but for the model data as well (compare caption of Fig. 4).**

Revised. We have included the r^2 and p-values of the evaluation of the ECHAM5-wiso simulations in the figure caption of Figure S3.

Line 212 ff.: “Repeating the calculations using the vapour-ice phase change (snow) instead results in calculated differences that are still too small to explain the observed differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ between the western- and eastern-most stations (not shown).” Please provide a bit more information on this. I do not request a detailed discussion, but in the present form, the reader would not be able to do the calculations themselves if they wanted to.

We have revised this sentence by clarifying what parameter is used, stated its value that reader can repeat the calculation and have included the reference from which the values were used.

Line 217 ff.: “By contrast with the observational (GNIP) data discussed above, the ECHAM5-wiso simulated differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ can largely be accounted for by model air-temperature differences alone.” ... “Overall, however, these results supports the conclusion above that the winter air temperature effect on the longitudinal winter $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients is insufficient to explain the observed difference between the western and eastern GNIP stations.” This is not clear to me and even appears contradictory. If the effects observed in the model data can be explained by the model temperatures, this does not support the conclusion derived from the data. Please clarify.

In addition: “It is intriguing that the observed (GNIP datasets) and simulated (ECHAM5- wiso simulations) temperature slopes differ (Figure 6), while the slopes for longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients are apparently similar (Figure 4).” As far as I understood the text, the model temperatures in Eastern Europe are colder than those of the station data. If the $\delta^{18}\text{O}$ and δD values (i.e., the gradient and its dependence on the state of the NAO) are similar in the model and the data, but the temperatures are different, this either means that the dependence of model temperature on the NAO is too strong or that the sensitivity of the model $\delta^{18}\text{O}$ and δD values is too weak. In any case, this is an important difference, which makes it difficult to use the model data to interpret the station data. Based on the discussion following below, however, this statement is not necessary.

We have revised this section and deleted the sentence that is commented by the reviewer. The statement of this sentence is already given in the previous paragraph and it was confusing in the paragraph discussing the ECHAM5-wiso simulations.

Line 330 ff.: “The reason for the different strength of these two mechanisms (temperature gradient and precipitation history) on the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients for the observed (GNIP) and simulated (ECHAM5-wiso) datasets remains unclear, suggesting that the ECHAM5-wiso simulations warrant further investigation.” I am not a climate modeller, but how representative are the precipitation data of the model for the 13 stations (still a relatively low number) considered here. As far as I know, simulating (high-resolution) precipitation patterns is still difficult. **Thus, the model data (which represent climate variability in a larger grid cell) may be more representative for the dependence of the west-east gradient in precipitation on the NAO than the station data. In summary, it is not very surprising for me that the precipitation data of the stations do not show a dependence on the state of the NAO, but the model data do.**

This is an interesting point that the reviewer highlights here. However, it is beyond the scope of this study to evaluate the robustness of the precipitation here. This would require a rigorous comparison between different models and model types (GCM vs. regional models) and observational as well as reanalysis data.

Line 334 ff.: I would suggest to strongly shorten the first paragraph of section 4.2. It only summarises results from the analysis, which has partly already been presented in section 3.2. I would move all these results in section 3.2, and briefly summarise the findings here in one or two sentences.

Revised.

Line 401 ff.: I would remove the reference to tree rings here, which mainly record summer climate (and water isotopes).

Revised.

Line 426 ff.: “Note that our assumption about the temperature change represents a limiting case, because it implies that the annual air temperature, to which the cave air temperature is usually equilibrated, also decreases by the same value.” As the authors state themselves, this assumption is not reasonable. It may be possible to find a correlation between winter and annual temperature (in the station and the model data). Based on that, one could try to estimate the dependence of annual temperature on the state of winter NAO. However, temperature is very stable in most caves, and an inter-annual change of 1.3 °C, as assumed in the example for the station Stuttgart, is almost impossible and may only occur in a strongly ventilated cave. In such caves, however, other effects, such as precipitation of CaCO_3 under conditions of disequilibrium stable isotope fractionation or evaporation, will probably dominate the $\delta^{18}\text{O}$ values of the speleothem. The temperature effect may only be visible on a decadal or even longer time scale. **Thus, the reference to persistent changes in the NAO on centennial to millennial time scales should be given at the beginning of the paragraph.** However, I would rather suggest to remove the calculation because the caveats may not be present to many readers.

Revised. We clarified the calculations and state some more words on the caveats.

Section 5 in general: Since this section discusses the potential of speleothems for an NAO reconstruction based on speleothems, **I miss a critical discussion of other potential “problems” of speleothems for NAO reconstruction (smoothing of the signals in the aquifer, contributions of**

different seasons than winter, disequilibrium stable isotope fractionation, dating uncertainties, etc.). I know that the authors are aware of these problems, so they should not be omitted from the discussion here. Mischel et al. (2015) have modelled some of these processes in detail. Their study could be referenced in this context.

Revised. We have included a discussion on these caveats subsequently to the discussion of the NAO reconstruction. The study of Mischel et al. (2015) is cited right at the beginning of the discussion of the NAO reconstructions.

Finally, as the two other reviewers question the stationarity of the NAO and the gradient in the past and, in particular, its dependence on the location of the pressure systems during the Early Holocene, it may be interesting to read and discuss the recent paper by Wassenburg et al. (2016).

Revised. This is an issue raised by all reviewers and its detailed comment can be found at the beginning of this document.

McDermott, F., Atkinson, T. C., Fairchild, I. J., Baldini, L. M., and Matthey, D. P.: A first evaluation of the spatial gradients in $\delta^{18}\text{O}$ recorded by European Holocene speleothems, *Global and Planetary Change*, 79, 275-287, 2011.

Wang, Y. H., Magnusdottir, G., Stern, H., Tian, X., and Yu, Y.: Decadal variability of the NAO: introducing an augmented NAO index, *Geophysical Research Letters*, 39, 2012.

Wassenburg, J. A., Dietrich, S., Fietzke, J., Fohlmeister, J., Jochum, K. P., Scholz, D., Richter, D. K., Sabaoui, A., Spötl, C., and Lohmann, G.: Reorganization of the North Atlantic Oscillation during early Holocene deglaciation, *Nature Geoscience*, 2016.

North Atlantic Oscillation controls on oxygen and hydrogen isotope gradients in winter precipitation across Europe; implications for palaeoclimate studies

5 Michael Deininger¹, Martin Werner², Frank McDermott^{1,3}

¹UCD School of Earth Sciences, University College Dublin, Belfield, Dublin 4, Ireland

²Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bussestraße 24, 27570 Bremerhaven, Germany

³UCD Earth Institute, University College Dublin, Belfield, Dublin 4, Ireland

10 Correspondence to: Michael Deininger (michael.deininger@ucd.ie)

Abstract. Winter (October to March) precipitation $\delta^{18}\text{O}_p$ and δD_p values in central Europe correlate with the winter NAO index (wNAOi), but the causal mechanisms remain poorly understood. Here we analyse the relationships between precipitation-weighted $\delta^{18}\text{O}_p$ and δD_p datasets ($\delta^{18}\text{O}_{pw}$ and δD_{pw}) from European GNIP and ANIP stations and the wNAOi, with a focus on isotope gradients. We demonstrate that longitudinal $\delta^{18}\text{O}_{pw}$ and δD_{pw} gradients across Europe (continental effect) depend on the wNAOi state, with steeper gradients associated with more negative wNAOi states. Changing gradients reflect a combination of air temperature and variable amounts of precipitable water as a function of the wNAOi. The relationships between the wNAOi, $\delta^{18}\text{O}_{pw}$ and δD_{pw} can provide additional information from palaeoclimate archives such as European speleothems that primarily record winter $\delta^{18}\text{O}_{pw}$. Comparisons between present-day and past European longitudinal $\delta^{18}\text{O}$ gradients inferred from Holocene speleothems suggest that atmospheric pressure configurations akin to negative wNAO modes dominated the early Holocene, whereas patterns resembling positive wNAO modes were more common in the late Holocene, possibly caused by persistent shifts of the relative locations of the Azores High and the Icelandic Low.

1 Introduction

Stable oxygen and hydrogen isotopes ($\delta^{18}\text{O}$, δD ; relative to Vienna Standard Mean Ocean Water) in precipitation have been analyzed extensively since the 1950s (Dansgaard, 1954; Craig, 1961; Craig and Gordon, 1965) to decipher environmental processes that control their variations and to facilitate their use in palaeoclimatology (Dansgaard, 1964; Gat, 1996; Rozanski et al., 1992; Aggarwal et al., 2012). The Global Network of Isotopes in Precipitation (GNIP) program initiated in 1958 by the International Atomic Energy Agency (IAEA) and the World Meteorological Organization (WMO) has operated since 1961, and has made great progress towards these aims (e.g. Dansgaard, 1964; Rozanski et al., 1992). For the European sector of the GNIP program, 438 stations have operated, with the oldest records from the GNIP stations at Valentia Observatory in Ireland (since 1958), and at Hohe Warte (Vienna, Austria) since 1960. The processes that drive $\delta^{18}\text{O}$ values in precipitation ($\delta^{18}\text{O}_p$)

Michael 2.11.2016 12:12

Gelöscht: Winter (October to March) precipitation $\delta^{18}\text{O}_p$ and δD_p values in central Europe correlate with the winter NAO index (wNAOi), but the causal mechanisms remain poorly understood. Here we analyse the relationships between precipitation-weighted $\delta^{18}\text{O}_p$ and δD_p datasets ($\delta^{18}\text{O}_{pw}$ and δD_{pw}) from European GNIP and ANIP stations and the wNAOi, with a focus on isotope gradients. We demonstrate that longitudinal $\delta^{18}\text{O}_{pw}$ and δD_{pw} gradients across Europe (continental effect) depend on the wNAOi state, with steeper gradients associated with more negative wNAOi states. Changing gradients reflect a combination of air temperature and variable amounts of precipitable water as a function of the wNAOi. The relationships between the wNAOi, $\delta^{18}\text{O}_{pw}$ and δD_{pw} can provide additional information from palaeoclimate archives such as European speleothems that primarily record winter $\delta^{18}\text{O}_{pw}$. Comparisons between present-day and past European longitudinal $\delta^{18}\text{O}$ gradients inferred from Holocene speleothems suggest that negative wNAO modes dominated the early Holocene, but positive wNAO modes were more common in the late Holocene.

Michael 15.8.2016 13:29

Gelöscht: with great effort

North Atlantic Oscillation controls on oxygen and hydrogen isotope gradients in winter precipitation across Europe; implications for palaeoclimate studies

5 Michael Deininger¹, Martin Werner², Frank McDermott^{1,3}

¹UCD School of Earth Sciences, University College Dublin, Belfield, Dublin 4, Ireland

²Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bussestraße 24, 27570 Bremerhaven, Germany

³UCD Earth Institute, University College Dublin, Belfield, Dublin 4, Ireland

10 Correspondence to: Michael Deininger (michael.deininger@ucd.ie)

Abstract. Winter (October to March) precipitation $\delta^{18}\text{O}_p$ and δD_p values in central Europe correlate with the winter NAO index (wNAOi), but the causal mechanisms remain poorly understood. Here we analyse the relationships between precipitation-weighted $\delta^{18}\text{O}_p$ and δD_p datasets ($\delta^{18}\text{O}_{pw}$ and δD_{pw}) from European GNIP and ANIP stations and the wNAOi, with a focus on isotope gradients. We demonstrate that longitudinal $\delta^{18}\text{O}_{pw}$ and δD_{pw} gradients across Europe (continental effect) depend on the wNAOi state, with steeper gradients associated with more negative wNAOi states. Changing gradients reflect a combination of air temperature and variable amounts of precipitable water as a function of the wNAOi. The relationships between the wNAOi, $\delta^{18}\text{O}_{pw}$ and δD_{pw} can provide additional information from palaeoclimate archives such as European speleothems that primarily record winter $\delta^{18}\text{O}_{pw}$. Comparisons between present-day and past European longitudinal $\delta^{18}\text{O}$ gradients inferred from Holocene speleothems suggest that atmospheric pressure configurations akin to negative wNAO modes dominated the early Holocene, whereas patterns resembling positive wNAO modes were more common in the late Holocene, possibly caused by persistent shifts of the relative locations of the Azores High and the Icelandic Low.

1 Introduction

Stable oxygen and hydrogen isotopes ($\delta^{18}\text{O}$, δD ; relative to Vienna Standard Mean Ocean Water) in precipitation have been analyzed extensively since the 1950s (Dansgaard, 1954; Craig, 1961; Craig and Gordon, 1965) to decipher environmental processes that control their variations and to facilitate their use in palaeoclimatology (Dansgaard, 1964; Gat, 1996; Rozanski et al., 1992; Aggarwal et al., 2012). The Global Network of Isotopes in Precipitation (GNIP) program initiated in 1958 by the International Atomic Energy Agency (IAEA) and the World Meteorological Organization (WMO) has operated since 1961, and has made great progress towards these aims (e.g. Dansgaard, 1964; Rozanski et al., 1992). For the European sector of the GNIP program, 438 stations have operated, with the oldest records from the GNIP stations at Valentia Observatory in Ireland (since 1958), and at Hohe Warte (Vienna, Austria) since 1960. The processes that drive $\delta^{18}\text{O}$ values in precipitation ($\delta^{18}\text{O}_p$)

Michael 2.11.2016 12:12

Gelöscht: Winter (October to March) precipitation $\delta^{18}\text{O}_p$ and δD_p values in central Europe correlate with the winter NAO index (wNAOi), but the causal mechanisms remain poorly understood. Here we analyse the relationships between precipitation-weighted $\delta^{18}\text{O}_p$ and δD_p datasets ($\delta^{18}\text{O}_{pw}$ and δD_{pw}) from European GNIP and ANIP stations and the wNAOi, with a focus on isotope gradients. We demonstrate that longitudinal $\delta^{18}\text{O}_{pw}$ and δD_{pw} gradients across Europe (continental effect) depend on the wNAOi state, with steeper gradients associated with more negative wNAOi states. Changing gradients reflect a combination of air temperature and variable amounts of precipitable water as a function of the wNAOi. The relationships between the wNAOi, $\delta^{18}\text{O}_{pw}$ and δD_{pw} can provide additional information from palaeoclimate archives such as European speleothems that primarily record winter $\delta^{18}\text{O}_{pw}$. Comparisons between present-day and past European longitudinal $\delta^{18}\text{O}$ gradients inferred from Holocene speleothems suggest that negative wNAO modes dominated the early Holocene, but positive wNAO modes were more common in the late Holocene.

Michael 15.8.2016 13:29

Gelöscht: with great effort

at these stations have been discussed extensively in the literature, focusing mainly on air temperature, altitude and the continental effect (Rozanski et al., 1982; Stumpp et al., 2014; Schürch et al., 2003; Lykoudis and Argiriou, 2011; Fischer and Baldini, 2011). In recent years, several studies have investigated the relationship between the North Atlantic Oscillation (NAO) – the major atmospheric mode of European winter climate, (Hurrell et al., 2003; Hurrell and VanLoon, 1997; Hurrell, 1995) – and $\delta^{18}\text{O}_p$ at specific stations (Baldini et al., 2008; Langebroek et al., 2011; Field, 2010). More recently, the effect of the East Atlantic pattern, the second major mode of atmospheric pressure variability in the North Atlantic region (Barnston and Livezey, 1987) on the relationship between the winter NAO index (wNAOi) and the $\delta^{18}\text{O}_p$ in Europe has been examined (Comas-Bru et al., 2016). A complementary approach, developed here, is to document and seek to understand the longitudinal gradients in winter rainfall $\delta^{18}\text{O}_p$ across Europe, and to link these to different states of the NAO.

Here we present a detailed station-based evaluation of the effects of the wNAOi on precipitation-weighted $\delta^{18}\text{O}_p$ and δD_p values ($\delta^{18}\text{O}_{pw}$ and δD_{pw}) during the winter season for continental Europe and the Alps. We demonstrate that the well-documented longitudinal gradient in $\delta^{18}\text{O}_{pw}$ and δD_{pw} across the European continent (often referred to as the ‘continental effect’) depends on the state of the winter NAO index (wNAOi), through a combination of air temperature and atmospheric precipitation history effects. The downstream effects of this wNAOi-linked variability can also be detected in the $\delta^{18}\text{O}_{pw}$ and δD_{pw} values at stations north of the Alpine divide. We also make use of the $\delta^{18}\text{O}_{pw}$ and δD_{pw} outputs from an isotope-enabled General Circulation Model ECHAM5-wiso (Werner et al., 2011; Langebroek et al., 2011) to better evaluate the NAO-dependence of the longitudinal isotope gradients. This study improves our understanding of the significance of inferred past changes in $\delta^{18}\text{O}_{pw}$ or δD_{pw} gradients across Europe, for example during the course of the Holocene (e.g. McDermott et al., 2011) by linking modern $\delta^{18}\text{O}_{pw}$ and δD_{pw} gradients, respectively, to different states of the wNAO and/or changes in the European continental temperature gradient and atmospheric precipitation histories.

2 Data and Methods

2.1 Station-based data

Monthly winter (October to March) $\delta^{18}\text{O}_p$ and δD_p , temperature and precipitation data from 37 European stations were used to investigate the dependence of $\delta^{18}\text{O}_{pw}$ and δD_{pw} values on the wNAOi (Figure 1).

Thirty-one of these stations are part of the GNIP network and six are part of the Austrian Network of Isotopes in Precipitation (ANIP). The data from the ANIP stations originate from Humer (1995) and the ANIP homepage (<http://wisa.bmlfuw.gv.at/daten.html>). The GNIP stations were selected based on recent findings, showing that the unweighted winter (December to March) $\delta^{18}\text{O}_p$ and δD_p values at these stations are significantly correlated with the wNAOi (Baldini et al., 2008). The ANIP stations were selected for their long records starting in the early 1970s and their location, which ensures that there are sufficient $\delta^{18}\text{O}_p$ and δD_p measurements in the Alps for the purposes of this study. For our

Michael 15.8.2016 17:06

Gelösch: on

Michael 15.8.2016 17:06

Gelösch: longitudinal

Michael 28.9.2016 15:06

Gelösch: The model data evaluation shows that, first, the observed NAO-dependence of the $\delta^{18}\text{O}_{pw}$ and δD_{pw} and longitudinal gradients are reproduced by the ECHAM5-wiso outputs, and second that the variability of the ECHAM5-wiso longitudinal gradients are in the same range as the longitudinal gradients observed in the station data.

Michael 28.9.2016 15:06

Gelösch: or the last 35 ka (e.g. Rozanski, 1985),

Michael 26.10.2016 13:26

Gelösch: the precipitation-weighted winter $\delta^{18}\text{O}_p$ and δD_p values (

Michael 26.10.2016 13:26

Gelösch:)

Michael 15.8.2016 14:05

Gelösch: Twenty-eight

analysis, the wNAOi is the average winter value calculated from the monthly PC-based NAOi values from December to March (Hurrell, 1995) downloaded from the Climate Data Guide¹. For every station, the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values were calculated for each winter, using the monthly $\delta^{18}\text{O}_{\text{p}}$ and $\delta\text{D}_{\text{p}}$ values, weighted by the monthly precipitation amounts, for the period October to March (6 month) and December to February (3 month). The studied periods are chosen on the one hand because during December to February the NAO exerts its strongest influence on the European winter climate (e.g. Hurrell, 1995) and we expect to observe the strongest influence on $\delta^{18}\text{O}_{\text{p}}$ and $\delta\text{D}_{\text{p}}$ in this 3-month period and on the other hand because the main infiltration period for precipitation in central Europe is from October to March and is, therefore, the period of interest for palaeoclimate archives recording the infiltrated $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ in this 6-month interval (e.g. speleothems). For further analysis, the winter $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values (6 month and 3 month) of every station are grouped into six classes depending on the wNAOi. The wNAOi was sub-divided into six classes (bins): (i) $1.6 \leq \text{wNAOi}$; (ii) $0.8 \leq \text{wNAOi} < 1.6$; (iii) $0 \leq \text{wNAOi} < 0.8$; (iv) $-0.8 \leq \text{wNAOi} < 0$; (v) $-1.6 \leq \text{wNAOi} < -0.8$; (vi) $\text{wNAOi} < -1.6$, resulting in six compilations of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values for every station. For every station, the median of every compilation was calculated. An example is given in Figure 2, which shows how the data were processed, resulting in six median $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values for the Garmisch-Partenkirchen station (#24 in Figure 1).

For the analysis of the processed $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values, the stations were also subdivided into continental stations, which includes all stations with an altitude $\leq 350\text{m}$ and having no (or negligible) Mediterranean influence (distance to the Mediterranean coastline > 100 km) (13 stations); and stations with an altitude $> 350\text{m}$. The latter stations were divided into two groups: continental stations with an altitude higher than 350m (3 stations) and Alpine stations (17 stations). Furthermore, four Mediterranean GNIP stations south of the Alps were analysed to better validate the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values of the Alpine regions (Figure 1).

3 Results

3.1 Results of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients (continental stations)

To study the relationship between the wNAOi classes and the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients during the winter season, $\delta^{18}\text{O}_{\text{pw}}$, $\delta\text{D}_{\text{pw}}$ datasets of a total of 13 'continental' GNIP stations are analysed. (This includes the two maritime sites Valentia Observatory, Ireland, (station #1) and Wallingford, England, (station #2).) The most westerly station is at -10.25°E (Valentia Observatory; Ireland) and the easternmost station is at 19.85°E (Krakow, Poland) (Figure 1). These continental stations show a positive sensitivity of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ to the wNAOi (Figure 3), i.e., a positive linear relationship with the wNAOi for the investigated continental station datasets, resulting in more positive median $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values for higher

¹ The Climate Data Guide: Principal Component based North Atlantic Oscillation (NAO) Index: <https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-pc-based>.

Michael 26.10.2016 13:27
Gelösch: (Principal Component)
Michael 26.10.2016 13:28
Gelösch: precipitation-weighted $\delta^{18}\text{O}_{\text{p}}$ and $\delta\text{D}_{\text{p}}$ values (
Michael 26.10.2016 13:28
Gelösch:)

Michael 15.8.2016 13:39
Gelösch: C
Michael 27.9.2016 10:38
Gelösch: p
Michael 27.9.2016 10:38
Gelösch: p
Michael 26.10.2016 13:29
Gelösch: Thus, t

Michael 15.8.2016 13:34
Gelösch:

Michael 26.9.2016 16:20
Gelösch: 2.2 Isotope modelling analysis (ECHAM5-wiso) ... [1]

Michael 26.9.2016 16:22
Gelösch: ; ECHAM5-wiso

145 | wNAOi classes, as in Figure 2 for Garmisch-Partenkirchen. An exception to this observation is the GNIP stations Valentia
Observatory (Station 1) where the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values do not show a clear linear relationship compared to the other
continental stations for either the 3-month (Figure 3a and 3c) or 6-month (Figure 3b and 3d) winter periods. Similar
behaviour is seen in the data for the Wallingford GNIP station (Station 2) for the 6-month winter period (Figure 3b and 3d).
150 | This behaviour is likely to be caused by the proximity of these two maritime stations to the North Atlantic. It is notable, that
for most stations, the sensitivity of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ to the wNAOi is higher for the 3-month winter period than for the 6-
month winter period (Figure 3). The median sensitivity for $\delta^{18}\text{O}_{\text{pw}}$ is 0.87 (3 month) and 0.57 (6 month) ‰/wNAOi unit, and
for $\delta\text{D}_{\text{pw}}$ 6.87 (3 month) and 4.44 (6 month) ‰/wNAOi unit. This likely reflects the fact that the mode of the wNAOi
(December to March) exerts the strongest influence on European winter meteorology (e.g. Hurrell, 1995; Hurrell et al., 2003),
and so on $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ (see discussion in Section 4.1).

155 | To investigate the dependence of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients from these 13 continental stations on the
wNAOi classes, the gradient slopes were calculated for each of the six wNAOi classes. For all winter month periods the
slope is negative (towards the east) indicating that the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values become more negative towards the east.
Furthermore, the slopes of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients of the continental stations becomes steeper for winters
160 | with more negative wNAOi classes (Figure 4). This means that ^{18}O and ^2H are more efficiently removed from the
atmospheric moisture with distance from the western margin, resulting in more strongly negative $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values
along the longitudinal transect during more negative NAO winters compared with more positive NAO winters. Furthermore,
the slope is steeper for the 3-month (DJF) averages compared with that for the 6-winter month averages.

165 | Comparison of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients derived from ECHAM5-wiso simulations with those from the
station-based data show that slopes from the ECHAM5-wiso data reproduce the observed station-based slopes quite well
(Figure 4). Only the slopes determined for the most negative wNAOi class of the 6-month winter period level off from the
empirically determined slopes (Figure 4a and 4c). Overall, however, there is good agreement between the ECHAM5-wiso
170 | isotope gradients and those derived from the observational datasets (see supplementary material for a detailed discussion on
ECHAM5-wiso simulations).

3.2 Alpine stations and other stations with an altitude > 350m

175 | Data from European stations with an altitude greater than 350m are separated into two groups: Alpine stations and non-
Alpine stations. The non-Alpine stations include Wasserkuppe-Rhoen (#20, 921m), Hof-Hohensaas (#28, 567m) and
Regensburg (#29, 377m), all located in Germany (Figure 1). These show similar $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ sensitivity to the wNAOi
as the other continental stations (Figure 3). The Alpine stations include a total of 17 stations that are well distributed over the
Alps (Figure 1). These data allow an evaluation of precipitation $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values and patterns in the entire Alpine
region as a function of the wNAOi. The sensitivity analysis of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ of the Alpine stations reveals a somewhat

Michael 26.10.2016 13:41

Gelösch: are

Michael 26.10.2016 13:41

Gelösch: at the

Michael 26.10.2016 13:42

Gelösch: it is

Michael 15.8.2016 12:57

Gelösch: depleted

Michael 26.10.2016 13:44

Gelösch: the

Michael 27.9.2016 10:45

Gelösch: is very good

185 more complex relationship to the wNAOi. For the 3-month winter period, the $\delta^{18}\text{O}_{\text{pw}}$ data from the Thonon-Les-Bains (14),
Längenfeld (21), Obergurgl (23), Böckstein (30), St. Peter (31), Villacher Alpe (32), Graz University stations (33) have a
weak or absent relationship to the wNAOi, while the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ data from the other Alpine stations have a similar
sensitivity to the wNAOi as the continental stations (Figure 3a). The $\delta\text{D}_{\text{pw}}$ data show similar results as $\delta^{18}\text{O}_{\text{pw}}$, but there are
different results for some other stations. For Thonon-Les-Bains (14) the relationship between $\delta\text{D}_{\text{pw}}$ and the wNAOi is
stronger compared to $\delta^{18}\text{O}_{\text{pw}}$ whereas the relationship for Grimsel is weaker (18) (Figure 3c). For the 6-month winter period,
190 the $\delta^{18}\text{O}_{\text{pw}}$ sensitivity to the wNAOi for most of the Alpine stations is comparably strong to that of the continental stations.
Only the stations at Thonon-Les-Bains (14), Längenfeld (21), Böckstein (30), St. Peter (31), Villacher Alpe (32), Graz
University (33) show a weak relationship to the wNAOi (like for the 3-month winter period). The Obergurgl station (23) has
a stronger relationship to the wNAOi compared with the 3-month winter period, but has a smaller average $\delta^{18}\text{O}_{\text{pw}}$ sensitivity
compared with the other Alpine stations. The highest $\delta^{18}\text{O}_{\text{pw}}$ sensitivity can be found for Hohenpeißenberg (22) and
195 Regensburg (29) and is 0.77 ‰/wNAOi unit (Figure 3b). The $\delta\text{D}_{\text{pw}}$ sensitivity of the Alpine stations for the 6-month winter
period shows a similar relationship as for the 3-month winter period. For stations in Bern (15), Guttannen (17), Obergurgl
(23) and Villacher Alpe (32), however, the conclusion is different compared with the 3-month winter period. The
relationship of $\delta\text{D}_{\text{pw}}$ to the wNAOi is weaker for Bern (17) whereas it is stronger for the other stations. The highest
sensitivity is observed for the Thonon-Les-Bains (14) and Meiringen (16) stations and is 7.42 and 8.84 ‰/wNAOi unit,
200 respectively (Figure 3d).

4 Discussion

4.1 Longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients

To explain the change in the slopes of the longitudinal winter $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients (Figure 4), the measured
temperatures and the amount of precipitation at the GNIP stations are evaluated. The GNIP station based continental
205 temperature and precipitation gradients for the different wNAOi classes show that the slopes of the longitudinal winter
temperature gradients become steeper for lower wNAOi values and are always negative (i.e. average winter temperatures are
always lower in the east), but no equivalent relationship is observed for the slopes of the winter precipitation gradients
(Figure 5). (The following conclusions are not hampered if the two maritime sites Valentia Observatory and Wallingford,
station #1 and #2 (Figure 1), are omitted from the compilation of all 'continental stations' (Figure S1).) The temperature
210 relationships suggests that while the winter air temperature clearly becomes colder from west to east, a higher average air
temperature gradient between western and eastern Europe occurs in more negative wNAOi winters. Furthermore, the
intercept of the linear regression is progressively smaller for more negative wNAOi modes, suggesting general cooler
conditions in central Europe during negative wNAOi modes. This is consistent with the general relationship between the
wNAOi and winter air temperatures for central Europe (e.g. Hurrell, 1995; Comas-Bru and McDermott, 2013). Curiously, for
215 the most negative wNAOi class, the slope of the temperature gradient does not follow the general trend and has a lower

Michael 26.10.2016 13:57

Gelösch: -

Michael 26.10.2016 13:57

Gelösch: -

Michael 26.10.2016 13:58

Gelösch: the

Michael 26.10.2016 13:58

Gelösch: stations

Michael 26.9.2016 16:25

Gelösch: and from the relevant ECHAM5-wiso model grid cells were evaluated. Furthermore, the total precipitable water was analysed for the ECHAM5-wiso dataset. These variables were also grouped according to wNAOi class of their respective winters, and the median of temperature and precipitation were calculated for each wNAOi class (Figures 5 and 6).

Similar temperature-NAO relationships are observed for the ECHAM5-wiso simulations, with increasingly steeper model air temperature gradients and a smaller temperature intercept for lower wNAO indices (Figure 6a and 6b). However, in comparison with the slopes derived from the GNIP datasets, the slopes derived from the ECHAM5-wiso simulations suggest steeper temperature gradients (Figure 6a).

The intercept of the temperature regression derived from the ECHAM5-wiso simulations is similar to that from the observational datasets (Figure 6b). Therefore, the longitudinal temperature gradients of the ECHAM5-wiso simulations suggest comparable temperatures at 0°E (the temperature intercept), but much cooler temperatures in eastern Europe (a steeper temperature slope) compared with the observed air temperatures in the GNIP datasets.

245 value, comparable to more positive winter wNAOi modes, suggesting a smaller temperature difference between western and eastern Europe under these conditions. The reason for this change is unclear and reflects possibly a relationship of the cyclone variability to the NAO (Gulev et al., 2001), which may increase the frequency of incursions of cold easterly winds into western Europe during very negative wNAO modes.

250 The observed temperature slopes can be used to calculate the expected air temperature-driven difference in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ between the western- and eastern-most GNIP stations using theoretical temperature sensitivities for $\delta^{18}\text{O}_{\text{p}}$ and $\delta\text{D}_{\text{p}}$ (e.g. from Dansgaard, 1964). The theoretical changes in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ between the western- and eastern-most GNIP stations (the longitudinal difference between the Valentia (Observatory) and Krakow (Wola Justowska) is 30.1°) were calculated as follows. First, the temperature difference between these two stations was calculated from the temperature slope for the different wNAOi classes. For example, for the highest wNAOi class, the slope of the observed temperature gradient is -0.19 K/ $^\circ\text{E}$ and -0.26 K/ $^\circ\text{E}$ for the 6-month and 3-month winter period, resulting in a temperature difference of 5.86 K and 7.78 K, respectively. Hence, the average winter temperature at Valentia Observatory from October to March (December to February) is about 5.86 K (7.78 K) warmer compared to Krakow Wola Justowska for the highest wNAO class. The effect of this eastward temperature decrease can be converted into an expected $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ difference between the two stations. For this estimate we apply an approximate estimation of the sensitivity of $\delta^{18}\text{O}_{\text{p}}$ and $\delta\text{D}_{\text{p}}$ on temperature changes based on theoretically derived values by Dansgaard (1964), assuming a Rayleigh-type moist adiabatic condensation (vapour-liquid) process. For the sensitivity, the average value for a cooling from an initial temperature of 0°C to -20°C is used, which is 0.64 ‰/K for $\delta^{18}\text{O}_{\text{p}}$ and 5.6 ‰/K for $\delta\text{D}_{\text{p}}$ (Dansgaard, 1964). Hence, for the highest wNAO class, the observed temperature difference would cause a calculated difference of 3.75 ‰ (6-month) and 4.98 ‰ (3-month) for $\delta^{18}\text{O}_{\text{pw}}$. $\delta^{18}\text{O}_{\text{pw}}$ values are therefore expected to be 3.75 ‰ and 4.98 ‰ lighter at the easternmost station compared to the westernmost station for the 3 winter month and 6 winter month averages, respectively.

260 These theoretical calculated values are now compared with the observed differences derived from the slopes of the linear regression of the GNIP $\delta^{18}\text{O}_{\text{pw}}$ datasets (Figure 4). For the highest wNAO class, the slope is -0.16 ‰/ $^\circ\text{E}$ and 0.20 ‰/ $^\circ\text{E}$ for the 6-month and 3-month winter period. This results in an observed difference of 4.78 ‰ (6-month) and 6.02 ‰ (3-month). Importantly, the observed differences (longitudinal gradients) are much larger than those calculated using the air-temperature driven Dansgaard-type model described above. The results of these calculations for $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$, based on the observed temperature slopes, and for all wNAO classes are listed in Table 1.

275 The most important result of this simple exercise is that the observed differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ between the western- and eastern-most GNIP stations are larger than can be accounted for by a simple air temperature driven Rayleigh distillation model alone (Table 1). Repeating the calculations using the sensitivity of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ for the vapour-ice phase change (snow) (0.73 ‰/K for $\delta^{18}\text{O}_{\text{p}}$ and 6.2 ‰/K for $\delta\text{D}_{\text{p}}$, Dansgaard, 1964) instead results in calculated differences that are still too small to explain the observed differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ between the western- and eastern-most stations (not shown).

Michael 26.9.2016 16:27

Gelöscht: -

Michael 26.9.2016 16:27

Gelöscht: Similar temperature-NAO relationships for the ECHAM5-wiso simulations, with increasingly steeper model air temperature gradients and a smaller temperature intercept for lower wNAO indices (Figure 6a and 6b). However, in comparison with the slopes derived from the GNIP datasets, the slopes derived from the ECHAM5-wiso simulations suggest steeper temperature gradients (Figure 6a). The intercept of the temperature regression derived from the ECHAM5-wiso simulations is similar to that from the observational datasets (Figure 6b). Therefore, the longitudinal temperature gradients of the ECHAM5-wiso simulations suggest comparable temperatures at 0°E (the temperature intercept), but much cooler temperatures in eastern Europe (a steeper temperature slope) compared with the observed air temperatures in the GNIP datasets.

Michael 26.10.2016 14:05

Gelöscht: (GNIP datasets) and simulated (ECHAM5-wiso)

Michael 26.9.2016 16:27

Gelöscht: (By comparison, in the ECHAM5-wiso output, the temperature difference is somewhat greater at 7.73 K and 9.64 K for the highest wNAO class for the 6-month and 3-month winter periods, respectively.)

Michael 26.10.2016 14:07

Gelöscht: (

Michael 26.10.2016 14:07

Gelöscht:

Michael 26.10.2016 14:07

Gelöscht:)

Michael 26.10.2016 14:08

Gelöscht: -

Michael 26.10.2016 14:08

Gelöscht: -

Michael 15.8.2016 17:11

Gelöscht: n

Michael 26.10.2016 14:12

Gelöscht: (Dansgaard, 1964

Michael 27.9.2016 10:53

Formatiert: Tiefgestellt

Michael 27.9.2016 10:53

Formatiert: Tiefgestellt

This strongly indicates that the observed longitudinal variations in winter air temperatures alone simply are insufficient to account for the observed difference between $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ in western and eastern Europe, and that additional processes must be considered.

315 Rozanski et al. (1982) pointed out that the precipitation history of air masses is an important control on observed longitudinal δD_p gradients. Precipitation history can be expressed as a numerical value f (fraction of remaining moisture). Hence, f depends on the balance between the amount of precipitation (P) that has already occurred along the longitudinal gradient and the initial amount of precipitable water [in the atmosphere](#) (Q_0). However, only weak relationships are found between the precipitation gradients calculated from the GNIP station precipitation datasets and the wNAOi classes, suggesting that

320 rainfall gradients between western and eastern Europe are fairly constant, within the range of uncertainty (Figure 5c and 5d). Furthermore, the intercept of the linear regression of the precipitation data shows no dependence with the class of the wNAOi. This is consistent with the findings of Baldini et al. (2008) who showed that precipitation data from continental GNIP stations have no systematic correlation to the wNAOi (opposite to temperature). As discussed below, this points to differences in the initial amounts of precipitable water (Q_0) as a possible control on isotope gradient-wNAOi relationships

325 (Figure 4), over and above those attributable to air temperature gradients alone.

To better constrain the governing physical mechanisms of the precipitation history f and their first-order effects on $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$, a simple one-box Rayleigh-type model for the atmosphere was used (Dansgaard, 1964; Eriksson, 1965):

$$330 \quad {}^xR = {}^xR_0 \cdot f^{x\alpha-1} = {}^xR_0 \left(\frac{Q}{Q_0}\right)^{x\alpha-1} = {}^xR_0 \left(1 - \frac{P}{Q_0}\right)^{x\alpha-1}. \quad (1)$$

Equation (1) is the classic Rayleigh distillation model (Rayleigh, 1902; see also Mook, 2006 for detail) that describes the evolution of an isotope ratio R (the subscript x is a place holder for $x = 2$ (^2H) or 18 (^{18}O), i.e., for $^{18}\text{R} = ^{18}\text{O}/^{16}\text{O}$ and $^2\text{R} = ^2\text{H}/^1\text{H}$) as a function of the precipitation history f . This is a function of the amount of initial (Q_0) and remaining (Q)

335 precipitable water in the atmosphere; Q is therefore the amount of precipitable water after a specific amount of precipitation P ($Q_0 - Q = P$) has formed. R_0 describes the initial isotope ratio. α is the equilibrium liquid-water isotope fractionation factor that depends only on temperature; the subscript x denotes, as above, the related isotope system. Because the slope of the longitudinal precipitation gradient does not change systematically with the wNAOi class, P does not change with the class of the wNAOi, and has, therefore, a negligible effect on $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$. Hence, the slope of the longitudinal gradient of

340 $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ in central Europe is driven only by the temperature dependent isotope fractionation factor α , and by the initial amount of precipitable water Q_0 in the atmosphere. Although the classic Rayleigh-type model, adopted here is unable to fully capture all atmospheric processes (e.g. mixing of atmospheric moisture with different isotope signatures and/or origins), it is nonetheless a useful first approximation to explain the deviations between the calculated (temperature effect)

Michael 26.9.2016 16:28

Gelöscht: By contrast with the observational (GNIP) data discussed above, the ECHAM5-wiso simulated differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ can largely be accounted for by model air-temperature differences alone. Only for the most negative winter NAO class does the expected temperature-driven west to east change in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ exceed the ECHAM5-wiso simulated differences derived from slope of the lowest wNAOi class. If the slope of the ECHAM5-wiso longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients were driven by winter air temperature gradients alone, the average temperature sensitivity for the remaining five wNAOi classes for the 6-month winter period for $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ calculated from the ECHAM5-wiso simulations varies between 0.59 and 0.63 ‰/K for $\delta^{18}\text{O}_{\text{pw}}$ and 4.44 and 4.87 ‰/K for $\delta\text{D}_{\text{pw}}$. For the 3-month winter period, equivalent temperature sensitivities from the ECHAM5-wiso simulations range between 0.51 and 0.58 ‰/K for $\delta^{18}\text{O}_{\text{pw}}$ and 4.09 and 4.65 ‰/K for $\delta\text{D}_{\text{pw}}$ and are smaller than for the 6-month winter period. Hence, the temperature sensitivities for $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ derived from the ECHAM5-wiso are apparently somewhat lower compared to the theoretically estimated ones based on the approach of Dansgaard (1964). One explanation for the larger temperature sensitivities derived by the Dansgaard (1964) approach may be that the initial temperature at which the moisture condensation begins in the atmosphere is actually greater than 0°C as assumed in our theoretical calculations. Higher condensation temperatures are reasonable considering that ... [2]

Michael 26.9.2016 16:30

Gelöscht: By contrast, in the ECHAM5-wiso simulations, a clear relationship exists between the slope and the intercept of the longitudinal precipitation gradient (Figure 6c and 6d). The slopes, as well as the intercepts of the precipitation gradients are smaller for lower wNAOi classes, suggesting that the west-east difference of the amount of precipitation decreases (increases) for lower (higher) wNAOi classes in the model. Furthermore, the decreasing (increasing) intercept of the precipitation gradient for lower (higher) wNAOi classes suggests a lower (higher) amount of rainfall in the western grid cells. Analysis of the total amount of precipitation for all grid cells reveals lower values for the total amount of rainfall in the selected grid cells (i.e. the sum of precipitation from all grid cells) for lower wNAOi classes. For the 6-month winter period, for example, the median of the total precipitation decreases from 1091 mm for the highest wNAOi class to 852.5 mm for the lowest wNAOi class. In the case of the 3-month winter period, the median of the total precipitation decreases from 1270 mm to 800 mm from the highest to the lowest wNAOi class. For the lowest wNAOi class, the total amount of precipitation of all grid cells is about 78 % and 63 % of that from the highest wNAOi class for the 6-month and 3-month winter periods, respectively. Thus, the ECHAM5-wiso output is not consistent with the observc ... [3]

and observed differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ between the western- and eastern-most stations. Therefore, our observations of the dependence of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradient on the class of the wNAOi can be explained if additionally the amount of precipitable water over central Europe is lower for more negative wNAOi values. As demonstrated previously in a comparison of very positive (wNAOi>1) and negative (wNAOi<-1) wNAOi values, the maximum amount of precipitable water in the atmosphere is shifted southward for very negative wNAOi (Trigo et al., 2002). This shift in the amount of precipitable water is associated with changing air temperature patterns, and smaller amounts of precipitable water are associated with cooler continental air temperatures. Accordingly, our simple model shows that the atmosphere in central Europe contains less (more) atmospheric moisture during more negative (positive) wNAOi states, and this is independently supported by analysis of the amount of precipitable water in the atmosphere from a NCEP/NCER reanalysis dataset (see below). Hence, atmospheric moisture $\delta^{18}\text{O}$ and δD values are likely to be more sensitive to the rainout history during more negative wNAOi modes, because f must change at a higher rate if Q_0 is smaller and P is held constant (Eq. 1). This effect is also confirmed by a multi-box exercise that assumes a Rayleigh-type condensation process, mimicking the longitudinal gradients. This multi-box exercise shows that $\delta^{18}\text{O}_p$ becomes progressively depleted in dependence on Q_0 from west to east across Europe, with steeper gradients (higher slopes) for smaller Q_0 (Figure S2).

The wNAOi-precipitable water relationship described by Trigo et al. (2002) is also evident in the NCEP/NCER reanalysis dataset (and the ECHAM5-wiso simulations, see supplementary material) (Figure 6a). The NCEP/NCER reanalysis dataset covers the period from 1948-2016 and is analysed for the winter months December to March; it has a spatial resolution of $2.5^\circ \times 2.5^\circ$. Precipitable water shows a positive correlation with wNAOi values in central and northern Europe (where all continental stations are located), and a negative correlation over the Mediterranean (including Iberia, the Balkans and Turkey) (Figure 6). (Interestingly, the boundary between the area of positive and negative correlation lies within the Alpine region, though the coarse resolution does not allow any detailed conclusion for individual Alpine stations.) This indicates that during positive wNAO modes, the amount of precipitable water increases over central Europe and decreases during negative wNAO modes. This finding is also confirmed by the amount of precipitable water of the analysed NCEP/NCER reanalysis grid cells (Figure 6b and 6c). The results of the regression analysis of the longitudinal gradient of the precipitable water shows that the slope of the precipitable water along the longitudinal gradient is rather independent (within the range of uncertainties, although a slight trend is visible indicating a shallower slope for more negative NAOi classes) of the wNAOi class (Figure 6c), while the intercept clearly decreases for smaller wNAOi classes (Figure 6b). Compared with the amount of precipitable water for the highest wNAOi class, the atmosphere contains only about 88.4% for the lowest wNAOi class for the 6-month winter period for example. As a result, the precipitation history f becomes more sensitive to the rainout history along the longitudinal gradient for lower wNAOi classes.

To summarise, the dependence of the observed longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradient on the class of wNAOi in the winter season results from two processes: (i) the changing continental temperature gradients via the temperature dependent isotope

Michael 15.8.2016 13:13

Gelöscht: confirmed

Michael 26.9.2016 16:32

Gelöscht: the

Michael 26.9.2016 16:31

Gelöscht: ECHAM5-wiso

Michael 26.9.2016 13:32

Gelöscht: 1

Michael 26.9.2016 16:33

Gelöscht: S2

Michael 27.9.2016 11:08

Gelöscht: ECAHMS-wiso

Michael 27.9.2016 11:08

Gelöscht: 6e

Michael 27.9.2016 11:08

Gelöscht: 6f

Michael 27.9.2016 11:09

Gelöscht: modelled

Michael 27.9.2016 11:09

Gelöscht: 6e

Michael 27.9.2016 11:09

Gelöscht: f

Michael 27.9.2016 11:14

Gelöscht:

Michael 27.9.2016 11:19

Gelöscht: 5

Michael 27.9.2016 11:19

Gelöscht: 7

Michael 27.9.2016 11:20

Gelöscht: In the ECHAM5-wiso simulations, the total amount of precipitation (78 % for the 6-month winter period) decreases more along the longitudinal transect than does the amount of precipitable water.

Michael 27.9.2016 11:20

Gelöscht: less

Michael 27.9.2016 11:27

Gelöscht: The differences between the theoretically derived temperature sensitivity by Dansgaard (1964) and the temperature sensitivity derived by the ECHAM5-wiso simulations could thus be ascribed to the changing ratio of total precipitation and amount of precipitable water as seen in the ECHAM5-wiso simulations.

565 fraction during condensation, which exerts the strongest influence on the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradient and (ii) the dependence of the amount of precipitable water or in general on the precipitation history, over central Europe on the wNAOi mode, which becomes important for more negative wNAOi classes. The latter mechanism is in agreement with recent findings of Aggarwal et al. (2012) who showed that, generally, more negative $\delta^{18}\text{O}_{\text{p}}$ values are associated with lower moisture residence times, where the moisture residence time is defined as the ratio between the amount of precipitable water and the precipitation (Aggarwal et al., 2012; Trenberth, 1998) and conclusions of (Rozanski et al., 1982) analysing summer and winter European $\delta\text{D}_{\text{p}}$ longitudinal gradients.

570 4.2 Alpine stations

By comparison with the low-altitude stations, the Alpine stations reveal more complex wNAOi - $\delta^{18}\text{O}_{\text{pw}}$, $\delta\text{D}_{\text{pw}}$ patterns (Figure 3). North of the Alpine divide, all stations show similar $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ - wNAOi class relationships as at the Garmisch-Partenkirchen GNIP station (Figure 2). The only exception to this relationship are the $\delta^{18}\text{O}_{\text{pw}}$ datasets for Thonon-Les-Bains, whose $\delta^{18}\text{O}_{\text{pw}}$ datasets have only a weak relationship to the wNAOi. The relationship between the $\delta^{18}\text{O}_{\text{pw}}$ - $\delta\text{D}_{\text{pw}}$ values and wNAOi from stations at and south of the Alpine divide, including Grimsel (#18) (western Alps); Längenfeld (#21), Obergurgl (#23), Patscherkofel (#27) (all central Alps); Böckstein (#30), St. Peter (#31), Villacher Alpe (#32) and Graz (#33) (eastern Alps) is more complex compared to the stations north of the Alpine divide (see section 3.2 for detail).

580 The $\delta^{18}\text{O}_{\text{pw}}$, $\delta\text{D}_{\text{pw}}$, temperature and precipitation datasets were also grouped into six wNAOi classes according to respective wNAOi as previously for the non-Alpine stations (Section 4.1). A detailed analysis of the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values of all Alpine stations shows that the median $\delta^{18}\text{O}_{\text{pw}}$ values become more negative for higher altitudes, irrespective of the wNAOi class. This observation is well known as the “altitude effect” (Dansgaard, 1954; Schürch et al., 2003). However, there is no obvious relationship ($p > 0.1$ for all datasets) between the class of the wNAOi and the altitude effect for the Alpine stations (Fig. 7a and 7b). On average, the altitude effect is -0.32 ‰/100m (6-month average) and -0.30 ‰/100m (3-month average) for $\delta^{18}\text{O}_{\text{pw}}$ and -2.55 ‰/100m (6-months average) and -2.28 ‰/100m (3-months average) for $\delta\text{D}_{\text{pw}}$ (Fig. 7a and 7b). Furthermore, the recorded air temperature at the Alpine stations change on average by about -0.58 and -0.56 K/100m for the 6-month and 3-month average, respectively, independent of the wNAOi class ($p > 0.1$ for all datasets) (Fig. 7c). The mean values of the temperature-altitude relationship correspond approximately to the moist adiabatic lapse rate. Hence, a strong relationship between $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values and air temperature is observed in the Alpine stations (e.g. Schürch et al., 2003). No relationship between rainfall amount and $\delta^{18}\text{O}_{\text{pw}}$ was observed (Fig. 7d).

595 Because there is no relationship between the lapse rate and and/or precipitation amount with the wNAOi class, we conclude that the observed relationships between $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ and the class of the wNAOi for our selection of Alpine stations are (i) caused by different air mass origins linked to wNAOi states and (ii) downstream effects of the varying central European continental effect that causes more negative $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values for more negative wNAOi classes. The weak or absent

Michael 27.9.2016 11:22

Gelösch: For the ECHAM5-wiso, air temperature gradients are clearly the most important factor that controls the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients. In ECHAM5-wiso, the precipitation history seems to be of relatively minor importance, because changes in the amount of precipitable water are mediated by changes in the simulated amount of precipitation. The reason for the different strength of these two mechanisms (temperature gradient and precipitation history) on the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients for the observed (GNIP) and simulated (ECHAM5-wiso) datasets remains unclear, suggesting that the ECHAM5-wiso simulations warrant further investigation.

Michael 28.9.2016 15:08

Gelösch: Precipitation at these stations is depleted in ^{18}O and ^2H for more negative wNAOi winters.

Michael 15.8.2016 13:14

Gelösch: s

Michael 28.9.2016 15:10

Gelösch: For the 3-month $\delta\text{D}_{\text{pw}}$ datasets only, the isotope data from Patscherkofel show a strong relationship to the wNAOi, while all other stations have weak or no relationships to the wNAOi (Figure 3c). In the 6-month dataset, only the southern-most stations (Längenfeld and St. Peter) show a weak relationship to the wNAOi. All other stations have a strong relationship to the wNAOi (Figure 3d). The $\delta\text{D}_{\text{pw}}$ dataset of the southern- and eastern-most Alpine stations, Villacher Alp (32) and Graz (33), respectively, exhibit no relationships with the wNAOi for the 3-month datasets; for 6-month winter period only the $\delta\text{D}_{\text{pw}}$ dataset of Villacher Alpe has a relationship to the wNAOi. For the $\delta^{18}\text{O}_{\text{pw}}$ datasets the behaviour of the relationships to the wNAOi is similar to the $\delta\text{D}_{\text{pw}}$ datasets. The 3-month $\delta^{18}\text{O}_{\text{pw}}$ datasets from Grimsel (#18) and Patscherkofel (#27) show a relationship to the wNAOi but all other stations exhibit no or only weak relationships to the wNAOi (Figure 3a). The 6-month datasets from Längenfeld (21) and St. Peter (31) have a weak relationship to the wNAOi (Figure 3b). The 6-month $\delta^{18}\text{O}_{\text{pw}}$ datasets from Villacher Alpe and Graz also show weak relationships to the wNAOi.

Michael 28.9.2016 15:10

Gelösch: -

Michael 26.10.2016 14:29

Gelösch: class

Michael 28.9.2016 15:10

Gelösch: -

... [4]

relationships for stations at the Alpine divide can be caused by air masses of different origins (e.g. variable influences of the Mediterranean sourced moisture) (Kaiser et al., 2001). However, detailed back trajectories of rainfall events for the entire Alpine region would be required to further evaluate this explanation. Our analysis indicates that the Alpine divide exerts an important influence on the winter hydrological cycle in the region, with precipitation north of the Alps sourced by atmospheric moisture originating from central Europe. In winter, this residual atmospheric moisture is already depleted in ^{18}O and ^2H when it reaches the northern part of the Alps, reflecting the ambient winter mode of the wNAOI, thereby determining the degree of the depletion in ^{18}O and ^2H in north Alpine winter precipitation.

To complete the above conclusion on the mixing of atmospheric moisture for stations at the Alpine divide, the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ of circum-Mediterranean stations were also analysed for their dependence on the wNAOI (a discussion of the NAO-relationships between $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ and temperature and precipitation can be found in the supplementary information). These stations are Avignon (34) (southwest of the Alps), Locarno (35) (south of Grimsel), Genoa (Setri) (south of the Alps) (36) and Zagreb (37) (southeast of the Alps) (Figure 1). For the 3-month $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ data only those from Avignon and Zagreb shows a strong relationship to the wNAOI (about 1 ‰/wNAOI unit for $\delta^{18}\text{O}_{\text{pw}}$) (Figure 3a and 3c). For the 6-month $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ data, only the $\delta^{18}\text{O}_{\text{pw}}$ dataset from Locarno and the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ datasets from Zagreb show a relationship to wNAOI. The NAO-relationships of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ for the 3-month winter period from the Mediterranean stations Locarno and Genoa (Setri) show that the NAO-fingerprint, which is observed for Alpine stations north of the Alpine divide, is not transferred to these two stations. The situation might change for $\delta^{18}\text{O}_{\text{pw}}$ from Locarno for the 6-month winter period where a relationship to the wNAOI is observed. The stronger relationship for this winter period could be caused by an increase of precipitation that results from air masses from central Europe. For Zagreb, it is difficult to explain the observed relationships between $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$, because the closest Alpine stations show no relationship to the wNAOI. To further investigate the mechanism that control the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ datasets in Avignon and Zagreb, the origin of the air masses in dependence on the wNAOI needs to be investigated further using isotope enabled regional climate models to better constrain the effect of local temperature and precipitation on $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$. In summary, the variable wNAO-relationship of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ datasets from the investigated Mediterranean stations support the observation of that the Alpine divide represents an important boundary region of the oxygen and hydrogen isotope system of Alpine precipitation.

5 Implications for palaeoclimate reconstructions from speleothems

5.1 Reconstructions from single speleothem-based carbonate $\delta^{18}\text{O}$

The results have important implications for palaeoclimate archives that record the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values of winter precipitation (October to March, the main period of infiltration in central Europe due to low evapo-transpiration rates). Potentially, such archives include speleothems and ground water. The following discussion focuses on speleothem carbonate $\delta^{18}\text{O}$ records, but is also applicable to speleothem fluid inclusion $\delta^{18}\text{O}$ and δD records, and is relevant for other palaeoclimate archives.

675 For a paleoclimate reconstruction that is based on a single speleothem from a cave site from central Europe or north
of the Alpine divide, $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values are typically lower for more negative wNAOi values (Figure 3). The median
sensitivity of $\delta^{18}\text{O}_{\text{pw}}$ from all continental stations is 0.57 ‰ per wNAOi unit. Hence, a persistent change of the average
wNAOi from e.g. +1 to -1 (i.e. -2 wNAOi units) would result in a reduction of the average $\delta^{18}\text{O}_{\text{pw}}$ by 1.14 ‰. Furthermore,
680 monthly average air temperatures are generally in phase with the wNAOi changes, resulting in a positive linear relationship
between $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ and air temperatures. Hence, air temperatures tend to be lower depending on the cave location, if
the average wNAOi is smaller. This temperature-relationship was used recently to reconstruct historic $\delta^{18}\text{O}_{\text{pw}}$ values
(Mischel et al., 2015) and its use has been suggested for wNAOi reconstructions (Casado et al., 2013). Crucially, however,
the $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ variability is controlled not only by changes in air temperature, but also by changes in the air mass
precipitation history (Section 4). Hence, the use of speleothem $\delta^{18}\text{O}$ values (or $\delta^{18}\text{O}$ and δD values from fluid inclusions or
ground water) to directly reconstruct the past variability of the wNAOi or winter temperatures should be undertaken
685 cautiously, because past changes of the hydrological cycle could result in a relationship between $\delta^{18}\text{O}_{\text{pw}}$ and temperature that
differs from the present day. This is particularly important for long-term reconstructions of atmospheric circulation. As for
the case of individual speleothem $\delta^{18}\text{O}$ records, caution should be exercised when interpreting changes in $\delta^{18}\text{O}$ gradients
inferred from multi-speleothem regression analysis because the continental precipitation $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients are
controlled by both the air temperature gradient and the air-mass precipitation history (Section 4).

690 Notwithstanding these caveats, the results from the sensitivity analysis of $\delta^{18}\text{O}_{\text{pw}}$ and air temperature theoretically
allow estimates of the possible influences of changes in $\delta^{18}\text{O}_{\text{pw}}$ and temperature for wNAOi changes on speleothem $\delta^{18}\text{O}$
records. For this exercise we use the station data of Stuttgart (Canstatt) (#8), located in southern Germany with a $\delta^{18}\text{O}_{\text{pw}}$ -
wNAOi sensitivity similar to the median of all continental stations. For the period from October to March, the NAO-
temperature relationship for this station suggests a sensitivity of 0.63 K/wNAOi unit, and an intercept of 4.38 °C ($r=0.57$); no
695 NAO-relationship is observed for precipitation amount. In the example above, a decrease in the NAO index from +1 to -1
would cause a change of -1.20 ‰ in $\delta^{18}\text{O}_{\text{pw}}$ in Stuttgart precipitation, with a concomitant decrease in air temperature of
approximately 1.26 K. Assuming that these changes in $\delta^{18}\text{O}_{\text{pw}}$ and air temperature on the surface are transmitted into a cave,
we can estimate both effects on speleothem $\delta^{18}\text{O}$ values. However, this is only reasonable for persistent changes of the NAO
on decadal to millennial time scales when there is sufficient time for thermal equilibration of the cave air temperature to
700 changes of the surface air temperature; and where any annual variability of the wNAO (i.e. 'noise') is overwhelmed by the
persistent change of the wNAO ('signal').

Applying the temperature sensitivity of the equilibrium oxygen-isotope fractionation factor of Kim and O'Neil
(1997) of about -0.22 ‰/K, the temperature change of -1.26 K would result in an increase in speleothem $\delta^{18}\text{O}$ by about
+0.28 ‰, but the simultaneous decrease by 1.20 ‰ in drip water $\delta^{18}\text{O}$ would dominate, resulting in an overall decrease in the
705 speleothem $\delta^{18}\text{O}$ by about 0.92 ‰. Therefore, the changes of precipitation $\delta^{18}\text{O}_{\text{pw}}$ would dominate the speleothem $\delta^{18}\text{O}$
record. We stress that these idealised conditions are rarely met in natural cave systems where other processes typically
influence the speleothem proxy values (e.g. isotope disequilibrium effects Deininger et al., 2012). Nonetheless, this exercise

710 emphasises that the sensitivity of $\delta^{18}\text{O}_{\text{pw}}$ to wNAOi is large and that persistent changes (i.e. centennial and millennial changes) in the mean state of the wNAO are likely to produce detectable and coherent changes in speleothem $\delta^{18}\text{O}$ time series in central Europe.

715 It is notable that for speleothems deposited in caves close to the Alpine divide where there is little or no relationship between $\delta^{18}\text{O}_{\text{pw}}$ and the wNAOi, speleothem $\delta^{18}\text{O}$ values are likely to be dominated by the temperature-dependent isotope fractionation factor during speleothem growth rather than by changes in $\delta^{18}\text{O}_{\text{pw}}$. Hence, if modern/historic temperature calibrations of speleothem $\delta^{18}\text{O}$ values are conducted, a relatively straightforward, but site-specific negative correlation between speleothem $\delta^{18}\text{O}$ and temperature is to be expected for sites close to the Alpine divide. This inference is supported by the two currently available studies: one at Spannagel Cave, Austria, which is located close to the Alpine divide in the central Alps where the temperature calibration is -0.44‰/K (Mangini et al., 2005) and one from Milandre Cave situated in the Swiss Jura Mountains (western Alps) approximately 150 km north of the Alpine divide, where the temperature calibration is 0.70‰/K (Fleitmann, 2016 personal communication). Taking the temperature sensitivity of $\delta^{18}\text{O}_{\text{pw}}$ of about 0.95‰/K for the GNIP station Konstanz, located at the similar latitude as Milandre Cave, and the temperature sensitivity of the equilibrium isotope fractionation between water and calcite of about -0.22‰/K , yields a net value of 0.73‰/K . This value is broadly consistent with the temperature calibration (0.70‰/K) found for Milandre Cave by Fleitmann.

725 The above discussion focused only on changes in $\delta^{18}\text{O}_{\text{pw}}$ and temperature that are transmitted into the cave. However, in natural cave systems, other environmental parameters linked to persistent changes of the wNAO could also influence speleothem $\delta^{18}\text{O}$. These include parameters such as cave air CO_2 concentrations (pCO_2), the Ca^{2+} concentration of drip waters or drip intervals. Modelling (e.g. Deininger et al., 2012; Mühlinghaus et al., 2009) and experimental work (e.g. Polag et al., 2010; Wiedner et al., 2008; Day and Henderson, 2011) reveal that these parameters can influence speleothem (calcite) $\delta^{18}\text{O}$ via isotope disequilibrium effects. These latter effects can in principle be as large as the NAO-related changes calculated above (see Deininger et al., 2012 for a thorough discussion on the sensitivity of all these parameters). Therefore, such reconstructions should be always undertaken cautiously, because they can include large uncertainties. Furthermore, $\delta^{18}\text{O}_{\text{pw}}$ variability can be smoothed during water percolation through the karst system, resulting in rather constant drip water $\delta^{18}\text{O}$ values which may further influence speleothem $\delta^{18}\text{O}$ (Mischel et al., 2015). The effects of these complications can be minimised through the use of well-monitored cave systems where the 'modern' (meaning over the duration of the monitoring period) cave system and the response of the diverse parameters monitored to natural changes of the surface conditions are well constrained, combined with appropriate forward models (e.g. ISOLUTION, Deininger et al., 2012) that allow estimates of the magnitude of these effect on speleothem $\delta^{18}\text{O}$. Further, the use of multiple coeval speleothem records from different cave systems can be regarded as a potential (new) approach to synthesise the common signal of all of these records (e.g. $\delta^{18}\text{O}$) using for example Principal Component Analysis (Deininger et al., 2016) or other statistical techniques (Rehfeld et al., 2013; Fischer, 2016).

5.2 Multiple speleothem-based reconstructions of Holocene longitudinal speleothem $\delta^{18}\text{O}$ gradients and implication on the evolution of the North Atlantic Oscillation

745 Although the aforementioned processes can modify speleothem $\delta^{18}\text{O}$ values, their effects might be negligible or at least less
important when the palaeoclimate signals of multiple coeval speleothem $\delta^{18}\text{O}$ records are used to reconstruct past climate
dynamics. As an example, McDermott et al. (2011) used $\delta^{18}\text{O}$ data for several coeval speleothems to show that longitudinal
speleothem $\delta^{18}\text{O}$ gradients across Europe changed systematically during the Holocene. To compare the observed slopes and
their temporal evolution during the Holocene and to evaluate a possible NAO-like mechanism the 6-month (October-March)
longitudinal $\delta^{18}\text{O}_{\text{pw}}$ gradients need to be converted into expected speleothem (calcite) $\delta^{18}\text{O}$ gradients. Such estimates require
750 that both the longitudinally variable winter temperature gradients (Figure 5a), as well as the $\delta^{18}\text{O}_{\text{pw}}$ slopes (Figure 4a) are
taken into account. The effects of varying temperature gradients are considered via the temperature-dependent oxygen
isotope fractionation between water and calcite. These estimates show that the slope of the longitudinal speleothem $\delta^{18}\text{O}$
gradient is expected to be greatest for the highest wNAOi class, decreasing for lower wNAOi classes. The calculated slopes
for speleothem calcite based on the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ slopes and air-temperature gradients of this study show remarkable
755 agreement with a recent reconstruction of speleothem based $\delta^{18}\text{O}$ gradients throughout the Holocene (McDermott et al.,
2011). (Figure 8).

Assuming that the present-day mechanisms that determine the observed relationships between the wNAOi and
temperature, precipitation and $\delta^{18}\text{O}_{\text{pw}}$ are relevant for the boreal winter of the entire Holocene, the evolution of the
reconstructed Holocene speleothem $\delta^{18}\text{O}$ gradients can be interpreted as reflecting changes in atmospheric pressure patterns
760 akin to predominantly negative wNAOi-like modes in the early Holocene until about c. 8 ka, and mainly positive wNAOi-
like modes during mid and late Holocene winters (this assumption includes caveats that are discussed further below). For
this assumption (NAO type forcing), the changes of the speleothem $\delta^{18}\text{O}$ gradients are caused by changing temperature
gradients (steeper west to east European temperature gradients in the early Holocene) and a changing precipitation history
(increased precipitation and/or reduced amount of moisture in the atmosphere) linked to a NAO-like forcing. However, this
765 interpretation requires that the NAO was operating akin to modern observations. In other words: it hypothesises that the
Azores High and the Icelandic Low were broadly stationary in space and time. A recent study has demonstrated recent
(1871-2008 AD) non-stationarity of the Azores High (AH) and the Icelandic Low (IL), (expressed as the NAO angle index,
which is positive when the IL is east of the AH and vice versa), but the latter is correlated with the wNAOi: a positive angle
index tends to coincide with a positive wNAOi (Wang et al., 2012). For the Holocene, stationarity is hypothesised for the last
770 8 ka by Wassenburg et al. (2016) who proposed that the NAO reorganised between 9 ka and 8 ka ago in response to the
melting Laurentide ice sheet and meltwater fluxes into the North Atlantic (Wassenburg et al., 2016). This study further
suggests that the Icelandic Low was shifted southwest of its modern position during the early Holocene. The timing of this
NAO reorganisation would be consistent with the finding of this study that the NAO switched from predominantly negative
to positive wNAOi-like at about 8 ka. However, in contrast to the findings of Wassenburg et al., (2016), our study based on

775 the results of McDermott et al. (2011) would suggest a gradual shift of the IL and/or AH towards the northeast or southwest, respectively, during the course of the Holocene until stable boundary conditions are reached 5 to 4 ka ago.

780 The different inferences between this study (and that of McDermott et al., 2011) and Wassenburg et al. (2016) may be that the observations of Wassenburg et al. (2016) are based on two stalagmite proxy records only, located in North Germany and North Morocco whose proxy records are sensitive for the local amount of precipitation. The modern precipitation amount of the two regions is driven mainly by the NAO, but with opposite correlations, positive in North Germany and negative in North Morocco (cf. Figure 1b in Wassenburg et al., 2016). Hence, if these two speleothem precipitation-proxy records were compared from these two regions we would expect that, if the modern precipitation regimes had not changed, these two proxy records would be anti-correlated. If, however, the modern precipitation regimes changed (e.g. they shifted in response to changes of the atmospheric circulation) and if the local precipitation amount at the two
785 locations change in the same direction, the two proxy records should be correlated. This scenario is exactly observed by Wassenburg et al. (2016) who show that the correlation between these two speleothem proxy records from North Germany and North Morocco changed from positive to negative between 9 ka and 8 ka (see Wassenburg et al., 2016 for details). However, the correlation of these two records is only a local perspective of the history on the amount of precipitation at these two individual locations and cannot resolve gradual changes of the precipitation pattern, but can only indicate whether the locations lie within the same hydrological regime or not. By contrast, the results of McDermott et al. (2011) that are further
790 evaluated by this study are based on multiple speleothem $\delta^{18}\text{O}$ records that are distributed throughout Europe and are potentially more reliable to detect gradual changes. However, we emphasise that the results of our study and those of Wassenburg et al. (2016) do not contradict, but rather complement each other, and give two different views on the same phenomena, a gradual change of the 'NAO' during the Holocene that is likely caused by a relative shift of the IL and AH, mimicking NAO-like conditions in the speleothem $\delta^{18}\text{O}$ gradients. The synthesis of the aforementioned studies and the results presented here is that the relative location of the IL and AH changed gradually during the Holocene: in the early Holocene the IL and/or the AH were located southwest and northeast, respectively, of their modern positions, eventually
795 mimicking negative NAO-like conditions and subsequently shifted northeastwards and southwestwards respectively during the Holocene, mimicking positive and modern NAO-like conditions, consistent with an independent study from the Iberian Peninsula (Walczak et al., 2015). This gradual change was potentially forced by the retreat of the Laurentide ice sheet during the early Holocene (Wassenburg et al., 2016) and possibly also by the changing Holocene wintertime insolation (McDermott et al., 2011;Walczak et al., 2015).

800 The approach applied by this study uses modern relationships between longitudinal precipitation $\delta^{18}\text{O}_{\text{pw}}$ gradients and the wNAOi and argues that the observed changes are caused i) by changing temperature gradients between west and east Europe and ii) a modified precipitation history f that accounts for changes in the amount of precipitable water in the atmosphere and in principle changing precipitation amounts. Though the latter (changes in the precipitation amount) was not evident in the analysed station datasets, there is no reason to exclude it for the Holocene interpretation of changing longitudinal speleothem
805

810 $\delta^{18}\text{O}$ gradients. This approach includes several caveats that need to be discussed in more detail. First, modern relationships
are derived for yearly winter values (6 month and 3 month average values) that are derived mainly post 1961 AD (when the
GNIP network begun). (The ECHAM5-wiso simulations used here also cover only the post 1960 AD period.) However in
815 during this period, the NAO angle index was rather stable and the relative location of the AH and IL did not vary much, compared
to the period before (Wang et al., 2012). Hence, we cannot rule out any effect of the relative location of the AH and IL on
the modern relationships between longitudinal $\delta^{18}\text{O}_{\text{pw}}$ gradients and the NAO and the conversion of the speleothem $\delta^{18}\text{O}$
820 gradients into wNAOi should be interpreted cautiously. Furthermore, recent studies have also highlighted that other
atmospheric pressure systems (like the East Atlantic pattern) also exert an influence on European $\delta^{18}\text{O}_p$ -NAO relationships
(Comas-Bru et al., 2016), but did not study its relationship to the angle index of the NAO. This would be, however,
interesting, because there might be relationships between the strength and location of the East Atlantic pattern and the angle
index of the NAO (i.e. to the relative location between the AH and the IL). Second, a requirement for the interpretation of
825 the changing longitudinal speleothem $\delta^{18}\text{O}$ gradients as variations of the NAO requires a time-stationary of the NAO back to
the early Holocene. However, this requirement can be questioned based on the recent findings of Walczak et al. (2015) and
Wassenburg et al. (2016) who demonstrated Holocene shifts of the AH and IL. Their conclusion is further supported by this
study. Hence, the gradual shift of the speleothem $\delta^{18}\text{O}$ gradients during the Holocene should be interpreted as relative shifts
of the AH and IL that are akin to the modes of the modern NAO (i.e. NAO-like) but likely have very different mechanism
(ice sheets, insolation) forcing mechanisms on these longer timescales. Further research is necessary to illuminate these
caveats extending on the one hand palaeoclimate investigations to generate new palaeoclimate proxy records and on the
other hand conducting climate model simulations to study in detail the relationships between the dominating pressure
systems and palaeoclimate proxy records as well as their underlying mechanism.

6 Conclusions and Summary

830 This study investigated the relationships between central European and Alpine $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values and the North
Atlantic Oscillation index for the boreal winter (wNAOi). $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ data for 37 meteorological stations distributed
over central Europe and the Alps were analysed. This study demonstrates that the European continental isotope effect
depends on the wNAOi, with steeper $\delta^{18}\text{O}_{\text{pw}}$ gradients associated with more negative wNAOi values. Hence, precipitation
835 $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values across central Europe are more negative for lower wNAOi winters. We argue that the strength of this
change is caused by (i) a steeper west to east continental air temperature gradient and (ii) a decrease in the precipitable water
content of the atmosphere during more negative wNAOi conditions. (An evaluation of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$
gradients from an isotope-enabled general circulation model, ECHAM5-wiso, shows that the simulated slopes are well
reproduced compared with those reconstructed from observational data).

840 The emerging picture for Alpine stations is that for stations north of the alpine divide, $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values show
a similar relationship to the wNAOi, as continental stations north of the Alps. However, in contrast with the processes that

Michael 2.11.2016 12:19

Gelöscht: 5 Implications for speleothem-based and other paleoclimate reconstructions and application to a reconstruction of Holocene longitudinal speleothem $\delta^{18}\text{O}$ gradients [5]

determine $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values at continental stations, $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ characteristics at Alpine stations are mainly a downstream residual effect of the changing continental effect. Furthermore, $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values of stations close to the Alpine divide in the central and eastern Alps exhibit a weak or absent relationship to the wNAOI. The results of this study have important implications for palaeoclimate reconstructions that are based on winter $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values, such as speleothem $\delta^{18}\text{O}$ records.

A simplistic interpretation of recent findings on the temporal evolution of longitudinal speleothem $\delta^{18}\text{O}$ gradients, assuming that modern NAO-related mechanisms are relevant for the Holocene, would be that the early Holocene was characterised by more negative NAO-like modes, while the mid and late Holocene would be characterised by more positive NAO-like modes. However, consistent with other independent studies (Wassenburg et al., 2016; Walczak et al., 2015) and an augmented NAO index that accounts for the relative position of the Azores High (AH) and Icelandic Low (IL) (Wang et al., 2012), we interpret the observed gradual Holocene change in longitudinal speleothem $\delta^{18}\text{O}$ gradients as a relative shift of the angular position of the AH and IL. These positions eventually evolve to situations akin to modern positive and negative NAO modes, but the millennial forcing mechanism during the course of the Holocene are likely to be different from the present day inter-annual to decadal variability. Further observational and modelling studies are necessary to better constrain the complex relationships between the atmospheric circulation and palaeoclimate proxy records.

Data availability

The processed data is available through supplementary material. Unprocessed data can be accessed through http://www-naweb.iaea.org/napc/ih/IHS_resources_gnip.html and <https://wasser.umweltbundesamt.at/h2odb/>.

Acknowledgement

We thank all operating organizations of the Global Network of Isotopes in Precipitation (GNIP) and the Austrian Network of Isotopes in Precipitation (ANIP) and all contributing researchers for their efforts to run and maintain the stations, which made this study possible. [We are grateful to three anonymous reviewers for their constructive comments that help to improve the manuscript and to the associated editor Dominik Fleitmann for handling our manuscript.](#) MD is funded by the Irish Research Council (IRC) by a Government of Ireland Postdoctoral Fellowship (GOIPD/2015/789). FMcD acknowledges support from Science Foundation Ireland through its Research Frontiers Program (RFP) Grants 07/RFP/GEOF265 and 08/FRP/GEO1184.

Michael 2.11.2016 12:19

Gelöscht: This study investigated the relationships between central European and Alpine $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values and the North Atlantic Oscillation index for the boreal winter (wNAOI). $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ data for 37 meteorological stations distributed over central Europe and the Alps were analysed. This study demonstrates that the European continental isotope effect depends on the wNAOI, with steeper gradients associated with more negative wNAOI values. Hence, precipitation $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values across central Europe are more negative for lower wNAOI winters. We argue that the strength of this change is caused by (i) a steeper west to east continental air temperature gradient and (ii) a decrease in the precipitable water content of the atmosphere during more negative wNAOI conditions. An evaluation of the longitudinal $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ gradients from an isotope-enabled general circulation model, ECHAM5-wiso, shows that the simulated slopes are well reproduced compared with those reconstructed from observational data. The mechanisms that cause the variability of the simulated slopes in the ECHAM5-wiso output are steeper compared with the observational datasets and suggest that the ECHAM5-wiso simulated $\delta^{18}\text{O}$ and δD values are largely controlled by the air temperature gradient, but this requires further investigation.

... [6]

References

- Aggarwal, P. K., Alduchov, O. A., Froehlich, K. O., Araguas-Araguas, L. J., Sturchio, N. C., and Kurita, N.: Stable isotopes in global precipitation: A unified interpretation based on atmospheric moisture residence time, *Geophysical Research Letters*, 39, 2012.
- 905 Baldini, L. M., McDermott, F., Foley, A. M., and Baldini, J. U. L.: Spatial variability in the European winter precipitation $\delta^{18}\text{O}$ -NAO relationship: Implications for reconstructing NAO-mode climate variability in the Holocene, *Geophysical Research Letters*, 35, L04709-L04709, 2008.
- Barnston, A. G., and Livezey, R. E.: Classification, seasonality and persistence of low-frequency atmospheric circulation patterns, *Monthly weather review*, 115, 1083-1126, 1987.
- 910 Casado, M., Ortega, P., Masson-Delmotte, V., Risi, C., Swingedouw, D., Daux, V., Genty, D., Maignan, F., Solomina, O., and Vinther, B.: Impact of precipitation intermittency on NAO-temperature signals in proxy records, *Climate of the Past*, 9, 871-886, 2013.
- Comas-Bru, L., and McDermott, F.: Impacts of the EA and SCA patterns on the European twentieth century NAO-winter climate relationship, *Quarterly Journal of the Royal Meteorological Society*, 2013.
- 915 Comas-Bru, L., McDermott, F., and Werner, M.: The effect of the East Atlantic pattern on the precipitation $\delta^{18}\text{O}$ -NAO relationship in Europe, *Climate Dynamics*, 47, 2059-2069, 2016.
- Craig, H.: Isotopic variations in meteoric waters, *Science*, 133, 1702-1703, 1961.
- Craig, H., and Gordon, L. I.: Deuterium and oxygen 18 variations in the ocean and the marine atmosphere, *Consiglio nazionale delle ricerche, Laboratorio de geologia nucleare*, 1965.
- 920 Dansgaard, W.: The O 18-abundance in fresh water, *Geochimica et Cosmochimica Acta*, 6, 241-260, 1954.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436-468, 1964.
- Day, C. C., and Henderson, G. M.: Oxygen isotopes in calcite grown under cave-analogue conditions, *Geochimica et Cosmochimica Acta*, 75, 3956-3972, 2011.
- Deininger, M., Fohlmeister, J., Scholz, D., and Mangini, A.: Isotope disequilibrium effects: The influence of evaporation and ventilation effects on the carbon and oxygen isotope composition of speleothems-A model approach, *Geochimica et Cosmochimica Acta*, 2012.
- 925 Deininger, M., McDermott, F., Mudelsee, M., Werner, M., Frank, N., and Mangini, A.: Coherency of late Holocene European speleothem $\delta^{18}\text{O}$ records linked to North Atlantic Ocean circulation, *Climate Dynamics*, 2016.
- Eriksson, E.: Deuterium and oxygen-18 in precipitation and other natural waters Some theoretical considerations, *Tellus A*, 17, 1965.
- 930 Field, R. D.: Observed and modeled controls on precipitation $\delta^{18}\text{O}$ over Europe: From local temperature to the Northern Annular Mode, *Journal of Geophysical Research: Atmospheres* (1984-2012), 115, 2010.
- Fischer, M. J., and Baldini, L. M.: A climate-isotope regression model with seasonally-varying and time-integrated relationships, *Climate dynamics*, 37, 2235-2251, 2011.
- 935 Fischer, M. J.: Predictable components in global speleothem $\delta^{18}\text{O}$, *Quaternary Science Reviews*, 131, 380-392, 2016.
- Gat, J. R.: Oxygen and hydrogen isotopes in the hydrologic cycle, *Annual Review of Earth and Planetary Sciences*, 24, 225-262, 1996.
- Gulev, S. K., Zolina, O., and Grigoriev, S.: Extratropical cyclone variability in the Northern Hemisphere winter from the NCEP/NCAR reanalysis data, *Climate Dynamics*, 17, 795-809, 2001.
- 940 Humer, G.: NIEDERSCHLAGSISOTOPENNETZ ÖSTERREICH. Daten, 1995.
- Hurrell, J. W.: Decadal Trends in the North-Atlantic Oscillation - Regional Temperatures and Precipitation, *Science*, 269, 676-679, 1995.
- Hurrell, J. W., and VanLoon, H.: Decadal variations in climate associated with the north Atlantic oscillation, *Climatic Change*, 36, 301-326, 1997.
- 945 Hurrell, J. W., Kushnir, Y., Ottensen, G., and Visbeck, M.: The North Atlantic Oscillation: climatic significance and environmental impact, *American Geophysical Union*, 2003.
- Kaiser, A., Scheifinger, H., Kralik, M., Papesch, W., Rank, D., and Stichler, W.: Links between meteorological conditions and spatial/temporal variations in long-term isotope records from the Austrian precipitation network, *C&SPaperSeries*, 13, 67-77, 2001.

- 950 Kim, S.-T., and O'Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochimica et Cosmochimica Acta*, 61, 3461-3475, 1997.
- Langebroek, P. M., Werner, M., and Lohmann, G.: Climate information imprinted in oxygen-isotopic composition of precipitation in Europe, *Earth and Planetary Science Letters*, 2011.
- 955 Lykoudis, S. P., and Argiriou, A. A.: Temporal trends in the stable isotope composition of precipitation: a comparison between the eastern Mediterranean and central Europe, *Theoretical and applied climatology*, 105, 199-207, 2011.
- Mangini, A., Spötl, C., and Verdes, P.: Reconstruction of temperature in the Central Alps during the past 2000 yr from a $\delta^{18}\text{O}$ stalagmite record, *Earth and Planetary Science Letters*, 235, 741-751, 2005.
- McDermott, F., Atkinson, T. C., Fairchild, I. J., Baldini, L. M., and Matthey, D. P.: A first evaluation of the spatial gradients in $\delta^{18}\text{O}$ recorded by European Holocene speleothems, *Global and Planetary Change*, 79, 275-287, 2011.
- 960 Mischel, S. A., Scholz, D., and Spötl, C.: $\delta^{18}\text{O}$ values of cave drip water: a promising proxy for the reconstruction of the North Atlantic Oscillation?, *Climate Dynamics*, 1-16, 2015.
- Mook, W. G.: *Introduction to isotope hydrology*, Taylor & Francis, 8, 2006.
- Mühlinghaus, C., Scholz, D., and Mangini, A.: Modelling fractionation of stable isotopes in stalagmites, *Geochimica et Cosmochimica Acta*, 73, 7275-7289, 2009.
- 965 Polag, D., Scholz, D., Mühlinghaus, C., Spötl, C., Schröder-Ritzrau, A., Segl, M., and Mangini, A.: Stable isotope fractionation in speleothems: Laboratory experiments, *Chemical Geology*, 279, 31-39, 2010.
- Rayleigh, L.: LIX. On the distillation of binary mixtures, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 4, 521-537, 1902.
- Rehfeld, K., Marwan, N., Breitenbach, S. F. M., and Kurths, J.: Late Holocene Asian summer monsoon dynamics from small but complex networks of paleoclimate data, *Climate dynamics*, 41, 3-19, 2013.
- 970 Rozanski, K., Sonntag, C., and Münnich, K. O.: Factors controlling stable isotope composition of European precipitation, *Tellus*, 34, 142-150, 1982.
- Rozanski, K., Araguás-Araguás, L., and Gonfiantini, R.: Relation between long-term trends of oxygen-18 isotope composition of precipitation and climate, *Science*, 258, 981-985, 1992.
- 975 Schürch, M., Kozel, R., Schotterer, U., and Tripet, J.-P.: Observation of isotopes in the water cycle—the Swiss National Network (NISOT), *Environmental Geology*, 45, 1-11, 2003.
- Stumpp, C., Klaus, J., and Stichler, W.: Analysis of long-term stable isotopic composition in German precipitation, *Journal of Hydrology*, 517, 351-361, 2014.
- Trenberth, K. E.: Atmospheric moisture residence times and cycling: Implications for rainfall rates and climate change, *Climatic change*, 39, 667-694, 1998.
- 980 Trigo, R. M., Osborn, T. J., and Corte-Real, J. M.: The North Atlantic Oscillation influence on Europe: climate impacts and associated physical mechanisms, *Climate Research*, 20, 9-17, 2002.
- Walczak, I. W., Baldini, J. U. L., Baldini, L. M., McDermott, F., Marsden, S., Standish, C. D., Richards, D. A., Andreo, B., and Slater, J.: Reconstructing high-resolution climate using CT scanning of unsectioned stalagmites: A case study identifying the mid-Holocene onset of the Mediterranean climate in southern Iberia, *Quaternary Science Reviews*, 127, 117-128, 2015.
- 985 Wang, Y. H., Magnusdottir, G., Stern, H., Tian, X., and Yu, Y.: Decadal variability of the NAO: introducing an augmented NAO index, *Geophysical Research Letters*, 39, 2012.
- Wassenburg, J. A., Dietrich, S., Fietzke, J., Fohlmeister, J., Jochum, K. P., Scholz, D., Richter, D. K., Sabaoui, A., Spötl, C., Lohmann, G., Andreae, M. O., and Immenhauser, A.: Reorganization of the North Atlantic Oscillation during early Holocene deglaciation, *Nature Geoscience*, 2016.
- 990 Werner, M., Langebroek, P. M., Carlsen, T., Herold, M., and Lohmann, G.: Stable water isotopes in the ECHAM5 general circulation model: Toward high-resolution isotope modeling on a global scale, *Journal of Geophysical Research*, 116, D15109-D15109, 2011.
- 995 Wiedner, E., Scholz, D., Mangini, A., Polag, D., Mühlinghaus, C., and Segl, M.: Investigation of the stable isotope fractionation in speleothems with laboratory experiments, *Quaternary International*, 187, 15-24, 2008.

1000


1005 **Tables**

wNAOi class	W-E difference for $\delta^{18}\text{O}_{\text{pw}}$									W-E difference for $\delta\text{D}_{\text{pw}}$							
	Observed		Calculated		Deviation			Observed		Calculated		Deviation					
	6m	3m	6m	3m	6m	3m	%	6m	3m	6m	3m	6m	3m	6m	3m		
	‰	‰	‰	‰	‰	%	‰	%	‰	‰	‰	‰	‰	%	‰	%	
I: 1.77	4.78	6.02	3.75	4.98	1.03	22.5	1.04	17.3	42.62	52.10	32.80	43.56	9.82	23.0	8.54	16.4	
II: 1.18	4.94	5.14	3.28	3.98	1.66	33.6	1.16	22.6	44.06	46.19	28.72	34.84	15.34	34.8	11.35	25.8	
III: 0.40	5.73	6.32	4.00	5.54	1.73	30.2	0.79	12.5	49.37	51.68	35.01	48.44	14.36	29.1	3.24	6.3	
IV: -0.46	6.66	7.89	4.18	5.58	2.48	37.2	2.31	29.3	56.93	65.10	36.58	48.81	20.35	35.7	16.29	25.0	
V: -1.17	6.22	7.38	4.44	6.55	1.79	28.8	0.83	11.2	53.44	61.11	38.82	57.31	14.62	27.4	3.80	6.2	
VI: -2.23	6.54	7.56	3.37	5.05	3.17	48.5	2.51	33.2	54.61	64.56	29.46	44.21	25.14	46.0	20.35	31.5	

Table 1. Differences in $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ (6 month (6m) and 3 month (3m) winter periods) between the western- (Valentia, Ireland) and eastern-most station (Krakow, Poland) for all wNAOi classes based on continental GNIP station datasets. The median wNAOi for classes I to VI respectively are: 1.77, 1.18, 0.40, -0.46, -1.17 and -2.23. The empirical estimates for the observed W-E difference of $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ presented here are based on linear regressions of the observed trends (Figure 4). The *calculated* $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values are based on the temperature difference between Valentia (Ireland) and Krakow (Poland) estimated from the linear regression of the continental GNIP station temperature datasets and temperature sensitivities for $\delta^{18}\text{O}_{\text{p}}$ and $\delta\text{D}_{\text{p}}$ from Dansgaard (1964). Deviation (bold numbers) means the difference between observed and calculated $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values. The italic numbers states the deviation relative to the observed difference given in %.

1010

1020



1025


1030 **Figures**

Figure 1. Map showing the location of all investigated stations: closed grey circles show the location of continental stations (altitude $\leq 350\text{m}$ with no Mediterranean influence); open triangles indicate alpine and high altitude ($>350\text{m}$) stations; closed squares show the location of ‘Mediterranean influenced’ stations. The grey line **in the box** indicates the Alpine Divide.

1035 Station codes for continental stations (from west to east): 1) Valentia (Observatory); 2) Wallingford; 3) Groningen; 4) Koblenz; 5) Karlsruhe; 6) Cuxhaven; 7) Bad Salzuflen; 8) Stuttgart (Cannstatt); 9) Würzburg; 10) Braunschweig; 11) Berlin; 12) Vienna (Hohe Warte); 13) Krakow (Wola Justowska). For high altitude stations: 14) Thonon-Les-Bains; 15) Bern; 16) Meiringen; 17) Guttannen; 18) Grimsel; 19) Konstanz; 20) Wasserkuppe-Rhoen; 21) Längenfeld; 22) Hohenpeisenberg; 23) Obergurgl; 24) Garmisch-Partenkirchen; 25) Scharnitz; 26) Neuherberg; 27) Patscherkofel; 28) Hof-Hohensaas; 29)

1040 Regensburg; 30) Bockstein; 31) St. Peter; 32) Villacher Alpe; 33) Graz Universität. For Mediterranean influenced stations:
34) Avignon; 35) Locarno; 36) Genoa (Sestri); 37) Zagreb.

1045






Figure 2. The upper part of the figure depicts how the precipitation weighted $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values are classified here into six classes depending on the wNAOI. Median values are calculated for every class, and this is used for further analysis. Panel a) to d) show the processed results for the six classes for one exemplar GNIP station (Garmisch-Partenkirchen, Germany), for which $\delta^{18}\text{O}_{\text{pw}}$ and $\delta\text{D}_{\text{pw}}$ values have a typical sensitivity to wNAOI (see Figure 3). The left panels show the box plots for the 6 months period (October to March) and the right panels for the 3 months period (December to February). The upper panels illustrate the $\delta^{18}\text{O}_{\text{pw}}$ values, the lower panels the $\delta\text{D}_{\text{pw}}$ values. Every box plot illustrates the statistical variables (median, min, max, 25 and 75% quantile) for every wNAOI class from lowest to highest (left to right). For the individual wNAOI classes, the red line illustrates the median of the data compilation; the edges of the blue rectangles mark the 25% and 75% quantile; the black bars illustrates the minimum and maximum values and the red cross denotes 'outliers'.

Michael 26.10.2016 15:05

Gelöscht: winter NAO index

1060






Figure 3. The panels illustrate the slopes of the linear regressions between yearly $\delta^{18}\text{O}_{pw}$ (a and b) and δD_{pw} (c and d) and the wNAOI for each individual station for the 3-month winter period (December-February) (left panels) and the 6-month winter period (October-March) (right panels). If the slope is illustrated in black the linear correlation coefficient is greater than 0.3; otherwise it is shown in orange. The blue lines indicate the median values of all continental stations; the upper and lower red lines highlight the 1-sigma standard deviation around the mean value. The grey dashed lines indicate a slope of 0 (i.e. no sensitivity). Station Garmisch-Partenkirchen (24) whose sensitivity on the wNAOI classes is shown in detail in Figure 2 is typical of the investigated stations. The cyan coloured arrows indicate stations closest to the Alpine Divide. Numbers indicate station codes as in Figure 1.



1075

Figure 4: Illustration of the slopes of the $\delta^{18}O_{pw}$ (a and b) and δD_{pw} (c and d) longitudinal gradients across Europe (filled circles) – and their respective standard errors – calculated from 13 continental GNIP stations for the 6-month (October-March) (a and c) and 3-month (December to February) (b and d) winter period. More negative wNAOi classes result in a steeper isotope gradient across Europe in winter and, therefore, more strongly depleted $\delta^{18}O_{pw}$ and δD_{pw} values with increasing distance (towards the east) from the European western margin. The coefficient for the linear regression between the observed slopes and the class of the wNAOi is 0.016 ± 0.004 ($r^2 = 0.79$; $p < 0.05$) for Fig. 1a, 0.019 ± 0.006 ($r^2 = 0.65$; $p = 0.0502$) for Fig. 1b, 0.111 ± 0.033 ($r^2 = 0.74$, $p < 0.05$) for Fig. 1c and 0.146 ± 0.048 ($r^2 = 0.70$, $p < 0.05$) for Fig. 1d (units are ‰ E⁻¹/wNAOi). These equations state the results from the linear regression where Y_{pw} is the slope of the $\delta^{18}O_{pw}$ or δD_{pw} gradient and X is the wNAOi. The filled grey squares show the median slopes of the ECHAM5-wiso simulations; the grey envelope indicates the 25% and 75% quantiles of the ECHAM5-wiso slopes.

1085



1090 **Figure 5.** Based on observational data only, these four panels illustrate the slope of the continental gradient for (a)
 1095 temperature and (c) precipitation as a function of the class of wNAOi that are calculated from GNIP station datasets. Panel
 b) and d) shows the intercepts of the linear regression for the continental temperature and precipitation gradients versus the
 wNAOi, respectively. Black symbols indicate the results for the 6-month winter period (October-March); grey symbols
 denote results for the 3-month (December-February) winter period. The slopes for temperature and precipitation show no
 relationship to the wNAOi if all six classes are analysed ($p > 0.1$). However, omitting the most negative wNAOi class yields a
 significant linear correlation of 0.71 and 0.67 ($p < 0.01$) for the 6 and 3 month averages, respectively. While there is no
 significant relationship between the intercept of the precipitation gradients with the wNAOi ($p > 0.1$), the intercept of the air
 temperature gradients shows a clear trend, with lower temperatures associated with lower wNAOi values.

1100




Figure 6: a) Correlation map between the wNAOI and the amount of precipitable water (PW) for the month December to March based on NCEP/NCER reanalysis data for the period 1948-2016 and the results of the longitudinal gradient (b) intercept; c) slope) for grid cells where continental stations are located (closed circles). Open triangles show the location of Alpine stations; open squares indicate the position of Mediterranean influenced stations.

1105

Michael 2.11.2016 12:20

Gelösch: Figure 6:

Michael 27.9.2016 11:31

Gelösch: ECHAM5-wiso outputs: The six panels illustrate the median slope of the gradient for a) temperature (T), c) precipitation (P) and e) the total precipitable water (TPW) depending on the class of the wNAOI from the analysed ECHAM5-wiso grid cells. Panel b), d) and f) shows the median intercept of the linear regression for the temperature, precipitation and total precipitable water gradient versus the class of wNAOI. The envelopes (straight line) indicate the 25% and 75% quantile. Back colours indicate the results for the 6-month winter period and grey colours the results from 3-month winter period. The dashed lines in panel a) to d) indicate the observed variability of these parameters derived from the observational datasets from the GNIP stations as illustrated in Figure 5.





Figure 7: The four panels show the rate of change of (a) $\delta^{18}\text{O}_{pw}$, (b) δD_{pw} , (c) temperature (T) and (d) precipitation (P) as a function of altitude for different wNAOi classes. (The lowest wNAOi class was not analysed because data of only four stations (out of 17) is available.) All Alpine stations are included. None of the correlations are statistically significant ($p > 0.1$). Black symbols indicate the results of the 6-month winter period; grey symbols denote the 3-month winter period.

1130



1135 **Figure 8:** The blue triangles illustrate the forward-modelled longitudinal slopes for speleothem records calculated for each wNAOi class from their respective 6-months (October-March) longitudinal $\delta^{18}\text{O}_{pw}$ gradients (closed circles) and the
 1140 observed temperature gradient converted into a speleothem $\delta^{18}\text{O}$ gradient using a sensitivity of 0.225 ‰/K (Kim and O'Neil, 1997). The blue and black line indicates the regression line from the longitudinal speleothem $\delta^{18}\text{O}$ and $\delta^{18}\text{O}_{pw}$ gradient, respectively. The blue bar highlights the range of reconstructed longitudinal speleothem $\delta^{18}\text{O}$ slopes during the Holocene (McDermott et al., 2011). The temporal evolution of the speleothem $\delta^{18}\text{O}$ slopes (McDermott et al. (2011) vary between -0.2208 ‰/°E and -0.1336 ‰/°E from 12 ka to 7 ka, and vary and between -0.1266 ‰/°E and -0.1046 ‰/°E from 6 ka to the present (shaded area). Comparison of the speleothem slopes with those calculated in this study suggests predominantly wNAO- like-modes in the early Holocene until about c. 8 ka and wNAO+ like-modes in the mid and late Holocene.

Michael 27.9.2016 16:18
 Gelöscht: E
 Michael 27.9.2016 16:18
 Gelöscht: Late