

## ***Interactive comment on “***

# **TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica” by D. Buiron et al.**

### **Anonymous Referee #1**

Received and published: 1 October 2010

The paper presents an age model for the Talos Dome ice core based on inverse modelling along the lines of Lemieux-Dudon et al., 2009. Roughly speaking, the inverse model seeks to adjust the flow, accumulation and  $\Delta$ age scenarios derived from simpler a priori model results so that the resulting gas and ice age scales agree optimally well with a set of chronological marker points that can represent ties between several cores being age-modelled at the same time (depth-depth ties) or independently dated horizons that are applied to the cores in question (age ties).

In my opinion, there are three main potentials for added value by the inverse modelling method: 1. The model allows parallel and consistent dating of several cores (if the tie

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



points are correct). 2. Both gas and ice tie points can be used, also in the case where the tie points are not fully internally consistent due to e.g. uncertainties. 3. On top of the age model itself, the model produces a consistent set of  $\Delta$ age, accumulation and thinning functions.

The application of the inverse model in this work is a simple version of the Lemieux-Dudon work, because only one core is being modelled (i.e. there are only age ties and no depth-depth ties), and because only gas-age ties are used for the young part and only ice age ties age used for the old part of the record. The advantage of using the model in this case is therefore almost entirely reduced to point (3) above.

I know this will sound provocative, but I would like to see a calculation / graph of how much the inverse model differs from a simple interpolation between the tie points (either linear interpolation depth vs. depth or linear interpolation of annual layer thicknesses), i.e. inverse model age minus interpolated age vs. depth. I have the feeling that the whole thing relies almost entirely on the tie points, and that one would get almost the same age model without applying the inverse model. If the differences (also in between tie points) are small compared to the uncertainties of the tie points, the inverse model does not add much value in this regard.

With regard to point (3) above, the consistent set of  $\Delta$ age, accumulation and thinning functions are valuable and may in themselves justify the use of the inverse model, but I'm not sure how well-constrained they are when only one core is being modelled and only gas age ties are used in the top and only ice age ties are used in the old part.

To summarize, I think the inverse model is almost overkill in this simple case. Provided that the differences between the inverse model and a simple interpolation are indeed small, the result of whole setup is an advanced transferral of the selected GICC05 and EDC time scales to the Talos Dome core using the visually selected tie points. There is nothing wrong with this, but the manuscript should reflect this and focus on the value of point (3) above.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

All this being said, I am confident that the presented age model is robust, and in this regard, the conclusions are justified. I acknowledge that the manuscript documents the dating methodology carefully, which is an important contribution, albeit it to some degree has the character of a technical report. I therefore recommend publication after the authors have satisfactorily dealt with the issues raised in the review. In line with the comments above, I encourage the authors to focus on point (3) above and if possible include more discussion of the results of the age model and the dated Talos Dome record and its relation to other records.

Detailed comments — 1734. 16: Bølling and Allerød should be spelled with ø. 16: The abrupt warming is at 14.6 ka BP (14692 b2k is 14642 bp1950). 16-17: The text should reflect that the dates given are for the onsets of the named periods, not the periods themselves. Also, the onset at 14.6 ka BP is the onset of Bølling, not Bølling-Allerød, as Allerød only starts some 700 years later (Lowe et al, INTIMATE protocol QSR 2008). 18: Severinghaus et al., 1998, sets the onset of the Holocene to 11.6 ka BP1950. 15-18: Consistency would increase if the dates of Greenland transitions were taken from one source, e.g. the GICC05 time scale. 22: Only a few hundred years? Which record is that? — 1735. First paragraph: mention that some of these differences could be related to dating. — 1736. 24: "Due to incorrect identification of missing seasonal signals and absence of absolute volcanic chronology before 1000 AD". Please clarify how one incorrectly identifies missing seasonal signals ... or rewrite. — 1738. 11: The authors may want to address the implications (and if possible, the magnitude) on this study of the concerns raised by Köhler recently (Clim. Past Discuss., 6, 1453-1471, 2010). 16: Add reference to Rasmussen et al, 2006 (JGR), which describes the dating in the interval 8-15 ka. The Vinther et al., 2006 (JGR), paper describes dating of the section above 8 ka, which is not used much here, but could be added for completeness. 20: Which EDC3 age model is used? The one with modifications by Lemieux-Dudon et al., 2009, or the original EDC3? And if not the first of these, why, given that this time scale is expected to be more consistent with GICC05 and the modelling approach applied? In general, the paper would benefit from consistent use of one or the other

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

EDC time scale and a discussion of which one is considered superior. — 1741. 12: It is not described how the authors estimate the uncertainty of the visual matching. This is an important shortcoming given that the result depends critically on these ties. Some kind of statistical modelling would be welcome to support the estimates, and the authors should at least describe in detail how they evaluate the synchronization uncertainty and how this value compares to the (variable) CH<sub>4</sub> data resolution. The different uncertainty contributions (esp. for the CH<sub>4</sub> tie points) should be listed individually as well as combined for each tie point. 14: The GICC05 uncertainties are not included in the error estimates of table 1, and I doubt they are for the EDC-derived tie points, either. This could be a perfectly reasonable approach (especially if the EDC age model used is the Lemieux-Dudon et al. version), with the resulting TALDICE model being a trade-off between a match to GICC05, a match to the used EDC age model, and the glaciological constraints. This would, I believe, follow the approach by Lemieux-Dudon, and imply that the TALDICE age model inherits any possible errors in the GICC05 time scale and EDC age model. This is also suggested in line 21 on page 1755, but contradicts the text here. — 1743. The authors could discuss here whether the chosen accumulation rate and temperature parameterisation realistically can capture both variations in A and T on short timescales and across glacial-interglacial transitions. I am aware that the accumulation rate is a free parameter of the more advanced inverse model of section 4, but deviations from the simple (background) model scenario are penalized (or at least, I guess they are, as described in Lemieux-Dudon et al., 2009), and it is thus essential that the a priori estimate is essentially correct. 19: The parameters  $p$  and  $\Delta H$  have not been introduced at this point. Elevation changes are discussed in section 3.2, but are not referred to as  $\Delta H$  in the text. — 1745. 3: Explain how/if the elevation changes are used in the full model and if the  $\delta D$  values are corrected accordingly before used for determination of past accumulation and/or temperatures. — 1746. 10: "(further studies are necessary to test this latter assumption)" is a strong understatement that would benefit from some qualified comments by the authors. Sec. 4.2: It seems to me that these rather crude assumptions about error

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

magnitude and correlation significantly reduce the value of the model. For example, the choice of constant correlation length parameters of 4000 yr / 50 m seems to be a stretch ... I would estimate that the errors are likely to have weaker (auto)correlation in periods of changeable climate conditions. I know that good estimates are not easy to obtain, but a more full discussion of the implications of these simplifications and – preferably – a sensitivity experiment that allows some quantification of the influence of these error magnitude and correlation estimates would greatly increase the reader's ability to assess the robustness and quality of the new age model. This is especially true if the model results are not very different from those of a simple interpolation (see general comments above). Why should we prefer a modelled time scale based on these simplifications more than a simple interpolation if the influence of the simplifications cannot be assessed? — 1747. 14: The line starting with "By comparison ..." is unclear. What is being compared, and what is "their" after the comma? 17: Strictly speaking, "The relatively small uncertainty" applies only to the MIS3 part, which is not clear from this line. 20: Clumsy sentence in the beginning. 22: "largely"? — 1748. 15: "The tie point assignment becomes more uncertain during this time interval and leads to larger changes in the thinning function deduced by the inverse method." If the tie point uncertainties appropriately reflected the larger uncertainties here, wouldn't one expect the model to produce a less varying thinning function at the cost of a less tight fit to the (more uncertain) age tie in question? — 1749. 7-24: These cases of good agreement are direct consequences of the tie point at  $14\,680 \pm 100$  yr BP used for the inverse model and a similar tie point used by Lemieux-Dudon et al., 2010. As both TALDICE-1 and the new EDC scales are tied very closely to GICC05, anything else than good correspondence so close to a tie point would be a sign of a huge problem ... my point is that this comparison cannot be used as an independent support of the validity of the inverse model, but is a direct consequence of a tie point with a low uncertainty assigned to it. Unless there is another (hidden) point with this comparison, I suggest that the section is removed. — 1750. As the resolution of the  $\delta^{18}\text{O}_{\text{atm}}$  profile in general is lower than the resolution of the data used for deriving tie points (maybe

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

except for the last glacial transition), the agreement does not add much information about the age model ... you can say that the agreement confirms that the tie points are overall correct (e.g. that the right stadials and interstadials have been matched together), but apart from that, this agreement cannot be used to validate the model. In summary, section 5.1.3 is very weak. The data presented are perfectly reasonable but cannot be used for a confident evaluation of the quality of the age model as the authors claim. — 1751. 15: The authors may want to note that the biggest difference between the resulting acc. rates and the simple model (purple) occurs over the 18-30 ka interval, in part of which also the simple Greenland  $\delta^{18}\text{O}$  – accumulation relationship breaks down (Svensson et al., 2006, QSR). — 1754. Whole section and Fig. 7: I would prefer the consistent use of inverse thinning function or just thinning function, not a mix. Also, I would suggest that the authors decide whether they trust the a priori thinning function from the ice-flow model that does or does not take into account altitudinal changes (and argue why), and consistently compare the fabric curve with that curve only. 7: I'm definitely not an expert on this, but the range is 1-3 in the text and 0.2-1 in the figure. Please correct / explain / make consistent. 12: "Around 700–750m (11.5–12.4 kyr BP) the fabric evolution shows an increasing slope (yes) at the time when the ice-flow and inverse thinning functions start to diverge in their main trend (only for the – unrealistic? – case of no altitudinal changes – the inverse model fits well with the solid grey line)" 15: "a clear increase in the rate of fabric clustering appears" ... could this "clear increase" not be caused by one low value (ca. 825 m)? Given that there is quite some noise in the record at this depth, I find the correlations between the fabric orientation and thinning function curves to be very bold. 24-25: I simply have no clue what this line means. 26: "Around 1100–1150m (42.8–46.6 kyr) both the fabric and the thinning function evolution record an increasing rate of change". I would say that the fabric curve is almost constant ... the changes happens below 1150 m. — 1755. 3: "could"? Do the data show this or not? In summary, my impression as a non-specialist in fabric analysis is that this section is based on rather bold interpretations that rely on the determination of changes of slopes of very short curve segments where error

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

on just one point can change the picture. I am not convinced about the validity of the conclusions, and recommend that the editor seeks expert advice. 5: rephrase ... especially "... come now to ..." — 1756. 2: Rephrase "and makes a clear bonus to use the" — Table 5. Rephrase "bibliographic descriptions". — Figures. The clarity of figures would benefit from a homogenous font size, consistent use of yr/kyr, and consistent use of labels A, B, etc. on both (sub-)figures and in captions. A use of a more diverse colour scheme (rather than shades of blue) would ease interpretation, as would legends on the figures, so that the reader can see what is presented by the different curves without having to read the caption for colour definitions. — Fig. 1: Make figure full width. — Fig. 4: Is there an explanation for the offset of the peaks at ~250 kyr BP? — Fig. 5: Mention if elevation change corrections have been applied. Write "atm" in subscript in label. — Fig. 6: Mention if elevation change corrections have been applied in any of the three cases presented. It is not clear how the 10Be stars relate to the explanation and accumulation rate ratios on page 1752-53. — Fig. 7: a1 has a strange font size/subscript in label. There is no mentioning of the 950 m grey-shaded section in the text.

---

Interactive comment on *Clim. Past Discuss.*, 6, 1733, 2010.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)