

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



³Laboratoire d'étude des Transferts en Hydrologie et Environnement, UMR5564, Université Joseph Fourier,-Grenoble 1, CNRS, G-INP, IRD, 38402, Grenoble, France
⁴Instituto de Investigaciones Geológicas y del Medio Ambiente (IGEMA), Universidad Mayor de San Andrés (UMSA), Calle 27, Pabellón 3, Campus Universitario Cota Cota, Casilla 35140, La Paz, Bolivia

Received: 29 May 2011 – Accepted: 8 June 2011 – Published: 24 June 2011

Correspondence to: P.-H. Blard (blard@crpg.cnrs-nancy.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

This paper describes several simple positive degree-day models (hereafter referred as “PDD models”) designed to provide past climatic reconstruction from tropical glacier paleo-equilibrium altitude lines (paleo-ELA). Several ablation laws were tested and calibrated using the monthly ablation and meteorological data recorded from 1997 to 2006 on the Zongo glacier (Cordillera Real, Bolivia, 16° S). The performed inversion analyses indicate that the model provides a better reconstruction of the mass balance if the ablation is modeled with different melting factors for snow and ice. The inclusion of short-wave solar radiations does not induce a substantial improvement. However, this type of model may be very useful to quantify the effects of local topographic (orientation, shading) and to take into account incoming solar radiation changes at geological timescale. The performed sensitivity test indicates that, in spite of the uncertainty in the calibrated snow-ice ablation factors, all models are able to provide paleotemperatures with $\sim 1^\circ\text{C}$ uncertainty for a given paleoprecipitation. This error includes a 50 m uncertainty in the estimate of the paleoELA. Finally, the models are characterized by different precipitation-temperature sensitivities: if a similar warming is applied, model including different ablation factors for snow and ice requires a lower precipitation increase (by $\sim 15\%$) than others to maintain the ELA.

1 Introduction

The physical mechanisms controlling the mass balance of mountain glaciers are primarily determined by the climatic conditions in the accumulation and ablation area (e.g. Ohmura et al., 1992). Additionally, because of their relative small size, alpine glaciers respond quickly (at the timescale of 10^0 to 10^1 yr, which is fast compared to the historical and geological timescales) to changes in main atmospheric variables, such as temperature or precipitation (e.g. Oerlemans, 2005). These two characteristics imply that mountain glaciers are very useful and precise proxies for paleoclimatic

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reconstructions (e.g. Blard et al., 2009; Flowers et al., 2007; Mark et al., 2005; Oerlemans, 2005). This is particularly true in high elevation areas where other climatic archives (such as pollen records) are scarce or inexistent. Because paleo-glaciers are by definition vanished, this climatic reconstruction must however be done through numerical models in which past glacial extents are considered as inputs and paleotemperature and paleoprecipitations as outputs. Accurate reconstruction of these main paleoclimate variables requires a proper modeling of the past glacier mass-balance (mass balance = accumulation – ablation). The computation of accumulation is generally quite straightforward (e.g. Hock, 2005; Lejeune et al., 2007; Vincent, 2002), so the main difficulty of paleoclimate studies is to model properly the ablation of ice and snow at the glacier surface.

In the recent decades, a large variety of melt models have been developed to link climatic variables with ablation ranging from (i) simple temperature-index models, also called positive-degree-days (PDD) models (e.g. Braithwaite, 1995; Braithwaite and Olesen, 1985; Hock, 1999, 2005; Johannesson et al., 1995; Vincent, 2002) that are based on the empirical relationship existing between ablation and the sum of positive air temperatures, to (ii) sophisticated energy balance models (e.g. Anslow et al., 2008; Brun et al., 1989; Greuell and Konzelmann, 1994; Hock and Holmgren, 2005; Klok and Oerlemans, 2002; Molg et al., 2003) that compute melt from the energy flux budget at the glacier surface.

Given the complexity and the large variety of surface processes involved in the mass balance of tropical glaciers, the most accurate models are those based on an energy balance-budget. However, these models generally require a large number of input data, such as, i.e., wind velocity, relative humidity, cloud cover, all these parameters being generally difficult to know over geological timescales, even for the Quaternary period (last 2 million years). Thus, although energy balance models are physically more relevant than PDD models, this sort of model is generally difficult to use in paleoclimatic studies. This issue could however be overcome through careful sensitivity tests, but such approach is time-consuming and can yield significant uncertainties

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and 5200 m) and to the East in the lower part (between 5200 and 4900 m) (Sicart et al., 2005, 2007; Wagnon et al., 1999). In 2006, the Zongo glacier itself covered about 1.96 km², for a total length of 3 km and a width of 0.75 km (Soruco et al., 2009) (Fig. 1). The Zongo has the characteristics of a temperate glacier (Francou et al., 1995). The hydrological year has been defined as starting on 1 September and ending on 31 August (Ribstein et al., 1995). During the wet season (summer, from September to March) solid accumulation and ice-snow melt occurs simultaneously. Contrarily, during the dry season (winter, from April to August) solid accumulation is sparse, and ice-snow melt is reduced but sublimation is significant (Favier et al., 2004).

2.2 Meteorological data

The meteorological data of the nine hydrological years studied in this article (Table 1) were collected by two weather stations located in the vicinity of the Zongo glacier. Locations of the weather station are shown on Fig. 1. For the 6 hydrological years ranging between 1997 and 2003, monthly temperature and precipitation data were recorded by the MEVIS station, which is located at an elevation of 4750 m, at a distance of approximately 1 km from the glacier terminus (16°16.9' S, 68°7.3' W). The climatic data for the 3 hydrological years between 2003 and 2006 are from a weather station located at 5050 m (16°17' S, 68°8.3' W), on the lateral moraine of the Zongo glacier (ORE GLACIOCLIM network), but far enough from the ice to avoid major perturbation due to change in the glacier surface conditions. Moreover, since the present model aims to infer regional temperatures from the inversion of past glacial extents, it is relevant to calibrate the model using local temperatures measured out of the Zongo glacier. Mean annual temperature is 0.4 ± 0.3 °C at 5050 m (ORE station – from 2003 to 2006) and 2.0 ± 0.7 °C at 4750 m (MEVIS station – from 1996 to 2003) (Table 1). The average annual amplitude is 3 ± 1 °C. Temperature standard deviations around monthly means are calculated from the high resolution time series (30' frequency record) of the ORE network recorded during the three hydrological years between 2003 and 2006. This standard deviation is used as an input data in the PDD model, it is used to calculate

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

a thin layer of oil to avoid evaporation and of opening area of 2000 cm²) has been manually measured every month since 1996. There is a good agreement between the corrected Geonor monthly precipitation and the monthly precipitation measured using the totalizer for both hydrological years 2004–2006 ($R^2 = 0.94$), but with a systematic under-estimation of 20 % for the totalizer. Consequently, for monthly precipitation before September 2004, we used the totalizer measurements corrected by a factor 1.2. Moreover, precipitation data recorded from the snow pits located at different elevations on the Zongo glacier (Fig. 1) do not indicate significant change in precipitation against elevation. The precipitation data measured on the moraine at 5050 m were thus used over the whole elevation range over which the Zongo mass balance is modeled (i.e. between 4900 and 6000 m).

Mean annual precipitation range from 1044 to 1529 mm yr⁻¹ for the 1997–1998 and 2000–2001 hydrological years, respectively (Table 1). All years are characterized by significant precipitation seasonality. Indeed, more than 70 % of precipitation on the Zongo occurs between November and March (Table 1). This seasonality arises from the location of the Cordillera Real: indeed, this mountain range belongs to the outer-tropics, in a zone of the Andes that is very sensitive to the South American monsoon (Garreaud et al., 2009).

2.3 Mass balance data

Mass balances of Zongo glacier are available from glaciological (since 1991 to 2006) hydrological (from 1975 to 2006) and geodetic methods (in 1956, 1963, 1975, 1983, 1997 and 2006) (Soruco et al., 2009). Annual mass balances of Zongo glacier have been systematically measured since 1991 (Francou et al., 1995). However, only the period from 1997 to 2006 have been corrected using the geodetic method by photogrammetry (Soruco et al., 2009). Therefore, for the present study, we have only used the mass balances data monitored over these 9 hydrological years. Only a short description of the glaciological data acquisition is given here, because the methods have

already been thoroughly described in previous articles (Francou et al., 1995; Sicart et al., 2007; Soruco et al., 2009).

The traditional glaciological method (Paterson, 1994) applied to the Zongo glacier aims to estimate the loss or gain in mass over the whole glacier surface from field measurements. Annual mass balances have been obtained using in average 14 stakes of monthly measurements over the ablation zone and 3 snow pits and drilled cores for yearly measurements over the accumulation zone. Moreover, snow height measurements are also required in the ablation area, because snow accumulation can occur at any time in the year over the entire glacier. These measurements were performed assuming that snow and ice densities are 400 and 900 kg m⁻³, respectively (Soruco et al., 2009). In the accumulation area, the snow density of the snow pits and drilled cores were measured at the end of the hydrological year. Finally, the specific glacier mass balances were computed using actual surface areas derived from photogrammetric measurements (Soruco et al., 2009).

The annual mass balance data were finally calculated by summing the monthly data for the 9 hydrological years between 1997 and 2006 (Table 2).

3 Description of the models

3.1 General considerations

The present study aims to evaluate the efficiency of 3 different types of models. Models 1 and 2 belong to the same category: both are derived from the classical PDD model (Braithwaite and Olesen, 1985). Model 3 is inspired from the enhanced versions developed by (Hock, 1999) and model 4 is a derivative of the model of (Pellicciotti et al., 2005). These 2 last types of models have the particularity to include the short wave solar radiations. The main difference between our study and these previous works is that the computation is here performed at a monthly timescale. Indeed, the Zongo mass balance data are only available at this temporal resolution. Moreover, the main motivation

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of the study is not to provide a detailed modeling of ablation at an hourly time scale, but rather to test a robust and simple approach to derive general paleoclimatic conditions from tropical glacier paleoELAs. Finally, (Sicart et al., 2008) has demonstrated that temperature-index models are not appropriate to simulate daily melting but that they can be relevant to describe the ablation of tropical glaciers at longer timescales (monthly or yearly). Although the model are run on a monthly time-step, all ablation parameters are reported here as daily values, to allow a direct comparison with the ablation parameters obtained from other calibration studies.

For all models, the required input data and the calibrated parameters are described in Table 3.

All models share a common structure: monthly snow accumulation, and snow/ice ablation are computed each 50 m over the relevant elevation range (from 4950 to 6050 m for the Zongo). Annual accumulation and ablation are calculated by summing the monthly values. Finally, the mass balance elevation law, MB (mm w.e. yr⁻¹) is computed at each elevation subtracting ablation (A in mm w.e. yr⁻¹) from accumulation (S in mm w.e. yr⁻¹):

$$MB = S - A \quad (1)$$

3.2 Modeling of snow accumulation

The annual snow accumulation, S (mm w.e. yr⁻¹), is computed in a similar way in all models. S is the sum of the monthly snowfalls:

$$S = \sum_{i=1}^{12} S_i \quad (2)$$

S_i (mm w.e. month⁻¹) is calculated for each month assuming precipitation falls as snow when temperature is below the rain-snow threshold T_s . The model assumes

that temperatures have a normal distribution around the monthly mean temperature T_{mi} , with a standard deviation σ_i (Johannesson et al., 1995):

$$S_i = P_i \frac{1}{\sigma_i \cdot \sqrt{2\pi}} \int_{-\infty}^{T_s} e^{-\frac{(T-T_{mi})^2}{2\sigma_i^2}} \cdot dT \quad (3)$$

where P_i (mm w.e. month⁻¹) is the monthly precipitation. Lejeune et al. (2007) has shown that, in the tropical Andes, 100 % of falls is snow below -1°C , and that 100 % is rain above 3°C . T_s is thus supposed to be 1°C in the following.

3.3 Modeling of ablation

Annual snow ablation, A (mm w.e. yr⁻¹), is computed in a similar way in all models. A is the sum of the monthly ablation:

$$A = \sum_{i=1}^{12} A_i \quad (4)$$

3.3.1 Calculation of the monthly positive temperature T_{Pi}

In each ablation model described below, the monthly positive temperature T_{Pi} ($^\circ\text{C}$) is calculated following the positive degree-day approach, which assumes that temperature follows a normal distribution characterized by the monthly mean T_{mi} ($^\circ\text{C}$) and the standard deviation σ_i ($^\circ\text{C}$). The average positive temperature is thus computed as the integral of this distribution function on the interval $(0, +\infty)$ $^\circ\text{C}$ (Johannesson et al., 1995):

$$T_{Pi} = \frac{1}{\sigma_i \cdot \sqrt{2\pi}} \int_0^{\infty} e^{-\frac{(T-T_{mi})^2}{2\sigma_i^2}} \cdot dT \quad (5)$$

3.3.2 Model 1: $A = MF \times T_p$

This model is the classical PDD model (e.g. Braithwaite and Olesen, 1985). In this high simplicity model, only one melting factor is involved, MF (mm w.e. °C day⁻¹). Monthly ablation is computed as:

$$A_j = \frac{365}{12} \times MF \times T_{Pi} \quad (6)$$

3.3.3 Model 2: $A = MF_{\text{snow/ice}} \times T_p$

In this model ablation of snow and ice is calculated with different degree-day factors:

- MF_{snow} (mm w.e. °C⁻¹ day⁻¹) for snow (if the glacier surface is snow).
- MF_{ice} (mm w.e. °C⁻¹ day⁻¹) for ice (if the glacier surface is ice).

At each monthly time step, total ablation, A_j , is the sum of snow ablation and ice ablation:

$$A_j = A_{\text{snow}_j} + A_{\text{ice}_j} \quad (7)$$

Ice melting only occurs if the ablation of snow was sufficient to remove all the snow cover. In order to correctly quantify the fraction of time during which the ablation of snow occur each month, the model first calculates a parameter α that is defined as:

$$\alpha = \frac{A_{\text{snow}_j}}{A_j} \quad (8)$$

As, for each month, the amount of total snow input is known (S_j), “ α ” is in practice computed as follow:

$$\alpha = \frac{12}{365} \times \frac{S_j}{MF_{\text{snow}} \times T_{Pi}} \quad (9)$$

- If $a > 1$, the snow ablation is smaller than accumulation, thus, in this case, only partial melting of the snow cover occurs. Monthly ablation A_j is then:

$$A_{\text{snow}j} = \frac{S_j}{\alpha} \quad (10)$$

- If $a < 1$, the total monthly ablation A_j is sufficient to remove all the snow cover and start melting the glacier ice. In this case, A_j is then calculated as:

$$A_j = S_j + (1 - \alpha) \times \frac{365}{12} \times \text{MF}_{\text{ice}} \times T_{\text{Pi}} \quad (11)$$

3.3.4 Model 3: $A_{\text{snow}} = \text{MF} \times T_p$ and $A_{\text{ice}} = (\text{MF} + a \times I) \times T_p$

This model is inspired from the one developed by (Hock, 1999). As in model 2, different melting factors are used for snow and ice. The particularity of this model is to incorporate the clear-sky short wave solar radiations. Similarly to the classical PDD model, ablation only occurs under positive temperature.

In this model, snow ablation is calculated following:

$$A_{\text{snow}j} = \frac{365}{12} \times \text{MF} \times T_{\text{Pi}} \quad (12)$$

Ice ablation is obtained from:

$$A_{\text{ice}j} = \frac{365}{12} \times (\text{MF} + a \times I_j) \times T_{\text{Pi}} \quad (13)$$

where MF (mm w.e. °C⁻¹ day⁻¹) is the temperature melt factor (MF is similar for snow and ice in model 3), a (mm w.e. d⁻¹ °C⁻¹ W⁻¹ m²) is the radiation-temperature melt factor and I_j (W m⁻²) is the monthly average short wave solar radiation. In the present

study, I_j was calculated assuming clear-sky conditions, according to the equation (Hock, 1999):

$$I_j = I_0 \left(\frac{R_m}{R_i} \right)^2 \psi \left(\frac{P}{P_0 \cdot \cos Z_i} \right) \cdot \cos \theta \quad (14)$$

where I_0 is the solar constant (1368 W m^{-2}), R_i the mean monthly distance Sun-Earth, R_m the mean Sun-Earth distance, $\psi = 0.75$ the atmospheric clear-sky transmissivity, P and P_0 are respectively the local and sea-level atmospheric pressure, Z_i the mean monthly zenith angle and θ the angle of incidence between the normal to the grid slope and the solar beams. θ was calculated for each altitude step (between 4950 and 6050 m) by using a 30 m digital elevation model of the Zongo surface. This type of models (i.e. models 3 to 4) thus takes into account the glacier geometry to compute the actual amount of incident solar beams reaching the glacier surface.

In practice, the model follows the same algorithm as Model 2 to compute, for each month, the respective proportion of snow and ice melting: $\alpha = A_{\text{snow}} / (A_{\text{snow}} + A_{\text{ice}})$.

3.3.5 Model 4: $A = \text{Mf} \times T_p + a \times I$

This model is directly derived from the one proposed by (Pellicciotti et al., 2005). The slight difference with model 3 is that the ablation arising from solar radiation, I , is fully independent from temperatures. The particularity of the model presented here is to allow ablation, even when temperatures are negative (i.e. $T_p = 0^\circ\text{C}$).

$$A_i = \frac{365}{12} \times (\text{MF} \times T_{Pi} + a \times I_j) \quad (15)$$

a is the solar radiation melting factor expressed in $\text{mm w.e. d}^{-1} \text{ W}^{-1} \text{ m}^2$. I is the short wave incoming solar radiation calculated as for model 3 (Eq. 14, Sect. 3.3.4).

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Calibration of models' parameters

4.1 Optimization method

For each hydrological year, the best-suited models parameters were determined by minimizing the efficiency criterion R^2 . This criterion is calculated as follow (Nash and Sutcliffe, 1970):

$$R^2 = 1 - \frac{\sum_{i=1}^n (\text{MB}_{\text{obs}}(i) - \text{MB}_{\text{sim}}(i))^2}{\sum_{i=1}^n (\text{MB}_{\text{obs}}(i) - \overline{\text{MB}_{\text{obs}}})^2}, \quad (16)$$

where MB is the mass balance, the subscripts “obs” and “sim” denote “observed” and “simulated”, respectively and i denotes the elevation step.

In practice, the parameters optimization was realized for all models and each hydrological years following the method described in (Hock, 1999). The R^2 values are mapped as a function of the considered ablation parameters value (Fig. 2). The retained best parameters are those yielding maximum R^2 values. Note that, for each of the 4 models, we also calculated the best R^2 for the whole 9 hydrological years (from 1997 to 2006) taken together. This approach permitted to obtain the most relevant ablation parameters for the whole dataset.

4.2 Results: calibrated parameters

All the calibrated parameters from the 9 hydrological years are displayed in Table 4. Figure 3 shows the plot of simulated and observed mass balance vs elevation for each of the 4 models, for the three hydrological years 1997–1998, 2003–2004 and 2004–2005. Figure 4 displays a plot of the simulated vs. observed mass balances data for each of the 4 models applied to the nine hydrological years (1997–2006).

4.2.1 Model 1: $A = MF \times T_p$

The calibrated melting parameters MF range between 9 (2003–2004) and 15.5 mm w.e. °C⁻¹ d⁻¹ (1998–1999). The efficiency criterion R^2 ranges between 0.77 and 0.99, which underlines that, despite its simplicity, this model is quite efficient in reproducing the observed mass balance (Table 4, Figs. 3 and 4). The altitudinal gradient of the simulated mass balance is indeed in very good agreement with the observed one (Fig. 3). Besides, tropical glaciers are characterized by much stronger vertical mass balance in their ablation area than other glaciers (e.g. Kaser, 2001; Wagnon et al., 1999) and the fact that this gradient is well simulated gives confidence in the ability of the model to reproduce the glacier mass balance. Similarly, the modeled and observed ELA are generally closely matching, within the observation uncertainty (<50 m). The model optimization performed from the whole dataset yielded a MF of 11.9 ± 1.3 mm w.e. °C⁻¹ d⁻¹, with a R^2 of 0.92.

4.2.2 Model 2: $A = MF_{\text{snow/ice}} \times T_p$

The calibrated MF_{snow} melting parameters range between 1.1 (2005–2006) and 23 (2003–2004) mm w.e. °C⁻¹ d⁻¹. MF_{ice} parameters range between 7.6 (2003–2004) and 25.7 (2005–2006) mm w.e. °C⁻¹ d⁻¹. Ice ablation parameters are systematically higher than those of snow (except for 2003–2004), which is consistent with the fact that, because of different albedo, ice melting is faster than snow melting. R^2 value range from 0.83 to 0.99, which indicates that model 2 is slightly more efficient than model 1 to match the observed mass balances (Fig. 3). The calibration performed from the whole dataset yielded 8.7 ± 0.6 mm w.e. °C⁻¹ d⁻¹ for MF_{snow} and 12.7 ± 1.4 mm w.e. °C⁻¹ d⁻¹ for MF_{ice} , with a R^2 of 0.93.

4.2.3 Model 3: $A_{\text{snow}} = \text{MF} \times T_p$ and $A_{\text{ice}} = (\text{MF} + a \times \text{I}) \times T_p$

Calibrated values for MF range between 0.8 and 12 mm w.e. $^{\circ}\text{C}^{-1} \text{d}^{-1}$. The solar-radiation parameter “ a ” is characterized by larger interannual fluctuations, from 0 to 12.6×10^{-2} mm w.e. $^{\circ}\text{C}^{-1} \text{W}^{-1} \text{m}^2 \text{d}^{-1}$. The efficiency criterion R^2 ranges from 0.82 to 0.99, which indicates that this model structure has also a good ability to fit the observed mass balance. However, the R^2 obtained from the whole dataset (0.93) is not significantly better than those of models 1 and 2. Thus the introduction of the parameter “ a ” did not result in a significant improvement of the model efficiency (Fig. 3). The “all-years” calibration yielded a value of 8.8 ± 1.0 mm w.e. $^{\circ}\text{C}^{-1} \text{d}^{-1}$ for MF and $0.92 \pm 0.21 \times 10^{-2}$ mm w.e. $^{\circ}\text{C}^{-1} \text{W}^{-1} \text{m}^2 \text{d}^{-1}$ for “ a ”, with a R^2 of 0.93.

4.2.4 Model 4: $A = \text{MF} \times T_p + a \times \text{I}$

Parameterized MF range between 8 and 15.4 mm w.e. $^{\circ}\text{C}^{-1} \text{d}^{-1}$. Best “ a ” parameters range between 0 and 20×10^{-4} mm w.e. $^{\circ}\text{C}^{-1} \text{W}^{-1} \text{m}^2 \text{d}^{-1}$. R^2 range from 0.77 to 0.99. The “all-years” calibration yielded a value of 11.8 ± 1.3 mm w.e. $^{\circ}\text{C}^{-1} \text{d}^{-1}$ for MF and $2.1 \pm 0.5 \times 10^{-4}$ mm w.e. $^{\circ}\text{C}^{-1} \text{W}^{-1} \text{m}^2 \text{d}^{-1}$ for “ a ”, with a R^2 of 0.93.

5 Discussion

5.1 Inter-model comparison

All models including the solar radiations (model 3 and 4) are characterized by a significant interannual variability of the “ a ” factors calibrated from each hydrological years (Table 4). Two main explanations could be invoked to explain such a large discrepancy: first, these low complexity models do not take directly into account the fluctuations of cloudiness, nor those of albedo. Only clear-sky radiations (input I in W m^{-2}) are considered in Model 3 and 4. Under cloudy conditions, the true incoming short

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

reproducing the observed mass balance (Fig. 3). “All years R^2 ” of model 2, 3 and 4 are equal to 0.93, respectively. Consequently, the introduction of solar radiations or the use of different melting factors did induce only a slight improvement of the model performance (Figs. 3 and 4, Table 4).

Correlation coefficients have also been computed considering separately the positive and negative mass balances (R_{pos}^2 and R_{neg}^2 , respectively). These efficiency criteria allow a detailed diagnostic of the ability of each model to fit the data in the accumulation or in the ablation zones (Table 4). For all models R_{neg}^2 (0.87) are significantly better than R_{pos}^2 (0.48 to 0.50), which shows that these models are more efficient to model the mass balance in the ablation zone than in the accumulation zone.

R_{neg}^2 are similar for each of the 4 models, suggesting that all of them have a comparable ability in fitting the data in the ablation zone. However, R_{pos}^2 are slightly different between each model. The highest value for R_{pos}^2 is obtained with model 4 ($R_{\text{pos}}^2 = 0.50$). This is probably because this model allows ablation in the accumulation zone. Indeed, Fig. 3 indicates that model 4 offers a much better fit between model and data above the ELA, while all of the 3 other models systematically overestimate the observed mass balance in the accumulation zone. Indeed, models 1 to 3 cannot yield ablation under negative temperatures. This observation is also confirmed by the slope of the regression line determined plotting the observed against the calibrated mass balance (Fig. 4): indeed models 1, 2 and 3 are characterized by a slope that is smaller than 1, while the MB simulated from model 4 define an almost perfect 1:1 line.

By comparing the efficiency of different type of models on a glacier in Switzerland, (Pellicciotti et al., 2005) obtained very similar results: models accounting separately the influence of temperature and short-wave radiations perform better. This characteristic may be potentially useful to take into account ablation mechanisms such as sublimation or snow erosion by wind. Although it is very difficult to parameterize these mechanisms, the possibility to take into account these processes in low-complexity models such as model 4 may represent a significant improvement of classical PDD models.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

It is thus tempting to suggest that model 4 should be favored for next climatic studies based on tropical glaciers. We also tested model having different “*a*” parameters for snow and ice, but this modification did not induce any improvement in the model performance and thus these models are not described in the present study. Moreover, considering solar-radiations in the ablation calculation (model 3 and 4) can be very useful in certain situations. First, this type of model is able to take into account specific effects, such as the influence of the glacier orientation, slope or shading in entrenched valleys. The way the solar radiation is calculated in the ablation equations of model 3 and 4 indeed permits to incorporate geometrical effects that can potentially modulate the incoming short wave radiation. These models can thus be used as efficient tools to evaluate and quantify the influence of topographic effects. Second, these models have the ability to incorporate the past fluctuations of solar radiations resulting from orbital forcing. This aspect is important for paleoclimatic reconstructions.

5.2 Comparison with previous calibration studies

5.2.1 Models 1 and 2

As previous calibration studies from tropical glacier are almost inexistent, it is tough to compare directly the temperature-index factors obtained here from the Zongo glacier with previously published melting index. The closest-to-equator calibration studies were located in the Himalaya (Kayastha et al., 2000; Singh and Kumar, 1996; Singh et al., 2000) between 27°N and 35°N latitude. In a review paper, (Hock, 2003) showed that these Himalayan glaciers yielded ablation factors ranging between 5.9 and 11.6 mm °C⁻¹ d⁻¹ (average 7.8 ± 2.4 mm °C⁻¹ d⁻¹) for snow and from 6.6 to 16.9 mm °C⁻¹ d⁻¹ (average 9.6 ± 3.4 mm °C⁻¹ d⁻¹) for ice respectively. These values are not significantly different from those obtained here from model 1 (MF = 11.9 ± 1.3 mm °C⁻¹ d⁻¹) and model 2 (MF_{snow} = 8.7 ± 0.6 mm °C⁻¹ d⁻¹ and MF_{ice} = 12.7 ± 1.4 mm °C⁻¹ d⁻¹). Some of the MF_{ice} factors obtained here are higher than previously determined ice ablation factors. However, this observation should be

considered with caution given the highest MF factors obtained in the present study are all determined from the 1997 to 2002 hydrological years. Indeed, these years of observations might be potentially affected by a lower reliability of the temperature data from the MEVIS stations.

5.2.2 Model 3

This model is very close to the one used in our study dedicated to the last glaciation of the Hawaiian summits (Blard et al., 2007). In this initial study, model parameters were calibrated from the 2003-2005 hydrological years record from the Zongo glacier. This study yielded optimized values of $3.7 \text{ mm } ^\circ\text{C}^{-1} \text{ d}^{-1}$ for MF_{snow} , $7.4 \text{ mm } ^\circ\text{C}^{-1} \text{ d}^{-1}$ for MF_{ice} and $\sim 4 \times 10^{-3} \text{ mm } ^\circ\text{C}^{-1} \text{ W}^{-1} \text{ m}^2 \text{ d}^{-1}$ for “a” (Blard et al., 2007). The new MF parameters are comparable, but the new “a” parameters is twice larger than the one obtained from the initial calibration study of (Blard et al., 2007) (Table 4). This discrepancy probably results from the fact that the 2 studies were performed by using different calibration years: 9 hydrological years (1997 to 2006) are used in the present calibration study, while only 2 yr (2003 to 2005) were considered in (Blard et al., 2007). More important, the precipitation data used here were significantly revised (by $\sim +20\%$) to account for a bias in the precipitation gauge that was not considered in (Blard et al., 2007).

The calibration of this type of model from glaciers in Sweden (Hock, 1999) and in Switzerland (Pellicciotti et al., 2005) also yielded ablation factors in the same order of magnitude than those obtained here.

5.2.3 Model 4

Only one study tested this model at an hourly timescale on a mountain glacier in Switzerland (Haut Glacier d’Arolla) (Pellicciotti et al., 2005). The MF factors calibrated in this study are significantly higher (about $\times 10$) than those obtained by Pellicciotti et al. (2005), while the “a” factors are significantly lower. However, this comparison may

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be considered with caution since Pellicciotti et al. (2005) used measured net radiation and not clear-sky radiations.

The ablation processes of high latitude glaciers are quite different than those occurring in the Tropics (Hock, 2003; Sicart et al., 2008). These discrepancies in calibrated parameters may thus reflect differences in the involved physical ablation processes. This underlines that any paleoclimatic or projection studies should benefit from using parameters calibrated in similar conditions than those of the studied zone, as already suggested by Hock (2003).

5.3 Physical limitations of the temperature-index models

In spite of any improvement that could be made, these PDD models remain empirical models. They are not based on a real energy balance approach and they have not a good ability in catching the complex interactions between the glacier surface and the atmospheric variables (relative humidity, albedo, cloudiness, wind velocity) at a short timescale (Sicart et al., 2005, 2008). However, the goal of the present study is to test and valid ablation models for paleoclimatic reconstruction, an exercise that requires keeping as low as possible the number of input data. The different models presented here should thus be used with caution, especially if the goal is to model the glacier mass balance under very different atmospheric conditions. For example, in very dry and low precipitation conditions, at high elevation (above 5500 m in the dry tropical Andes) sublimation becomes the main ablation process (Favier et al., 2004; Rupper and Roe, 2008; Wagnon et al., 2003). In the case of the Zongo glacier (precipitation $> 1 \text{ m yr}^{-1}$), the contribution of sublimation on total ablation is less than 10% (Fig. 6 in Rupper and Roe, 2008). Sublimation however becomes the major ablation process under drier conditions. Indeed, the sublimation/melt ratio is higher than 1 when precipitation is lower than 0.5 m yr^{-1} (Rupper and Roe, 2008). In that particular case, classical PDD models (1 and 2) and models 3 will not be able to correctly capture the real amount of ablation. The structure of model 4 clearly represents a more realistic approach to calculate the mass balance of tropical glaciers fed by very low precipitations.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ideally, the calibration of the “*a*” parameter of model 4 should be done using mass balance data from a glacier developed in a drier tropical zone. Unfortunately, there is a lack of mass balances records in such areas.

5.4 Sensitivity tests: precipitation-temperatures curves

5 We used the 4 calibrated models to determine the precipitation-temperature solution able to reproduce the present day-observed ELA at 5400 m (Fig. 6). This calculation was done using an “average year” calculated from the temperatures and precipitation data of the 9 hydrological years reported in Table 1. A 3°C range in annual average temperature (+1.5°C and –1.5°C the present value) was explored to determine the
10 corresponding precipitation conditions. This was done to evaluate (i) the uncertainty arising from the calibrated parameters and (ii) the precipitation-temperature sensitivity of the models. The 4 obtained precipitation-temperature curves are characterized by different slopes, which reflects contrasted sensitivities of the models. Under warmer conditions (+1.5°C), models including different ablation factors for snow and ice (Model 2 and 3) require lower precipitation (×3.2 for instead of ×3.7) to maintain the ELA at
15 5400 m a.s.l. than those having similar ablation factors for snow and ice. A comparison of models 1 and 2 with models 3 and 4 indicates that including solar radiation does not significantly change the precipitation temperature sensitivities. The error arising from the propagation of the parameters uncertainty is not significant. Indeed, even for
20 models characterized by a high factor uncertainty (Model 3), the error envelop indicates that the temperature uncertainty remains below 1°C, for a given precipitation.

The shape and slope of these obtained precipitation temperature curves can be compared with similar curves established by previous works. The curves yielded by the 4 models tested here match quite well with the curve established by Shi (2002) by
25 plotting the summer temperature vs the annual precipitation measured at the ELA of several mid-latitudes glaciers from the Alps and Western China. Our models yield a dP/dT slope that is slightly lower than the average curve built by Ohmura et al. (1992) from a large worldwide dataset. The P vs. T curves predicted by our models remains

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



however consistent with the precipitation-temperature sensitivity range reported by this author, suggesting that our approach does not yield unrealistic results.

6 Conclusions

The ablation parameters of several glacier-climate models were calibrated by using monthly time-scale meteorological and mass balance data from the Zongo glacier (Bolivia, 16° S). The majority of the so-obtained degree-day ablation factors are within the range of other values obtained from other worldwide glaciers (Hock, 2003). The “short wave ablation factors” of models including solar radiation are quite different from the parameters calibrated from mid latitude glaciers (Hock, 1999; Pellicciotti et al., 2005).

The models presented here are based on a semi-empirical approach, which hampers precise and accurate physical interpretations. In particular, it is important to keep in mind that classical temperature-index models are not suited for high elevation (> 6000 m) tropical glaciers, where wind erosion and sublimation are the dominant ablation mechanisms (Wagnon et al., 2003). However, (i) our study shows that the inclusion of different melting parameters for snow and ice only induce a slight improvement of the model ability in fitting the observed mass-balance. Calibrated ablation parameters of snow are systematically smaller than those of ice, suggesting that the effect of albedo is, at least partially, accounted by the model structure. (ii) Models with short wave radiations offer the advantage to include a quantitative estimate of spatial and temporal changes in solar radiations. This property can be of major interest to quantify the effects of shading (e.g. in deeply incised valley) or the influence of orbital insolation changes.

Reconstructed precipitation-temperature curves able to fit a given ELA show that these models are able to yield temperature reconstructions with $\sim 1^\circ\text{C}$ uncertainty. However, the temperature-precipitation sensitivity can be different between each model: models with different ablation parameters for snow and ice require lower precipitation at the same temperature.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The performance of these empirical models could certainly be improved by further studies, notably by considering new and larger dataset acquired under different atmospheric and climatic conditions. Another way to improve these types of glacier-models would be to increase their sophistication. For example, it could be interesting to include a parameterization of cloudiness modulating the incoming solar radiation. Eventually, it could be appropriate to move toward energy balance models under certain favorable conditions. But it must be kept in mind that it is quite tough to increase the complexity of the model without adding extra parameters that cannot be constrained accurately for the past.

Acknowledgements. This work greatly benefited from the observation program of tropical glaciers (ORE – GLACIOCLIM program) undertaken by the Institut de Recherches pour le Développement (IRD) in Bolivia.



The publication of this article is financed by CNRS-INSU.

References

- Anslow, F. S., Hostetler, S., Bidlake, W. R., and Clark, P. U.: Distributed energy balance modeling of South Cascade Glacier, Washington and assessment of model uncertainty, *J. Geophys. Res.-Earth*, 113, F02019, doi:10.1029/2007JF000850, 2008.
- Blard, P.-H., Lave, J., Pik, R., Wagnon, P., and Bourles, D.: Persistence of full glacial conditions in the central Pacific until 15,000 years ago, *Nature*, 449, 591–594, 2007.
- Blard, P.-H., Lavé, J., Farley, K. A., Fornari, M., Jiménez, N., and Ramirez, V.: Late local glacial maximum in the Central Altiplano triggered by cold and locally-wet conditions during

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the paleolake Tauca episode (17–15 ka, Heinrich 1), *Quaternary Sci. Rev.*, 28, 3414–3427, 2009.

Braithwaite, R. J.: Positive degree-day factors for ablation on the Greenland ice-sheet studied by energy-balance modeling, *J. Glaciol.*, 41, 153–160, 1995.

5 Braithwaite, R. J. and Olesen, O. B.: Ice ablation in West Greenland in relation to air temperature and global radiation, *Z. Gletscherkd. Glazialgeol.* 20, 155–168, 1985.

Brun, E., Martin, E., Simon, V., Gendre, C., and Coleou, C.: An energy and mass model of snow cover for operational avalanche forecasting, *J. Glaciol.*, 35, 333–342, 1989.

10 Favier, V., Wagnon, P., and Ribstein, P.: Glaciers of the inner and outer tropics : a different behaviour but a common response to climatic forcing, *Geophys. Res. Lett.*, 31, L16403, doi:10.1029/2004GL020654, 2004.

Flowers, G. E., Bjornsson, H., Geirsdottir, A., Miller, G. H., and Clarke, G. K. C.: Glacier fluctuation and inferred climatology of Langjokull ice cap through the Little Ice Age, *Quaternary Sci. Rev.*, 26, 2337–2353, 2007.

15 Forland, E., Allerup, P., Dahlström, B., Elomaa, E. T., Perälä, J., Rissanen, P., Vedin, H., and Vejen, F.: Manual for operational correction of Nordic precipitation data, Det Norske Meteorologiske Institutt, Report No. 24/96, 1996.

Francou, B., Ribstein, P., Saravia, R., and Tiriau, E.: Monthly balance and water discharge of an intertropical glacier – Zongo glacier, Cordillera Real, Bolivia, 16°S, *J. Glaciol.*, 41, 61–67, 1995.

20 Garreaud, R. D., Vuille, M., Compagnucci, R., and Marengo, J.: Present-day South American climate, *Palaeogeogr. Palaeocli.*, 281, 180–195, 2009.

Greuell, W. and Konzelmann, T.: Numerical modeling of the energy-balance and the englacial temperature of the Greenland ice-sheet – calculations for the ETH-camp location (West Greenland, 1155 m asl), *Global Planet. Change*, 9, 91–114, 1994.

25 Hock, R.: A distributed temperature-index ice and snowmelt model including potential direct solar radiation, *J. Glaciol.*, 45, 101–111, 1999.

Hock, R.: Temperature index melt modelling in mountain areas, *J. Hydrol.*, 282, 104–115, 2003.

30 Hock, R.: Glacier melt: a review of processes and their modelling, *Prog. Phys. Geog.*, 29, 362–391, 2005.

Hock, R. and Holmgren, B.: A distributed surface energy-balance model for complex topography and its application to Storglaciaren, Sweden, *J. Glaciol.*, 51, 25–36, ISSN: 0022-1430,

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2005.

Johannesson, T., Sigurdsson, O., Laumann, T., and Kennett, M.: Degree-Day Glacier Mass-Balance Modeling With Applications To Glaciers In Iceland, Norway And Greenland, *J. Glaciol.*, 41, 345–358, 1995.

5 Kageyama, M., Harrison, S. P., and Abe-Ouchi, A.: The depression of tropical snowlines at the last glacial maximum: What can we learn from climate model experiments?, *Quatern. Int.*, 138, 202–219, 2005.

Kaser, G.: Glacier-climate interaction at low latitudes, *J. Glaciol.*, 47, 195–204, 2001.

Kaser, G., Hastenrath, S., and Ames, A.: Mass balance profiles on tropical glaciers, *Z. Gletscherkd. Glazialgeol.*, 32, 75–81, 1996.

10

Kayastha, R. B., Ageta, Y., and Nakawo, M.: Positive degree-day factors for ablation on glaciers in the Nepalese Himalayas: case study on glacier AX010 in Shoron Himal, Nepal. *Bull. Glaciol. Res.*, 17, 1–10, 2000.

Klein, A. G., Seltzer, G. O., and Isacks, B. L.: Modern and last local glacial maximum snowlines in the Central Andes of Peru, Bolivia, and Northern Chile, *Quaternary Sci. Rev.*, 18, 63–84, 1999.

15

Klok, E. J. and Oerlemans, J.: Model study of the spatial distribution of the energy and mass balance of Morteratschgletscher, Switzerland, *J. Glaciol.*, 48, 505–518, 2002.

Lejeune, Y.: Apports des modèles de neige CROCUS et de sol ISBA à l'étude du bilan glaciologique d'un glacier tropical et du bilan hydrologique de son bassin versant, 2009.

20

Lejeune, Y., Wagnon, P., Bouilloud, L., Chevallier, P., Etchevers, P., Martin, E., Sicart, J. E., and Habets, F.: Melting of snow cover in a tropical mountain environment in Bolivia: Processes and modeling, *J. Hydrometeorol.*, 8, 922–937, 2007.

Mark, B. G., Harrison, S. P., Spessa, A., New, M., Evans, D. J. A., and Helmens, K. F.: Tropical snowline changes at the last glacial maximum: A global assessment, *Quatern. Int.*, 138, 168–201, 2005.

25

Molg, T., Hardy, D. R., and Kaser, G.: Solar-radiation-maintained glacier recession on Kilimanjaro drawn from combined ice-radiation geometry modeling, *J. Geophys. Res.-Atmos.*, 108, 4731, doi:10.1029/2003JD003546, 2003.

Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models. Part 1. A discussion of principles, *J. Hydrol.*, 10, 282–290, 1970.

30

Oerlemans, J.: Extracting a climate signal from 169 glacier records, *Science*, 308, 675–677, 2005.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ohmura, A., Kasser, P., and Funk, M.: Climate at the equilibrium line of glaciers, *J. Glaciol.*, **38**, 397–411, 1992.
- Paterson, W. S. B.: *The physics of glaciers*, Pergamon Press, Elsevier Science Ltd, Oxford, 1994.
- 5 Pellicciotti, F., Brock, B., Strasser, U., Burlando, P., Funk, M., and Corripio, J.: An enhanced temperature-index glacier melt model including the shortwave radiation balance: development and testing for Haut Glacier d’Arolla, Switzerland, *J. Glaciol.*, **51**, 573–587, 2005.
- Plummer, M. A. and Phillips, F. M.: A 2-D numerical model of snow/ice energy balance and ice flow for paleoclimatic interpretation of glacial geomorphic features, *Quaternary Sci. Rev.*, **22**, 1389–1406, 2003.
- 10 Ribstein, P., Tiriau, E., Francou, B., and Saravia, R.: Tropical climate and glacier hydrology – A case study in Bolivia, *J. Hydrol.*, **165**, 221–234, 1995.
- Rupper, S. and Roe, G.: Glacier changes and regional climate: a mass and energy balance approach, *J. Climate*, **21**, 5384–5401, 2008.
- 15 Shi, Y.: Characteristics of late Quaternary monsoonal glaciation on the Tibetan Plateau and in East Asia, *Quatern. Int.*, **97–8**, 79–91, 2002.
- Sicart, J. E., Wagnon, P., and Ribstein, P.: Atmospheric controls of the heat balance of Zongo Glacier (16°S, Bolivia), *J. Geophys. Res.-Atmos.*, **110**, D12106, doi:10.1029/2004JD005732, 2005.
- 20 Sicart, J. E., Ribstein, P., Francou, B., Pouyaud, B., and Condom, T.: Glacier mass balance of tropical Zongo glacier, Bolivia, comparing hydrological and glaciological methods, *Global Planet. Change*, **59**, 27–36, 2007.
- Sicart, J. E., Hock, R., and Six, D.: Glacier melt, air temperature, and energy balance in different climates: The Bolivian Tropics, the French Alps, and northern Sweden, *J. Geophys. Res.-Atmos.*, **113**, D24113, doi:10.1029/2008JD010406, 2008.
- 25 Singh, P. and Kumar, N.: Determination of snowmelt factor in the Himalayan region, *Hydrol. Sci. J.*, **41**, 301–310, 1996.
- Singh, P., Kumar, N., Ramasastri, K. S., and Singh, Y.: Influence of a fine debris layer on the melting of snow and ice on a Himalayan glacier In *Debris-covered Glaciers*, Proceedings of the Workshop on Debris-covered Glaciers, edited by: Nakawo, M., Raymond, C. F., and Fountain, A., 63–69, IAHS, Seattle, 2000.
- 30 Soruco, A., Vincent, C., Francou, B., Ribstein, P., Berger, T., Sicart, J.-E., Wagnon, P., Arnaud, Y., Favier, V., and Lejeune, Y.: Mass balance of Glaciar Zongo, Bolivia, between 1956 and

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- 2006, using glaciological, hydrological and geodetic methods, *Ann. Glaciol.*, 50, 1–8, 2009.
- Vincent, C.: Influence of climate change over the 20th Century on four French glacier mass balances, *J. Geophys. Res.-Atmos.*, 107, 4375, 10.1029/2001JD000832, 2002.
- Wagnon, P., Ribstein, P., Kaser, G., and Berton, P.: Energy balance and runoff seasonality of a Bolivian glacier, *Global Planet. Change*, 22, 49–58, 1999.
- Wagnon, P., Sicart, J.-E., Berthier, E., and Chazarin, J.-P.: Wintertime high-altitude surface energy balance of a Bolivian glacier, Illimani, 6340 m above sea level, *J. Geophys. Res.-Atmos.*, 108, 4177, 10.1029/2002JD002088, 2003.
- Wagnon, P., Lafaysse, M., Lejeune, Y., Maisinsho, L., Rojas, M., and Chazarin, J. P.: Understanding and modelling the physical processes that govern the melting of the snow cover in a tropical mountain environment in Ecuador, *J. Geophys. Res.*, 114, D19113, doi:10.1029/2009JD012292, 2009.
- WMO: Intercomparison of models for snowmelt runoff, Operational Hydrology Report WMO No. 646, 1986.

Table 1. Meteorological data near the Zongo glacier 1 for the 1997–2006 hydrological 2 yr. Hydrological years range from 1 September to 31 August.

| Hydrological year | Weather station | Month | Annual T ($^{\circ}\text{C}$) / Precipitation (mm) | | | | | | | | | | | | |
|--|-----------------|------------------------------|--|-----|-----|-----|------|------|------|------|------|-------|-------|-------|------|
| | | | Sep | Oct | Nov | Dec | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | |
| 1997–1998 | MEVIS – 4750 m | T_m ($^{\circ}\text{C}$) | 0.8 | 3.2 | 3.0 | 4.0 | 3.8 | 4.1 | 3.9 | 3.9 | 3.2 | 1.7 | 1.9 | 1.9 | 3.0 |
| | | 1σ | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 2.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.4 | 2.3 | |
| | | P (mm) | 120 | 90 | 84 | 102 | 144 | 180 | 168 | 48 | 0 | 72 | 0 | 36 | 1044 |
| 1998–1999 | MEVIS – 4750 m | T_m ($^{\circ}\text{C}$) | 2.1 | 2.1 | 2.3 | 3.0 | 1.4 | 1.7 | 1.6 | 1.4 | 1.6 | 1.3 | 0.2 | 0.8 | 1.6 |
| | | 1σ | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 2.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.4 | 2.3 | |
| | | P (mm) | 24 | 72 | 180 | 108 | 192 | 216 | 252 | 86 | 0 | 0 | 24 | 12 | 1166 |
| 1999–2000 | MEVIS – 4750 m | T_m ($^{\circ}\text{C}$) | 1.0 | 1.0 | 1.6 | 2.4 | 1.5 | 1.7 | 2.2 | 2.1 | 2.5 | 0.7 | -0.3 | 0.5 | 1.4 |
| | | 1σ | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 2.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.4 | 2.3 | |
| | | P (mm) | 108 | 84 | 72 | 156 | 300 | 264 | 144 | 36 | 24 | 36 | 0 | 72 | 1296 |
| 2000–2001 | MEVIS – 4750 m | T_m ($^{\circ}\text{C}$) | 2.0 | 1.0 | 2.8 | 2.1 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.5 |
| | | 1σ | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 2.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.4 | 2.3 | |
| | | P (mm) | 30 | 168 | 54 | 360 | 234 | 246 | 240 | 48 | 36 | 36 | 0 | 77 | 1529 |
| 2001–2002 | MEVIS – 4750 m | T_m ($^{\circ}\text{C}$) | 1.2 | 2.1 | 3.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.1 | 2.1 | 1.0 | -0.9 | 0.4 | 1.7 |
| | | 1σ | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 2.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.4 | 2.3 | |
| | | P (mm) | 12 | 60 | 126 | 66 | 204 | 168 | 329 | 19 | 96 | 17 | 60 | 46 | 1202 |
| 2002–2003 | MEVIS – 4750 m | T mens | 1.2 | 1.4 | 2.4 | 2.6 | 2.7 | 2.1 | 2.1 | 2.6 | 1.8 | 1.7 | 0.9 | 0.6 | 1.8 |
| | | $1s$ | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 2.2 | 2.0 | 2.2 | 2.3 | 2.3 | 2.4 | 2.3 | |
| | | P | 96 | 120 | 96 | 204 | 300 | 156 | 240 | 36 | 24 | 0 | 60 | 24 | 1356 |
| 2003–2004 | ORE – 5050 m | T_m ($^{\circ}\text{C}$) | 0.1 | 0.5 | 1.9 | 0.4 | 0.5 | 1.5 | 1.4 | 0.7 | 0.1 | -1.5 | -1.1 | -1.9 | 0.2 |
| | | 1σ | 2.8 | 2.4 | 2.1 | 1.5 | 1.6 | 2.3 | 2.1 | 2.2 | 2.3 | 2.0 | 2.3 | 2.1 | |
| | | P (mm) | 90 | 106 | 46 | 297 | 233 | 160 | 142 | 52 | 14 | 14 | 42 | 47 | 1243 |
| 2004–2005 | ORE – 5050 m | T_m ($^{\circ}\text{C}$) | -0.6 | 0.3 | 1.1 | 1.4 | 0.90 | 1.34 | 1.77 | 1.49 | 1.30 | -0.04 | 0.12 | -0.11 | 0.7 |
| | | 1σ | 2.1 | 2.2 | 2.1 | 2.0 | 1.9 | 2.1 | 2.1 | 2.1 | 2.3 | 2.5 | 2.5 | 2.5 | |
| | | P (mm) | 83 | 83 | 106 | 121 | 325 | 302 | 53 | 54 | 2 | 20 | 23 | 15 | 1187 |
| 2005–2006 | ORE – 5050 m | T_m ($^{\circ}\text{C}$) | -1.4 | 0.6 | 1.2 | 0.2 | 0.31 | 1.94 | 0.57 | 0.85 | 0.42 | -0.15 | -0.01 | -0.59 | 0.3 |
| | | 1σ | 2.3 | 1.9 | 2.1 | 1.7 | 1.5 | 2.3 | 1.7 | 2.4 | 2.4 | 2.3 | 2.5 | 2.3 | |
| | | P (mm) | 92 | 302 | 118 | 340 | 206 | 127 | 137 | 15 | 7 | 3 | 0 | 5 | 1352 |
| Monthly lapse rate ^a ($^{\circ}\text{C km}^{-1}$) | | | 6.7 | 6.3 | 6.3 | 6.0 | 5.5 | 5.2 | 5.3 | 5.8 | 6.5 | 6.8 | 6.7 | 6.8 | 6.2 |

^a Determined from 10 weather station ranging between 1900 and 5050 m in the Bolivian Andes (source: www.worldclimate.com).

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Mass balance data (m yr^{-1}) measured on the Zongo 1 glacier for the nine hydrological years between 1997 and 2006. Each hydrological year starts on 1 September and ends on 31 August. Data were collected by the ORE Glacioclim IRD program.

| Altitude (m) | 1997–1998 | 1998–1999 | 1999–2000 | 2000–2001 | 2001–2002 |
|--------------|---------------|-----------|-----------|-----------|-----------|
| 4950 | −8.74 | −6.00 | −5.00 | −3.80 | −5.40 |
| 5050 | −5.82 | −3.00 | −1.50 | −0.99 | −2.60 |
| 5150 | −4.26 | −1.32 | −0.10 | 0.15 | −0.80 |
| 5250 | −3.10 | −0.76 | 0.20 | 0.66 | 0.10 |
| 5350 | −1.93 | −0.19 | 0.45 | 1.01 | 0.40 |
| 5450 | −0.77 | 0.37 | 0.74 | 1.35 | 0.70 |
| 5550 | 0.39 | 0.93 | 0.98 | 1.69 | 0.82 |
| 5650 | 0.69 | 0.96 | 1.07 | 1.57 | 0.93 |
| 5750 | 0.69 | 0.99 | 1.15 | 1.37 | 0.93 |
| 5850 | 0.69 | 0.99 | 1.15 | 1.37 | 0.93 |
| 5950 | 0.69 | 0.99 | 1.15 | 1.37 | 0.93 |
| 6050 | 0.69 | 0.99 | 1.15 | 1.37 | 0.93 |
| Altitude (m) | 2002–2003* | 2003–2004 | 2004–2005 | 2005–2006 | |
| 4950 | (5050 m) −5.0 | −4.00 | −8.00 | −7.20 | |
| 5050 | (5070 m) −3.0 | −2.50 | −5.92 | −3.60 | |
| 5150 | (5110 m) −1.0 | −1.69 | −3.80 | −0.54 | |
| 5250 | (5180 m) 0.3 | −1.08 | −2.10 | 0.38 | |
| 5350 | (5350 m) 0.8 | −0.47 | −1.05 | 0.40 | |
| 5450 | (5450 m) 0.91 | 0.14 | −0.35 | 0.41 | |
| 5550 | 0.91 | 0.75 | 0.35 | 0.42 | |
| 5650 | 0.91 | 0.76 | 0.41 | 0.49 | |
| 5750 | 0.91 | 0.80 | 0.46 | 0.57 | |
| 5850 | | 0.80 | 0.52 | 0.64 | |
| 5950 | | 0.80 | 0.52 | 0.64 | |
| 6050 | | 0.80 | 0.52 | 0.64 | |

* The mass balance data measured in 2002–2003 are characterized by atypic elevations.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Models description, input data and 1 calibrated parameters.

| Model | Formulation of ablation | Input data | Parameters |
|-------|--|---|---|
| 1 | $A = MF \times T_p$ | Monthly temperature – average and standard deviation Monthly precipitation | MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) |
| 2 | $A = MF_{\text{snow/ice}} \times T_p$ | Monthly temperature – average and standard deviation Monthly precipitation | MF_{snow} ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) MF_{ice} ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) |
| 3 | $A_{\text{snow}} = MF \times T_p$ $A_{\text{ice}} = (MF + a \times I) \times T_p$ | Monthly temperature – average and standard deviation Monthly precipitation Clear-sky shortwave radiations | MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) a ($\text{mm } ^\circ\text{C}^{-1} \text{W}^{-1} \text{m}^2 \text{d}^{-1}$) |
| 4 | $A = MF \times T_p + a \times I$ | Monthly temperature – average and standard deviation Monthly precipitation Clear-sky shortwave radiations | MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) a ($\text{mm } \text{W}^{-1} \text{m}^2 \text{d}^{-1}$) |

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 4. Calibrated parameters. R^2 (all years) are the correlation coefficient calculated using the whole set of modeled and observed mass balances of all hydrological years. R_{pos}^2 and R_{neg}^2 are calculated considering the positive and the negative mass balance, respectively.

| Model 1 | | | Model 2 | | | | |
|------------------------------|--|--|-------------------------------------|---|---|---|------------|
| $A = MF \times T$ | | | $A = MF_{\text{ice/snow}} \times T$ | | | | |
| Year | MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) | Best R^2 | Year | MFsnow ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) | MFice ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) | Best R^2 | |
| 1997–1998 | 11.6 | 0.99 | 1997–1998 | 10.4 | 11.7 | 0.99 | |
| 1998–1999 | 15.5 | 0.98 | 1998–1999 | 7.5 | 19.0 | 0.99 | |
| 1999–2000 | 11.8 | 0.94 | 1999–2000 | 3.8 | 22.4 | 0.98 | |
| 2000–2001 | 10.3 | 0.92 | 2000–2001 | 2.2 | 20.0 | 0.98 | |
| 2001–2002 | 12.3 | 0.95 | 2001–2002 | 4.0 | 19.8 | 0.99 | |
| 2002–2003 | 12.6 | 0.77 | 2002–2003 | 6.2 | 20.1 | 0.83 | |
| 2003–2004 | 9.0 | 0.96 | 2003–2004 | 23.0 | 7.6 | 0.98 | |
| 2004–2005 | 13.0 | 0.98 | 2004–2005 | 11.8 | 13.2 | 0.99 | |
| 2005–2006 | 11.9 | 0.84 | 2005–2006 | 1.1 | 25.7 | 0.91 | |
| All years | 11.9 ± 1.3 | 0.92 | All years | 8.7 ± 0.6 | 12.7 ± 1.4 | 0.93 | |
| | R2pos | 0.48 | | | R2pos | 0.49 | |
| | R2neg | 0.87 | | | R2neg | 0.87 | |
| Model 3 | | | | Model 4 | | | |
| If snow: $A = MF \times T_p$ | | If ice: $A = (MF + a \times l) \times T_p$ | | $A = MF + a \times l$ | | | |
| Year | MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) | $a \times 10^{-2}$ ($\text{mm } ^\circ\text{C}^{-1} \text{W}^{-1} \text{m}^2 \text{d}^{-1}$) | Best R^2 | Year | MF ($\text{mm } ^\circ\text{C}^{-1} \text{d}^{-1}$) | a ($10^{-4} \text{mm } \text{W}^{-1} \text{m}^2 \text{d}^{-1}$) | Best R^2 |
| 1997–1998 | 10.9 | 0.18 | 0.99 | 1997–1998 | 11.5 | 1.4 | 0.99 |
| 1998–1999 | 0.8 | 12.6 | 0.89 | 1998–1999 | 15.4 | 1.1 | 0.98 |
| 1999–2000 | 4 | 4.4 | 0.82 | 1999–2000 | 11.8 | 0 | 0.94 |
| 2000–2001 | 1 | 9.4 | 0.86 | 2000–2001 | 10.2 | 0 | 0.92 |
| 2001–2002 | 3.1 | 4.8 | 0.84 | 2001–2002 | 12.3 | 0 | 0.95 |
| 2002–2003 | 6.6 | 2.8 | 0.93 | 2002–2003 | 12.6 | 0 | 0.77 |
| 2003–2004 | 9 | 0 | 0.96 | 2003–2004 | 8 | 20 | 0.99 |
| 2004–2005 | 12 | 0.28 | 0.98 | 2004–2005 | 12.4 | 15 | 0.99 |
| 2005–2006 | 7.7 | 1.31 | 0.84 | 2005–2006 | 11.6 | 4 | 0.84 |
| All years | 8.8 ± 1.0 | 0.92 ± 0.21 | 0.93 | All years | 11.8 ± 1.3 | 2.1 ± 0.5 | 0.93 |
| | | R2pos | 0.48 | | | R2pos | 0.50 |
| | | R2neg | 0.87 | | | R2neg | 0.87 |

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.




Fig. 1. Map of the Zongo glacier. Locations of the weather stations are shown, as well as those of the snow pits and the stakes used to measure the mass balance during the 2005–2006 year.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏴

⏵

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Values of the efficiency criterion R^2 vs. ablation parameters (MF, a) for models 2 to 4 inferred for the 9 hydrological years between 1997 and 2006. Red star shows the best parameters, which yields the maximum R^2 value.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.




Fig. 3. Plot of data and best-fit modeled mass balance for the hydrological years 1997–1998, 2003–2004 and 2004–2005. Dots are observations, the red line are the simulated mass balances (MB in mm w.e. yr⁻¹ vs. altitude in m). Blue line is the annual ablation; black line is the annual accumulation.

Title Page

| | |
|--------------------------|--------------|
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Printer-friendly Version | |
| Interactive Discussion | |



Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. For each of the 4 models, plot of the observed vs. best-fit modeled annual mass balances for the 9 hydrological years between 1997 and 2006.

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.




Fig. 5. (a) Mean annual positive temperature (T_p , in $^{\circ}\text{C}$, calculated from the positive degree day approach) and mean annual clear sky incoming short-wave radiations on the Zongo glacier surface (I , in kW m^{-2} , calculated taking into account the glacier slope and orientation) vs. elevation (m). **(b)** For Model 4, shown are the respective proportions of total ablation arising from the $(a \times I)$ and the $(MF \times T_p)$ terms vs. elevation (m). The MF and a values used in the calculation are $11.8 \text{ mm } ^{\circ}\text{C}^{-1} \text{ d}^{-1}$ and $2.1 \times 10^{-4} \text{ mm W}^{-1} \text{ m}^2 \text{ } ^{\circ}\text{C}^{-1} \text{ d}^{-1}$. This calculation shows that the ablation arising from the $a \times I$ term becomes significant only for elevations higher than 5600 m.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Degree-day melt models for paleoclimate reconstruction

P.-H. Blard et al.




Fig. 6. Precipitation – Temperature curves able to maintain the ELA at 5400 m for models 1 to 4. Input temperature and precipitation are the mean of the 9 hydrological years. Used parameters are those calibrated from the present study (Table 4). Envelope shows the 1σ error calculated by taking into account the parameters uncertainty given in Table 4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion