



Pleistocene glacial history of the New Zealand subantarctic islands

Rainsley, Eleanor^{1*}, Turney, Chris S.M.^{2,3}, Golledge, Nicholas R.^{4,5}, Wilmshurst, Janet M.^{6,7}, McGlone, Matt S.⁶, Hogg, Alan G.^{8,9}, Li, Bo^{10,11}, Thomas, Zoë A.^{2,3}, Roberts, Richard^{10,11}, Jones, Richard T.^{12, †}, Palmer, Jonathan G.^{2,3}, Flett, Verity¹³ de Wet, Gregory¹⁴, Hutchinson, David K.¹⁵, Lipson, Mathew J.²,
5 Fenwick, Pavla¹⁶, Hines, Ben R.¹⁷, Binetti, Umberto¹⁸ and Fogwill, Christopher J.^{1,2}

1. ICELAB, School of Geography, Geology and the Environment, University of Keele, Staffordshire, ST5 5BG
2. Climate Change Research Centre, School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, Australia
3. Australian Research Council (ARC) Centre of Excellence for Australian Biodiversity and Heritage, University of New
10 South Wales, NSW 2052, Australia
4. Antarctic Research Centre, Victoria University of Wellington, Wellington 6140, New Zealand
5. GNS Science, Avalon, Lower Hutt 5011, New Zealand
6. Long Term Ecology Laboratory, Landcare Research, Lincoln, New Zealand
7. School of Environment, University of Auckland, Private Bag 92019, Auckland 1142, New Zealand
- 15 8. Waikato Radiocarbon Laboratory, University of Waikato, Private Bag 3105, Hamilton, New Zealand
9. Australian Research Council (ARC) Centre of Excellence for Australian Biodiversity and Heritage, University of Waikato, Hamilton 3240, New Zealand
10. Centre for Archaeological Science, School of Earth and Environmental Sciences, University of Wollongong, Australia
11. Australian Research Council (ARC) Centre of Excellence for Australian Biodiversity and Heritage, University of
20 Wollongong, NSW 2522, Australia
12. Department of Geography, University of Exeter, Exeter EX4 4RJ, UK
13. School of the Environment, University of Dundee, Nethergate DD1 4HN, UK.
14. Institute of Arctic and Alpine Research, University of Colorado, Boulder, Boulder, CO, USA
15. Bolin Centre for Climate Research and Department of Geological Sciences, Stockholm University, Stockholm, Sweden
- 25 16. Gondwana Tree-Ring Laboratory, PO Box 14, Little River, Canterbury 7546, New Zealand.
17. School of Geography, Environment and Earth Sciences, Victoria University of Wellington, Wellington, New Zealand
18. Centre for Ocean and Atmospheric Studies, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK.

† Deceased

30 *Correspondence to:* Eleanor Rainsley (eleanor.rainsley@gmail.com)



Abstract. The New Zealand subantarctic islands of Auckland and Campbell, situated between the Subtropical Front and the Antarctic Convergence in the Pacific sector of the Southern Ocean, provide valuable terrestrial records from a globally-important climatic region. Whilst the islands show clear evidence of past glaciation, the timing and mechanisms behind Pleistocene environmental and climate changes remain uncertain. Here we present a multidisciplinary study of the islands – including marine and terrestrial geomorphological surveys, extensive analyses of sedimentary sequences, a comprehensive dating program, and glacier flowline modelling – to investigate multiple phases of glaciation across the islands. We find evidence that the Auckland Islands hosted a small ice cap at $384,000 \pm 26,000$ years ago (384 ± 26 ka), most likely during Marine Isotope Stage 10, a period when the Subtropical Front was pushed northwards by seven degrees, and consistent with hemispheric-wide glacial expansion. Despite previous interpretations that suggest the maximum glacial extent occurred in the form of valley glaciation at the Last Glacial Maximum (LGM; ~ 21 ka) age, our combined approach suggests minimal LGM glaciation across the New Zealand Subantarctic Islands, and that no glaciers were present during the Antarctic Cold Reversal (ACR; ~ 15 - 13 ka). Instead, our flowline modelling, constrained by field evidence, implies that despite a regional mean annual air temperature depression of $\sim 5^\circ\text{C}$ during the LGM, a combination of high seasonality and low precipitation left the islands incapable of sustaining significant glaciation. We suggest that northwards expansion of winter sea ice during the LGM and subsequent ACR led to precipitation starvation across the mid to high latitudes of the Southern Ocean, resulting in restricted glaciation of the subantarctic islands.



1 Introduction

The Southern Ocean plays a critical role in the climate system, connecting the three main ocean basins (i.e. Atlantic, Pacific and Indian Oceans) and modulating regional to global atmospheric temperatures and weather patterns on historic to millennial timeframes (Anderson et al., 2009; Jones et al., 2016; Marshall and Speer, 2012; Steig et al., 2009; Turney et al., 2016a; WAIS, 2015). During the late Pleistocene (110,000 to 11,650 years ago, 110-11.65 ka), marine and terrestrial records across the mid-latitudes demonstrate broadly similar trends to those reported from Antarctic ice core sequences (EPICA, 2006; Kaplan et al., 2008; McGlone et al., 2010; Moreno et al., 2009; Pahnke et al., 2003; Pedro et al., 2015). Following the Last Glacial Maximum (LGM, 21 ± 3 ka), these records show climatic amelioration from ~ 17 ka through to the onset of the Holocene (Walker et al., 2009) interrupted by a two-thousand year cooling event described as the Antarctic Cold Reversal (ACR), approximately 14.7 to 12.7 ka (Hogg et al., 2016; Pedro et al., 2015).

In the south Pacific, greatest cooling was focussed on ~ 28 -20 ka (the so-called early- or eLGM) (Denton et al., 2010; Fogwill et al., 2015; Hein et al., 2010; Newnham et al., 2007), with sea-surface temperatures (SSTs) apparently between 6-10 °C lower than present day (Barrows et al., 2007; Hayward et al., 2008; Pahnke and Sachs, 2006; Panitz et al., 2015), and terrestrial proxies suggesting comparable air temperature reductions of the order of 5-6 °C over southern New Zealand (Golledge et al., 2012; Newnham et al., 1989). The prevailing winds and precipitation belts appear to have varied both spatially and temporally with these changes throughout the late Pleistocene (Fogwill et al., 2015). Whilst there is abundant evidence of widespread glaciation across mainland New Zealand during these periods (Porter, 1975; Williams et al., 2015), there are relatively few high-resolution records available from the Southern Ocean (McGlone et al., 2010). The subantarctic islands that span the Southern Ocean have the potential to provide valuable palaeo-climatic and -environmental records. To gain a fuller understanding of climate dynamics over the region it is therefore vital we exploit these currently under-utilised archives fully.

1.1 The New Zealand Subantarctic Islands



5 Located in the Pacific Sector of the Southern Ocean, the New Zealand subantarctic Auckland (50.70°S, 166.10°E) and Campbell (52.54°S, 169.14°E) Islands (Fig. 1) are the eroded remnants of extensive Oligocene to Miocene basaltic volcanism (Quilty, 2007), with rugged topography, reaching up to 569 m above sea level (a.s.l) (Campbell Island) and 664 m a.s.l. (Auckland Island). At present the islands are
10 blanketed with peat deposits (up to 10 m thick) to at least 400-500 m a.s.l. The islands lie in the southwest Pacific in the core of Southern Hemisphere westerly winds and experience the same climate regime with relatively mild mean annual temperatures. Situated today between the Subtropical Convergence (approximately 45° S) and the cool Antarctic waters of the Antarctic Convergence (around 55° S), these ocean fronts may have shifted by up to 5° further northwards during the LGM (Nelson et al., 1993) leading
15 to a mean annual air temperature depression over the islands of ~6°C (McGlone, 2002). Campbell and Auckland Islands are therefore highly sensitive to regional climate change during the late Pleistocene, which is potentially archived in their extensive peat deposits and preserved landscape features (McGlone, 2002; Turney et al., 2016b).

15 The extent to which the subantarctic islands in the southwest Pacific were glaciated during the LGM remains highly uncertain, leaving many questions regarding the late Pleistocene climate in the region unanswered. Previous research carried out on the Auckland Island archipelago reported exposed sediment sections on Enderby Island (Fig. 1) with two distinct layers of glacial till, separated by laminated silt deposited in a pro-glacial lake environment (Fleming et al., 1976). The so-called ‘Enderby Formation’
20 was interpreted as having been deposited by the advance of a glacier in the U-shaped valley of Port Ross, and it was therefore assumed that the numerous fjord and cirque features along the east and south coasts of the main Auckland Island also sustained significant glaciation (Fleming et al., 1976; Hodgson et al., 2014a; McGlone, 2002). The presence of pollen and spores in the silt between the two till layers on
25 Enderby Island (Fleming et al., 1976) hints towards an extended period between two glacial events, but they could equally represent minor glacier fluctuations of the same event. However, little data have been produced constraining the extent and timing of glaciation on the Auckland Islands; in the absence of a dated chronology these features have been assumed to be evidence of an extensive valley glaciation occurring during the LGM (Fleming et al., 1976; Hodgson et al., 2014a; McGlone, 2002), although it is



also acknowledged that the features are undated and may be of older origins (Hodgson et al., 2014b). Campbell Island, c. 300 km further south and with a comparable elevation, exhibits a similar geomorphology to that of the Auckland Islands, with wide U-shaped valleys and cirque-like features (Oliver et al., 1950). Possible glacial depositional features such as till (overlain by peat with basal dates of ~13,000 yr BP), erratic boulders, and kame terraces have also been reported, posited to be of LGM origin (Hodgson et al., 2014a; McGlone et al., 1997); Campbell (1981), however, suggested some of them may have been formed by fluvial processes. Furthermore, palaeoecological reconstructions suggest that endemic flora and fauna on the islands prevailed throughout the late Pleistocene (McGlone et al., 2010; Mitchell et al., 2014; Van der Putten et al., 2010), conflicting with the paradigm of extensive LGM glaciation (Hodgson et al., 2014a; McGlone, 2002) and raising fundamental questions about our understanding of LGM climatic conditions across this sector of the Southern Ocean.

Here we report the results of a multidisciplinary study of the New Zealand subantarctic Campbell and Auckland Islands that aims to constrain the extent and timing of glacial activity during the late Pleistocene and provide insights into past shifts in climate regimes across the Southern Ocean.

2 Methods

To understand the glacial history of the islands we undertook a multidisciplinary field approach during the Australasian Antarctic Expedition 2013-14 (AAE) and subsequent fieldwork in November 2014, building on previous work led by MM and JW (McGlone et al., 1997; McGlone et al., 2000). This research programme encompassed geomorphological and aerial surveys, an extensive radiocarbon (^{14}C) and optical dating program, and the analysis of multiple marine and terrestrial sediment sequences, allowing us to constrain the outputs of targeted glacier flowline modelling. We summarise the methodology below; further technical details can be found in the extended methodology in Appendix 1.

25

2.1 Geomorphological survey

Initial surveying was carried out using topographic maps and satellite imagery to identify possible sites of former glaciation, which were then ground-truthed during fieldwork. Additional sites showing



potential glacial features were identified from Fleming et al. (1976), as well as his field diaries (Hince, 2007). Relevant sites were targeted with an Unmanned Aerial Vehicle (UAV) to produce high-resolution digital elevation models (DEMs.) To investigate the potential presence of relict glacial geomorphology including moraine features on the inner shelf areas of the New Zealand subantarctic islands, we undertook a ship-based multibeam survey within Perseverance Harbour, Campbell Island (Fig. 1), a valley with an over-deepened U-shaped profile typical of formerly glaciated regions.

2.2 Sedimentary analysis

Lacustrine and peat sediments were sampled in two key locations; from outside the relict glacial limits above Musgrave Harbour, Auckland Island (Core 10, Fig. 1, Figs. S1 and S2, Table 1), and from inside and outside the moraine limits of a distinct glacial cirque on Auckland Island. Named here as Lower 'Lake' Speight (although not itself hosting a lake), the cirque is situated above Coleridge Bay in Carnley Harbour, Auckland Island, immediately south of glacial Lake Speight. The cores are identified as Lower 'Lake' Speight (LLS) Cirque and LLS Terrace cores (Cores 14 and 15, respectively; Figs. 1 and 2, Table 1). In addition, we report here a new exposed sedimentary sequence on a north-facing cliff near Pillar Rock, Auckland Island (50.518°S, 166.217°E; 30 m a.s.l., Figs. 1 and 3) from which the organic units were sampled with monolith tins. Key sedimentary sequences were analysed for organic content by loss-on-ignition (LOI). The LLS Cirque core was subsampled (sample size 0.5 cm) every 2 cm from 64-104 cm and every 1 cm thereafter. The LLS Terrace core was subsampled (sample size 0.5 cm) every 5 cm from 0-95 cm, and every 2.5 cm thereafter. In addition, two shallow marine sediment cores were taken from Carnley Harbour (50.82°S, 166.01°E) (Cores 11 and 12, Fig. 1, Table 1). Over 1 m of sediment was recovered from each deployment of the corer, with the cores consisting mostly of peats overlain by fine silts containing occasional marine mollusk shells in life position. Samples were taken from the face of the Enderby Formation laminated silt at intervals of approximately 10 cm (Fig. 4). These were prepared for particle size analysis using a Saturn Digitiser and the results categorised using the Udden-Wentworth Scale (McCave and Syvitski, 1991).



2.3 Chronology

2.3.1 Radiocarbon dating

To provide a robust geochronological framework we also undertook an extensive radiocarbon dating programme of new material combined with previously collected sequences (Fig. 1, Tables 1 and 2) to complement previously reported ages (Moar, 1958; McGlone, 2002). The latter sites include the peat sequences from the Auckland Islands at Ranui Cove in Port Ross (Core 8) and two closely adjacent profiles (Deas Head Forest and Deas Head Bog, Core 4) from peats on McCormacks Peninsula, Port Ross. Five other unpublished sites collected by MM and JW in the Port Ross area in 1994 have new basal ages reported here: a bog from Mt Hooker at shrubline (Core 9); blanket peats from the cliffs of North Point Bay, northern Auckland Island (Core 2); a peat sequence adjacent to Pillar Rock, northern McCormacks Peninsula (Core 3); and two cores from a raised bog complex “O”, in the interior of McCormacks Peninsula (Cores 6 and 7). A previously published basal ^{14}C date from Deas Head (NZA4509; $15,170 \pm 140$ ^{14}C yr BP, $\delta^{13}\text{C}$ -34.1‰) (McGlone, 2002) was obtained from a sandy ‘peat’, but with $\sim 1\%$ carbon content using this age for the onset of the peat growth is considered highly uncertain (Lowe and Walker, 2000). A new core was therefore taken from Deas Head, and the basal age reported here (Core 4). All samples carbon dated in this study have a carbon content $>10\%$. On Campbell Island, the west-facing Hooker Valley (Core 16) peat and silt cliff section was published in McGlone *et al.* (1997) and the upland Mt Honey Saddle raised bog (Core 21) and Homestead Scarp blanket (Core 18) peat sites were published in McGlone *et al.* (2010). Several other Campbell Island sites have been dated but only the last 2000 years of the sequences published (McGlone *et al.*, 2007). Here we report new basal ages from two of these Campbell Island sites: the lowland raised bog at Homestead Ridge (Core 19) and the upland, west-facing Col Ridge blanket peat (Core 17).

2.3.2 Optical dating

With no organic material present, to provide an age constraint for the Enderby Formation (Figure 1), we collected four samples (Enderby-1, -2, -3 and -4) for optical dating from the sediments overlying the till, hammering opaque plastic tubes (5 cm in diameter) into the cleaned section faces. The tubes were removed and wrapped in lightproof plastic for transport to the Luminescence Dating Laboratory at the



University of Wollongong. Given the potential for the samples to lie beyond the dating range of quartz optically stimulated luminescence, we instead used infrared stimulated luminescence (IRSL), measuring equivalent dose (D_e) values for individual potassium-rich feldspar (K-feldspar) grains using a pIRIR (post-infrared IRSL) procedure, made possible by recent developments in the field (Buylaert et al., 2009; Li et al., 2014; Li and Li, 2011; Thiel et al., 2011; Thomsen et al., 2008). Details of sample preparation methods are provided in Supplementary Information, together with D_e and dose rate measurement procedures and supporting data. The optical dating results are summarised in Table 3. The pIRIR ages for the four samples are 400 ± 64 , 343 ± 43 , 284 ± 101 , 413 ± 39 ka for Enderby-1, -2, -3 and -4, respectively. Enderby-1 and -3 have large age uncertainties due mainly to the small number of grains (< 10 grains) suitable for D_e determination. See Appendix 1 and Supplementary Figures 3-6 for more details.

2.4 Flowline modelling

To investigate the impact of changing climatic conditions on glaciation across the southwest Pacific subantarctic islands during the late Pleistocene, we used a glacier flowline model ensemble approach. Due to the complex topography of the Auckland Islands we focussed our modelling on the west-east aligned Perseverance Harbour (Campbell Island), a U-shaped valley likely to have been an independent ice catchment and ideally suited for flowline modelling (Fig. 1D).

The glacier model employed is essentially the one-dimensional shallow-ice approximation (SIA) flowline model of Golledge & Levy (2011), with two modifications implemented for this study. Firstly, the positive degree-day (PDD) scheme formerly used to calculate surface mass balance (SMB) has been replaced by a simplified energy-balance scheme, following the insolation-temperature melt method (Eq. 16 of Robinson et al., (2010)). The second modification to the model is the way in which basal sliding velocities are calculated. Previously we used a triangular averaging scheme (Kamb and Echelmeyer, 1986) to smooth calculated driving stress values, which were then used to calculate sliding velocities using a rate factor and sliding exponent (Eq. 8 of Golledge and Levy (2011)). Here we follow instead the approach of Bueler & Brown (2009), and superpose velocity solutions from the SIA with those from the shallow-shelf approximation (SSA).



Input data necessary for our modelling experiments are bed topography (taken from topographic maps) and present-day air temperature and precipitation data (obtained from NIWA). Time-series data for the study period (125 ka to present) are used for summer and winter insolation values and for air temperature and eustatic sea level perturbations from present. For air temperature anomalies we use two forcing patterns, both scaled to a range of prescribed minima at the LGM. Given the parallel changes across the Southern Ocean region and the Antarctic (Fogwill et al., 2015), we use the $\delta^{18}\text{O}$ record from EPICA Dome C (EDC) (Parrenin et al., 2007) as a proxy for Southern Hemisphere atmospheric temperature changes. In addition, we used a proximal sea-surface temperature (SST) record from core DSDP 594 (Hayward et al., 2008), located to the east of New Zealand's South Island (43.632°S 179.379°E, Fig. 1). Starting at 125 ka we initialize our model with present-day topographic and climatological conditions, and simulate glacial advance and retreat according to the calculated SMB and glaciological parameterisation described above.

Since there are few empirical constraints on glacier extent in Perseverance Harbour, our modelling employs full factorial sampling so that results from the mutual combinations of a range of model and climatological parameters can be evaluated with respect to the sparse data, without a priori assumptions about glacier geometry or palaeoclimate. Our parameter-space spans mean annual air temperature (MAAT) scalings of 3, 4, 5, 6, 7, and 8 °C, annual temperature ranges of 5, 8 and 11 °C, and atmospheric transmissivity values of 0.245, 0.25, 0.255, and 0.26 (based on winter values of Robinson *et al.*, (2010) to account for greater year-round cloudiness at Campbell Island than over the Greenland Ice Sheet). We perform each of these parameter combinations both with and without an air temperature / precipitation coupling based on the Clausius-Clapeyron relationship. This relationship parameterizes the processes that lead to a reduction in moisture-holding capacity as air cools, and equates to a change in precipitation of approximately 7% per degree C air temperature change. Finally, each of the parameter combinations is used in runs forced by each of the EDC and SST timeseries data. The ensemble therefore results in 288 unique simulations.



3 Results

3.1 Geomorphology

Surveys using satellite imagery, topographic maps and ground-truthing highlight glacial features on both islands, despite the thick peat cover that subdues the glacial geomorphology across much of the islands.

5

The glacial till described by Fleming *et al.* (1976) on Enderby Island appears to also be represented in similar extensive drift deposits up to 6 m thick, which we identified at Pillar Rock on the north coast of Auckland Island and along the shoreline at Emergency Bay within Carnley Harbour (Fig. 1B and C, Fig. S7A-C and S8). For ease of reference, we refer to this till as ‘Enderby Till’, wherever it is found on the island. The Enderby Formation outcropping on Enderby Island is the most complete exposure of glacial sediments on the islands, and in common with all other outcrops of the sequence is highly indurated. As described by Fleming *et al.* (1976), the formation consists of a lower diamict (Enderby Till Fig. S8A) overlain conformably by massive sands and silts which grade upwards into laminated silts with dewatering structures. Above the laminated silts the sequence becomes coarser, with an increasing percentage of sand evident in laminations which grade into sandy gravels and finally laminated gravels that grade into an overlying upper diamict (Enderby Till). The laminated silts and sands are not found elsewhere on the islands, but the Enderby Till diamict is mirrored at Pillar Rock and Emergency Bay, (Fig. S7 and S8), suggesting that the exposures at Pillar Rock and Emergency Bay record the same glacial event.

20

The pIRIR ages of the laminated sediment in the Enderby Formation are listed in Table 3. All four ages have large relative uncertainties (10–35% at 1σ), mainly because of the small number of grains accepted for D_e estimation, but they are consistent at 2σ with a common value. The samples are effectively from a single unit/event, and statistically indistinguishable from each other, and we therefore compute the weighted-mean age: 384 ± 26 ka. This spans the time interval from about 330 to 440 ka at the 95% confidence interval, corresponding to Marine Isotope Stages 9 to 12.

25



Hydrographic charts hint at the presence of drowned moraine features within the over-deepened valleys of the eastern coasts of both islands, including Norman Inlet on Auckland Island (Fig. S9), and Perseverance Harbour on Campbell Island, which also have a number of promontories shown in the topographic maps that may signify the presence of preserved moraines. Multibeam data show a distinct pair of now-submerged moraines at Shoal and Boyack Points in Perseverance Harbour (Fig. 5), consistent with the over-deepening of the inlet, but no such features were observed at other promontories such as De la Vire and Davis Points (Fig. 5). These moraines provide the first evidence of past glaciation within Perseverance Harbour and we suggest that these moraines mark the greatest preserved extent of valley glaciation, at ~6 km from the valley head.

In addition to the above, we report a number of cirques at higher elevations on both Auckland and Campbell islands, with cirque floor elevations of ~115-233 m a.s.l. and maximum of lengths of ~ 1km; the locations of these are shown in Fig. 1. A UAV survey of one of the most distinct cirques, Lower 'Lake' Speight on Auckland Island (LLS, immediately south of glacial Lake Speight but not itself hosting a lake) captures the geomorphological feature in detail; the cirque is over-deepened, with a vertical back wall and distinct terminal moraines, with no evidence of nested moraines within their limits (Fig. 2). Unfortunately, extensive vegetation or thick peat deposits extending to the highest altitude of the islands bury any potential erratic boulders or glaciated bedrock at our surveyed sites, precluding direct dating of these features, or in-depth geomorphological mapping of the cirque sites. No other terrestrial glacial geomorphology outside the limits of these cirques was observed on either the Auckland or Campbell islands.

3.2 Sediment sequences

The Pillar Rock section provides important constraints on the onset of late Pleistocene glaciation on the Auckland Islands. The exposed section (Fig. 3) consists of a laterally continuous (~10 m), horizontally bedded section exposed by extensive cliff erosion along the north coast of Auckland Island at ~30 m.a.s.l. (Fig. 1C). In places the section is over four m thick, and rests on several metres of deeply indurated diamicton, that we correlate with the upper till exposed on Enderby Island. The 'Enderby Till' is overlain



non-conformably by stratified sands and gravels, which are iron banded. This unit is overlain by a ~2.4 m thick conformable sedimentary succession, that grades upwards from massive silts and sands into laminated silts and clays and into an organic-rich silt band (Fig. S3D). We have dated this organic-rich silt band, with the lower boundary, at a depth of 250 cm in the sequence of $42,230 \pm 1200$ ^{14}C yr BP (45,900 \pm 1250 cal yr BP), and at the upper boundary, at a depth of 237 cm, to $36,860 \pm 600$ ^{14}C yr BP (41,350 \pm 500 cal yr BP). Above this organic silt the succession continues, with a conformable, non-erosive boundary into sandy silts, which become increasingly clast rich and less dense. This conformable sequence is topped by a 10 cm thick clast-supported gravel, topped by a 2 cm layer of flat lying gravel / pebbles. This succession is overlain unconformably by a 27 cm thick sequence of humified dark brown peat, dated between $10,090 \pm 30$ ^{14}C yr BP (11,550 \pm 110 cal yr BP) and 8110 ± 30 ^{14}C yr BP (8960 \pm 80 cal yr BP) at the lower and upper boundaries respectively. Above the peat, there is a series of horizontally bedded silty sandy loams and low organic soil horizons, with pebble horizons, overlain by ~20 cm of sandy loam at the surface.

The two marine sediment cores from Carnley Harbour (Cores 11 and 12, Fig. S2) gave basal peat ages of $12,620 \pm 40$ cal yr BP and $11,690 \pm 120$ cal yr BP. Calibrating the basal radiocarbon ages of peat from across Campbell and Auckland Islands for all sequences with ^{14}C ages exceeding 9 ^{14}C ka BP – including the Carnley Harbour cores – provides an age constraint for the onset of peat growth across the islands (Table 1). The OxCal Phase age calibration of the basal peat samples suggests the onset of peat growth commenced at 17,230-16,110 cal yr BP.

Sediment sequences from key cores both within and directly outside the limits of key cirques (LLS Cirque, LLS Terrace, and Musgrave Harbour, Fig. 1) were dated throughout their profiles and analysed for organic content using LOI. Organic lake mud from the base of the Musgrave Harbour core (Core 10, Fig. 1, Table 2) at 175 cm is dated to $10,340 \pm 60$ cal yr BP. LOI and radiocarbon dating from a core from within the LLS cirque (Site 14, Figs. 1 and 6, Table 2) suggests onset of peat growth in the cirque commenced at $11,700 \pm 140$ cal yr BP; however, an age inversion in the silty material further down the core suggests a relocation of young carbon along the contact zone between the silt and organic layers,



and this constraint is therefore questionable. The LLS Terrace core, however, gives a coherent age-depth profile (Site 15, Figs. 1 and 6, Tables 1 and 2), and shows that organic matter content in the outwash zone of the LLS Cirque increased markedly between $12,310 \pm 50$ and $11,950 \pm 40$ ^{14}C yr BP ($14,200 \pm 120$ and $13,710 \pm 70$ cal yr BP) with no evidence of a reversal in trend. No glacial material was found at the base of the cores.

3.3 Flowline modelling

The 288 model runs span a large range of climatic conditions under which Perseverance Harbour (Campbell Island) could have been glaciated during the late Pleistocene. Figure 7A illustrates the glacier-length predictions of all 288 model runs, coloured according to the MAAT depression used to scale the EDC or SST time series forcing. Simulations that extended beyond the full length of Perseverance Harbour were terminated, consistent with our multibeam survey, which suggested the limits of Late Pleistocene glaciation were well within the constraints of the Harbour; this does not preclude the possibility of extensive glaciation across the island during an earlier period. To constrain the configurations that represent realistic scenarios, the comprehensively radiocarbon-dated peat sequences from Homestead Bog, Campbell Island (Fig. 1D, Table 1) (McGlone et al., 2010; Turney et al., 2006) were used as a constraint. At 40 m a.s.l. on the northern shore of Perseverance Harbour, 3.8 km from the head of the flowline (Fig. 1D, Table 1), the presence of peat in the valley indicates glacier absence from ~ 15 ka through to present. We allowed a 5% tolerance to account for a glacier snout to be slightly more extended in the centre of the valley than on the flanks where the ^{14}C dates were obtained, and to account for any spatial or temporal uncertainties not explicitly accounted for elsewhere. This constraint allowed us to select those simulations that model glacier length to less than 4 km at 15 ka.

The selection described above results in a suite of 102 simulations (Fig. 7B). Of these, the large majority (69%, Fig. 7B inset) – including those which model unrealistic glacial lengths during the Last Interglacial – occur with MAAT depressions that are either smaller than 5 degrees or greater than 7 degrees. These values are inconsistent with the LGM cooling constrained to $\sim 6^\circ\text{C}$ on Auckland and Campbell (McGlone, 2002). By excluding runs with 3, 4 and 8°C MAAT depressions, and those that take no account of the



5 temperature control on precipitation, we further reduced our sample subset to 25 simulations (Fig. 6 and Fig. S8). Each of these runs produced a different pattern of glacier advance and retreat. By taking an ensemble modelling approach and comparing these patterns with the geomorphological and chronological data available from the last 40 ka, it is possible to refine our interpretation of the most likely climatic conditions that prevailed across the New Zealand subantarctic islands during the last glacial cycle.

4 Discussion

Our results show unequivocal evidence of past glaciation, but challenge previous suggestions as to the timing and extent of ice during the LGM (Fleming et al., 1976; Hodgson et al., 2014a; McGlone, 2002).
10 Here we propose the sequence of late Pleistocene events that best explains our field and modelling datasets from the New Zealand subantarctic Campbell and Auckland Islands.

4.1 The Middle Pleistocene

We interpret the Enderby Formation as a glacial succession that records an oscillating ice margin, with
15 two extensive deposits of subglacial diamicton separated by deposition of ice-proximal glaci-fluvial sedimentation that was overrun by the later ice advance evidenced by the upper diamicton (Fig. 4, Figs. S7B and S8A). The conformable nature of the sequence suggests that there was not a significant hiatus between the two glacial advances recorded by the diamictons; rather the glaciofluvial sediments sandwiched between the two diamictons were formed during a brief period of recession and subsequent
20 advance during a sustained glaciation. The location of this glacial till on the north shore of Enderby Island has led to previous studies (Fleming et al., 1976; Hodgson et al., 2014a; McGlone, 2002) suggesting this was deposited by a valley glacier emanating from Port Ross, Auckland Island. However, this study has identified similar till at multiple sites across the Auckland Islands; at Pillar Rock these deposits (stratigraphically below those described in Fig. 3) are found exposed in high coastal cliffs, with those at
25 both Pillar Rock and Emergency Bay (Fig. 1) located outside valley settings. This suggests a period of extensive glacial cover of the Auckland Islands (henceforth referred to as the ‘Enderby Glaciation’), perhaps as a small ice cap, far beyond the scope of the valley glaciers previously suggested to have been the greatest extent of glaciation on the islands.



The highly indurated nature of the Enderby Formation suggests antiquity, but in the absence of direct chronological control the exposure has been interpreted to be LGM in age. This is overturned by the IRSL dating of the Enderby Formation glaciofluvial sediments to 384 ± 26 ka (Table 3). This places the Enderby
5 Glaciation firmly in the Middle Pleistocene, most likely during MIS-10 (~ 374 ka), or potentially MIS-12 (~ 424 ka), both significant glacial stadials.

The timing of this glaciation is consistent with other locations in the Southern Hemisphere. Notably, recent work on the Boco Plain in Tasmania (Augustinus et al., 2017) has identified a succession of glacial
10 advances, the most recent of which (prior to the LGM) was 378 ± 22 ka during Marine Isotope Stage 10 (the Boco Glaciation). Intriguingly, the recognition of two lateral moraines, one overlying the other, suggests an oscillating ice margin similar to that observed on Enderby Island, and may represent a common regional climate-driver. Similarly, ^{10}Be -dated glacial moraines in the Lago Buenos Aires region also indicates maximum glaciation prior to the LGM between ca. 450 and 340 ka (Augustinus et al., 2017;
15 Kaplan et al., 2005).

The hemispheric-wide nature of this major glaciation appears to be associated with extreme southern glacial stadials between 420 and 340 ka (Marine Isotope Stages 10 and 12) (Bard and Rickaby, 2009b; Li et al., 2010). Off southern Africa, a SST reconstruction suggests that the Subtropical Front (and the associated westerly winds) migrated north by some 7° of latitude during both glacial periods, limiting the
20 leakage of the Agulhas Current (warm, saline water) into the south Atlantic Ocean, weakening the Atlantic Meridional Overturning Circulation and amplifying the severity of glaciations in the Southern Hemisphere and possibly globally (Bard and Rickaby, 2009a), the opposite to that suggested for periods of super-interglacial warmth (Turney and Jones, 2010). SSTs during MIS-10 and -12 were 6°C cooler
25 than the Holocene, compared to a depression of just 4°C at the LGM/MIS-2. Given the numerous glacial advances between ~ 370 and 390 ka across the mid to high-latitudes of the Southern Hemisphere, and our weighted-mean age of 384 ± 26 ka for the laminated sediments, we favour attributing the Enderby Formation to MIS-10. Regardless of the glacial stage attribution, our results from the New Zealand



subantarctics imply that extensive glacial deposits on other subantarctic islands cannot be assumed to be LGM in age (Balco, 2007; Hodgson et al., 2014b) and may similarly be of middle Pleistocene origin.

4.2 The Late Pleistocene at Pillar Rock, Auckland Island

5 The interpretation of the rare exposed sedimentary sequence at Pillar Rock and the chronological control it affords is crucial to the glacial history of the islands, given that at present, the interpretation rests on the previously held assumption that the lower indurated glacial diamicton, the ‘Enderby Till’, is of LGM age. The stratigraphy and chronology of this sequence suggest this cannot be the case. The presence of a conformable succession of unindurated soft sediments at Pillar Rock overlying the ‘Enderby Till’ and
10 clearly predating the LGM is important. The depositional environment of the sequence is a relatively flat coastal plain, currently peat bog; we therefore interpret this unit as capturing the gradual infill of a depression with minimal slope processes at work, most likely during the last glacial cycle. Lower unstratified sands, prograding up into laminated silts and sands, are overtopped by a ~13 cm thick organic silt band, which has calibrated ages of ~46 cal ka BP and ~41 cal ka BP, eliminating the possibility that
15 the underlying Enderby Till is coeval with the LGM. We suggest that this organic unit was formed under anoxic conditions, as the basin or depression filled and ponded water and organic material over several millennia, and may correspond to Antarctic Isotopic Maximum 12, which represents significant warming in the EPICA Dome C Core in East Antarctica (EPICA, 2006) at ~48 ka (Fig. 9).

20 The Pillar Rock sequence preserves a record of ongoing sedimentation, most likely during the LGM, with the upper layer of the succession being indicative of a winnowed lag deposit, formed under cold, periglacial conditions. This interpretation is supported by the presence of an overlying thick humified peat deposit, lying unconformably above this succession, formed between ~11.6 cal ka BP and ~9 cal ka BP, during the early Holocene climatic amelioration. The lack of peat above this unit suggests the area
25 had become free draining soon after ~9 cal ka BP, most likely due to erosion of the cliffs surrounding the site of the basin or depression and led to a change in the depositional environment, alternating between periods of soil formation and low organic sedimentation at this exposed location. This could be caused



by increasing strength in the westerly winds throughout this period, which may have increased cliff erosion and led to the transfer of sands and silts inland (McGlone, 2002).

4.3 Evidence for limited Late Pleistocene glaciation?

5 The geomorphological evidence reported here suggests that at least one glaciation took place subsequent to the Enderby Glaciation. High-altitude cirques, with pronounced terminal moraines and no evidence of nested inner moraines, show a period of limited but sustained glaciation; organic lake sediments from outside the limits of the Musgrave Harbour cirque dated to ~10 ka cal BP suggest that there was no glacial sediment input from this time. Most significant is the dated sediment sequence from the LLS Terrace core
10 (Core 15, Fig. 1, Table 2), directly outside the limit – and within the sediment outwash zone – of the LLS Cirque. The sharp increase in organic matter content observed in the Terrace core (Fig. 6) continues uninterrupted through the Antarctic Cold Reversal (ACR; 14,500-12,500 cal yr BP (EPICA, 2006)), which was a period of extensive glacier growth in mainland New Zealand and Patagonia (Fogwill and Kubik, 2005; Pedro et al., 2015; Putnam et al., 2010). That the LLS Terrace core shows no inorganic
15 sediment input throughout this period strongly indicates that there was no substantial ACR glaciation in the cirque above this site. This interpretation corresponds with temperature reconstructions from Campbell Island (McGlone et al., 2010) (Fig. 9), which show minimal temperature depression during the ACR. The mean age for the onset of basal peat growth across Auckland and Campbell islands shows that significant climate amelioration had commenced by ~16.6 ka cal BP (Fig. 9), signifying the end of the
20 periglacial conditions across lower latitudes of the islands, and constraining the widespread onset of peat growth, which continued uninhibited by subsequent post-glacial cool periods, including the ACR. Furthermore, peat dated to ~12.6 ka cal BP from two marine cores taken from Carnley Harbour (Cores 11 and 12, Table 1) show that sea level was significantly lower and that this low-lying area was ice-free at the time. It is possible that the later onset of peat growth and imperfect age models within the cirques
25 were caused by the presence of perennial snow patches throughout the post-glacial and early Holocene period (Watson, 1966); however, it is clear that the expression of the ACR was limited over the subantarctic islands in the southwest Pacific, and did not result in glaciation.



We suggest that the lack of evidence for ACR glaciation strongly implies that the pronounced cirques observed across the Auckland and Campbell Islands (Fig. 1) originated during an earlier glacial; we suggest this to be either the LGM or Southern Hemisphere eLGM. There are two potential scenarios for this cirque glaciation: one, a limited valley glaciation of the islands during the eLGM/LGM, followed by retreat to the cirques before final deglaciation; or two, the cirques represent the maximum extent of eLGM/LGM glaciation. Given the lack of glacial geomorphology observed outside the limits of the cirques, we suggest the latter scenario is more likely, but more work is needed to confirm this interpretation. It must be noted that as the cirques (and associated features i.e. moraines) are not directly dated, it is possible that cirque formation predates the eLGM/LGM. Regardless of which scenario is correct, however, it is clear that glaciation of Campbell and Auckland Islands during the LGM was far more restricted than previously supposed (Fleming et al., 1976; Hodgson et al., 2014a; McGlone, 2002).

4.4 Modelling Late Pleistocene glaciation

The Perseverance Harbour moraines revealed on Campbell Island by the multibeam survey indicate the presence of a glacier terminating 6 km from the head of the valley and close to sea level; however, although more extensive than the LGM Auckland and Campbell cirques, this valley glaciation is on a much smaller scale than the Enderby Glaciation. The high level of preservation of these moraines strongly implies that another glacial event in the New Zealand subantarctic islands occurred prior to the LGM but after the Enderby Glaciation. To test this, and the hypothesis that the LGM glaciation was limited, we ran a series of flowline model experiments. 25 model runs of Late Pleistocene glaciation in Perseverance Harbour fit the constraints of the field data (Fig. 8 and Fig. S10), most importantly, the constraint from Homestead Bog, situated at 3.8 km from the head of the model flow line, and shown to be deglaciated by at least ~15ka. The vast majority of these 25, however, reach a maximum extent of just 3-4 km, which is incompatible with the length of glacier needed to form the Perseverance Harbour moraines reported here. Importantly, though, one model run does capture a significant glacier advance to >5 km at 67.7 ka (Fig. 8A), followed by retreat to 2.3 km and a subsequent limited advance to 3.2 km during the LGM, before rapid deglaciation at 15 ka prior to the ACR (Fig. 8A).



Our model simulations suggest that the Perseverance Harbour moraines (and similar valley glaciers on the east coast of Auckland Island) were formed around 68 ka, consistent with the New Zealand glacial maximum of the so-called Otira Glaciation at 62-72 ka during MIS-4 (Williams et al., 2015). The model suggests that the islands were incapable of hosting a more extensive glaciation than this at any point over the past 125 ka, consistent with our interpretation that it took a step-change in the regional climate system to support the extensive MIS-10 Enderby Glaciation. Another important consideration is the loss of catchment to the west through marine erosion and glacial attrition or tectonic subsidence (Summerhayes, 1967), which may have diminished the available accumulation area for more ‘recent’ glacials, further reducing the capability of the New Zealand subantarctic islands to sustain glaciation on the scale seen during MIS-10.

4.5 Late Pleistocene climatic implications

Given the close correspondence of the field data, we therefore propose that the climatic conditions of the model run shown in Fig. 8A provide a reasonable first-order representation of change over the New Zealand subantarctic islands during the late Pleistocene. For the LGM, our study implies a MAAT depression of 5°C, a similar level of cloudiness as today, an annual temperature range of 11°C, and a substantial precipitation reduction of ~30%. Crucially, seasonality appears to have played an important role in glaciation on these islands; our simulation with identical parameters to the above but with less severe winter cooling (Fig. 8B) fails to reach the glacier length necessary to form the Perseverance Harbour moraines.

The same MAAT depression of 5°C obtained by the colder summers and milder winters of the modelled scenario in Fig. 8B would have placed considerable pressure on the viability of tall shrubs, which would never have reached the required number of degree growing days to sustain growth. Crucially, the high seasonality implied by the 11°C temperature range invoked in the modelled scenario in Fig. 8A – biased towards the winter – would have allowed for the continued survival of plant life on the islands suggested by the high number of endemic species (McGlone et al., 2010; Mitchell et al., 2014; Van der Putten et al., 2010). The milder summers in this scenario means that the lowland shrubs and herbs found across the



islands would have been able to survive through the colder winters. Some woody species (e.g. *Dracophyllum* spp. and *Myrsine divaricata*) are likely to have survived through the LGM on both Auckland and Campbell Islands as they both appear very early in the postglacial period and can be shown to have been present pre-LGM on Auckland Island (Fleming et al., 1976; McGlone, 2002). The current absolute limit to shrubland containing these species on Campbell Island is c. 300 m (McGlone et al., 1997), which equates to a warmest month temperature of $\sim 7^{\circ}\text{C}$. If allowance is made for a fall in mean sea level of around 120 m, the maximum depression in MAAT at the LGM that would permit these shrubs and small trees to survive is $\sim 3^{\circ}\text{C}$. As pointed out by McGlone (2002) there is a mismatch here with previous estimates of LGM cooling of $\sim 6^{\circ}\text{C}$ as such a temperature depression, if it affected summer and winter equally, would exceed the tolerance of these woody species. However, a MAAT depression of 5°C , combined with an increased warmest month/coolest month temperature range of 11°C (versus the current 4.7°C), would not only ensure the survival of these woody species but also other species including the functionally flightless Campbell Island and Auckland Island teal (*Anas nesiotis* and *A. aucklandicus*) which, on the basis of molecular dating, have occupied the islands over several glacial-interglacial cycles (Mitchell et al., 2014).

We therefore propose that reduced winter precipitation and high seasonality shown by our flowline modelling combined to limit the LGM glaciation of the New Zealand subantarctic islands despite the large MAAT depression. This is consistent with work showing an expansion of winter sea ice in the Southern Ocean, reducing the moisture available to the atmosphere in this region (Fogwill et al., 2015; Otto-Bliesner et al., 2006). Whilst palaeo records of Antarctic sea-ice extent are incomplete, our flowline modelling is consistent with most proxy and model data, which suggest a significant northwards expansion of sea ice to within the region of the New Zealand subantarctic islands both during the LGM (Roche et al., 2012) and the ACR (Pedro et al., 2015) with a strong seasonal cycle (Gersonde et al., 2005). These studies are further supported by genetic analysis of kelp across the Southern Ocean that shows that kelp recolonized the subantarctic islands, including Campbell and Auckland Islands, relatively recently, a finding that has been interpreted to suggest the presence of sea ice across the region at the LGM (Fraser et al., 2009). Complementary studies on atmospheric circulation during the LGM suggest an equatorward



5 shift with a specific focus on mid-latitude westerly airflow across this region (Hesse, 1994; Jaccard et al., 2013; Kohfeld et al., 2013); this could have reduced sea ice break-up and storm frequency over the subantarctic islands, further reducing precipitation. Comparing our results to other subantarctic islands at similar latitudes, we note that southwest Pacific Macquarie Island (54.62°S) and south Atlantic South Georgia (54.25°S) may have experienced similarly restricted glaciation during the late Pleistocene (Bentley et al., 2007; Hodgson et al., 2014a), whilst Stewart Island (47°S) to the south of the New Zealand mainland also had a very limited LGM extent compared to mainland New Zealand (Brook, 2009), suggesting this may have been a Southern Ocean-wide mechanism.

10 5 Conclusions

Palaeo-archives and modelling from the New Zealand subantarctic islands provide important new insights into the glacial history and changing climate of the Pacific sector of the Southern Ocean. We find evidence for multiple distinct phases of glaciation on both Auckland and Campbell Islands, and hypothesize the following sequence of events: 1) the ‘Enderby Glaciation’ at ~380 ka consisting of extensive glaciation covering large portions of the islands (potentially as small ice caps), most likely occurring during MIS 15 10; 2) a more restricted glacial period prior to the LGM that formed substantial valley glaciers on Campbell and Auckland Islands at around 72-62 ka; and 3) a limited eLGM/LGM glacial advance, comprising of either valley glaciation with glaciers no longer than a few kilometers prior to retreat to high-altitude cirques, or of severely restricted glaciation limited to these cirques. We find no evidence for 20 glaciation on the islands after the LGM.

The disparity between limited late Pleistocene glaciation of the New Zealand subantarctic islands and the widespread expansion of glaciers across southern Pacific landmasses (such as Patagonia, Tasmania and New Zealand) raises important questions about our understanding of changing climatic conditions across 25 the Southern Ocean. The Auckland and Campbell Islands were apparently incapable of sustaining significant glaciation during the LGM or ACR despite the pronounced and sustained atmospheric and oceanic temperature depressions, suggesting other factors had a greater influence than temperature changes alone. Our results are consistent with precipitation starvation caused by the northwards expansion



of winter sea ice in the Southern Ocean, which overwhelmed regional atmospheric and oceanic temperature depressions and resulted in limited glaciation on the subantarctic islands, providing important new insights into the significance of sea-ice extent and precipitation regime shifts across the Southern Ocean. In marked contrast, during the Middle Pleistocene, the minimum in eccentricity (100 kyr and 400 kyr cycles) appears to have driven a northward migration of the Subtropical Front. The resulting substantial reduction in Agulhas Current leakage into the Atlantic Ocean depressed southern SSTs over an extended period, sufficient to sustain significant glaciation on the low-altitude New Zealand subantarctic islands. This varying response of Pleistocene glaciers on the Auckland and Campbell Islands, highlighting the sensitivity of the Southern Ocean region, and suggesting that step-changes in atmospheric and oceanic regimes can have extreme climatic and environmental impacts.

Appendix 1 – extended methodology

1. Geomorphological survey

Relevant sites were targeted with an Unmanned Aerial Vehicle (UAV) to produce high-resolution digital elevation models (DEMs.) Using a Sensefly ‘eBee’ mapping drone, imagery was acquired, taking aerial photographs with significant longitudinal and latitudinal overlaps (80% and 85% respectively) to produce a high-resolution (8 cm per pixel) image mosaic and digital surface model. Imagery was fully georeferenced using GPS data and senseFly’s eMotion post-processing flight planning software. Digital elevation models (DEMs) of survey sites were compiled using Post Flight Terra 3D 3, and imagery was produced with Global Mapper v16.

Multibeam mapping was undertaken with a Kongsberg EM3002 multibeam echo sounder, which was deployed from a side mounted swinging arm. The system is well-suited for detailed seafloor mapping in water depths from less than 1 metre up to typically 200 metres in cold oceanic conditions. The resultant images were fully georeferenced and simultaneously processed at sub-decimetre accuracy with an integrated differential GPS system.



2. Sedimentary analysis

Lacustrine and peat sediments were recovered using a 5-cm wide ‘D’-section (‘Russian’) corer. The organic units of the Pillar Rock formation were sampled with monolith tins. Sediments were described in the field, with samples wrapped and transported back to the laboratory where they were cold-stored (4°C) prior to analysis. Key sedimentary sequences were analysed for organic content by loss-on-ignition (LOI). The LLS Cirque core was subsampled (sample size 0.5 cm) every 2 cm from 64-104 cm and every 1 cm thereafter. The LLS Terrace core was subsampled (sample size 0.5 cm) every 5 cm from 0-95 cm, and every 2.5 cm thereafter. These samples were dried overnight in an oven at 105°C and then heated in a muffle furnace at 550°C for 4 hours. The LOI was calculated as the percentage dry weight (Heiri et al., 2001).

The shallow marine sediment cores from Carnley Harbour (Cores 11 and 12, Fig. 1, Table 1) were taken using a small gravity piston corer with 150cm plastic core barrels deployed over the side of the expedition vessel. The piston corer has an impact-triggered seal that creates a vacuum seal on the core barrel once the corer is retrieved, capturing the water sediment interface and preventing ‘wash-out’ of fine sediment during retrieval. Over 1 m of sediment was recovered from each deployment of the corer, with the cores consisting mostly of peats overlain by fine silts containing occasional marine mollusk shells in life position.

Samples were taken from the face of the Enderby Formation laminated silt at intervals of approximately 10 cm (Fig. 4). These were prepared for particle size analysis using a Saturn Digitiser and the results categorised using the Udden-Wentworth Scale (McCave and Syvitski, 1991).

3. Chronology

3.1. Radiocarbon dating

The peat sediments analysed here were given an acid-base-acid (ABA) pretreatment and then combusted and graphitized in the University of Waikato accelerator mass-spectrometry (AMS) laboratory, with $^{14}\text{C}/^{12}\text{C}$ measurement by the University of California at Irvine (UCI) on a NEC compact (1.5SDH) AMS



system. The pretreated samples were converted to CO₂ by combustion in sealed pre-baked quartz tubes, containing Cu and Ag wire. The CO₂ was then converted to graphite using H₂ and a Fe catalyst, and loaded into aluminium target holders for measurement at UCI. To estimate the timing of the onset of peat growth we used the Phase model option in OxCal v.4.2.4 (Bronk Ramsey and Lee, 2013) with General
5 Outlier analysis detection (probability = 0.05) (Ramsey, 2009). Importantly, the Phase option is a grouping model which assumes no geographic relationship between samples, simply that the ages represent a uniform distribution between a start and end boundary. The ¹⁴C ages were calibrated against the Southern Hemisphere calibration (SHCal13) dataset (Hogg et al., 2013) and reported here as mean calendar years (yr) or thousands of years (ka) cal BP ± 1σ uncertainty (Tables 1 and 2). Radiocarbon ages
10 are distinguished from calibrated ages by being expressed as ¹⁴C yr/ka BP.

3.2. Optical dating

Each sample was treated using routine procedures to extract sand-sized grains of K-feldspar (Aitken, 1998), treated sequentially with HCl acid and H₂O₂ to remove carbonates and organic matter, and then
15 dried. Grains in the size range of 90–125 μm (Enderby-OSL2, -OSL3 and -OSL4) or 180–212 μm (Enderby-OSL1) were isolated by dry sieving, and the K-feldspar grains separated from heavier minerals using a sodium polytungstate solution with a density of 2.58 g/cm³. K-feldspar grains were etched using 10% HF acid for 40 min to clean their surfaces and reduce the thickness of the alpha-irradiated outer layer of each grain. The IRSL measurements were made on an automated Risø TL-DA-20 reader using both
20 single-aliquot and single-grain techniques. Aliquots consisting of several hundred grains were prepared by mounting the grains as a monolayer (~5 mm in diameter) on stainless steel discs, using ‘Silkospray’ silicone oil as the adhesive, and were stimulated using infrared diodes (870 Δ 40 nm, 135 mW/cm²). For the single-grain measurements, we used discs drilled with 100 holes, each 300 μm in diameter and depth, and stimulated the grains individually using a focussed infrared laser (830 Δ 10 nm, 400 W/cm²) (Bøtter-
25 Jensen et al., 2003). Radiation doses were given using a ⁹⁰Sr/⁹⁰Y beta source mounted on the reader and calibrated for both multi-grain aliquots and individual grain positions. The IRSL signals were detected using a photomultiplier tube, after passing through a filter pack containing Schott BG-39 and Corning 7-59 filters to transmit wavelengths of 320–480 nm. We used single aliquots to conduct laboratory tests of



anomalous fading, but estimated the D_e values from measurements of individual grains of K-feldspar (Duller, 2008; Jacobs and Roberts, 2007), enabling the identification and elimination of any grains with aberrant luminescence characteristics (David et al., 2007; Feathers, 2003; Feathers et al., 2006; Jacobs et al., 2006; Jacobs et al., 2011; Jacobs et al., 2008; Roberts et al., 1998; Roberts et al., 1999) prior to age determination. We did not apply a residual-dose correction, because our samples have high natural doses (>400 Gy).

In this study, we adopted a two-step single-grain pIRIR regenerative-dose procedure (Blegen et al., 2015; Guo et al., 2016) to determine the D_e values for individual sand-sized grains of K-feldspar. In this procedure (see Table S1 for details), an initial infrared stimulation is made at 200 °C for 200 s using infrared diodes, so that all 100 grains on each single-grain disc are stimulated simultaneously. Li and Li (2012) showed that, compared to an initial infrared stimulation at 50 °C, the fading component is removed more effectively at a temperature at 200 °C. The pIRIR signals used for D_e determination were then measured for individual grains at 275 °C, with each grain stimulated by an infrared laser for 1.5 s; this duration proved sufficient to reduce the pIRIR intensity to a low and stable level (Fig. S3). We chose a stimulation temperature of 275 °C, rather than 290 °C (Thiel et al., 2011), to avoid any detrimental effects from significant thermal erosion during the single-grain measurements. The latter can occur at stimulation temperatures close to the preheat temperature (320 °C), because laser stimulation of the final grain on each 100-grain disc begins more than 2.5 min after the first grain on the disc. Figure S3 shows a typical pIRIR decay curve and associated dose response curve for a single grain of K-feldspar from Enderby-OSL1.

We conducted a dose recovery test (Galbraith et al., 1999; Roberts et al., 1999) on Enderby-OSL1 to validate the suitability of the pIRIR experimental conditions used to measure these samples (Table S1). A total of 600 grains were bleached for ~6 hr using a Dr Hönle solar simulator (model: UVACUBE 400) and then given a dose of ~300 Gy, before being measured using the procedure of Table S1. To select reliable single grains for D_e determination and reject unsuitable grains, we applied quality-assurance criteria similar to those proposed for single grains of quartz (Jacobs et al., 2006) and K-feldspar (Blegen



et al., 2015). Grains were rejected if they exhibited one or more of the following five properties: 1. Weak test-dose signal; 2. High level of recuperation; 3. Poor recycling ratio; 4. Poorly fitted dose response curve (DRC); or 5. Natural pIRIR signal equal to or greater than the saturation limit of the DRC. We found that most of the grains (583 grains) were rejected by criterion 1 because of weak T_n signals. Among the remaining 17 grains, two had L_n/T_n ratios greater than the saturation limit of the DRC, two had poorly fitted DRCs, and one grain suffered from high recuperation. As a result, only 12 grains were accepted for dose estimation. The measured-to-given dose ratios for these grains are shown in Fig. S4a. The weighted mean of these 12 dose recovery ratios is 1.06 ± 0.07 , which is consistent with unity and suggests that the pIRIR procedure in Table S1 yields reliable D_e values for the Enderby samples. We also tested the Enderby samples for anomalous fading of the pIRIR signal. Twelve aliquots of Enderby-OSL1 were measured using a single-aliquot IRSL procedure similar to that described by Auclair et al. (2003), but based on fading measurements of the pIRIR signal. As our samples emitted relatively dim signals, we used the procedure of Thiel et al. (2011), in which the initial infrared stimulation is made at 50 °C and the pIRIR signal is subsequently stimulated at 290 °C. Doses of ~200 Gy were administered using the laboratory beta source, and the irradiated aliquots were then preheated and stored for periods of up to 1 week at room temperature (~20 °C). Figure S4b shows the decay in the sensitivity-corrected pIRIR signal as a function of storage time, normalised to the time of prompt measurement. The corresponding fading rate (g -value: (Aitken, 1985; Huntley and Lamothe, 2001) is $0.6 \pm 0.7\%$ per decade, which is consistent with zero fading at 1σ . Li and Li (2012) demonstrated that a higher initial infrared stimulation temperature (e.g., 200 °C) could remove the fading component more effectively than stimulating at 50 °C, so the pIRIR signal measured after an initial infrared bleach at 200 °C (Table S1) should be stable. Hence, no fading correction has been made to the measured D_e values for the Enderby samples.

We consider the D_e values obtained for all samples as effectively single-grain estimates. Grains of 90–125 μm diameter were used for Enderby-OSL2, -OSL3 and -OSL4, so each hole on the disc may contain ~8 grains for these samples. For Enderby-OSL1, each hole was occupied by a single grain of 180–212 μm in diameter, and more than 95% of the total pIRIR signal was contributed by fewer than 5% of the



measured grains (Fig. S3b). For Enderby-OSL2, -OSL3 and -OSL4, therefore, we deduce that the measured pIRIR signal arises from only one or two grains in each of the holes.

Given the paucity of K-feldspar grains available for each of the Enderby samples, only 600, 600, 300 and 500 grains were measured for Enderby-OSL1 to -4, respectively. After applying the same five rejection criteria as were used in the dose recovery test, we found that ~20% of the accepted grains had natural signals equal to or greater than the saturation limit of their DRCs; finite D_e values could not be estimated for these grains. As a result, we expect that the D_e estimates for the remaining accepted grains will likely underestimate the true D_e value for each sample, because the rejection of the saturated and ‘oversaturated’ grains will lead to a truncated D_e distribution (Duller, 2012; Guo et al., 2017; Li et al., 2016; Thomsen et al., 2016). To avoid this problem, we applied the new method proposed by Li et al. (2017), in which a standardised growth curve (SGC) is established and the weighted-mean L_n/T_n ratio for all accepted grains is projected on to the corresponding SGC to determine the sample D_e . In this method, no grains are rejected due to saturation issues, so a full (untruncated) distribution of L_n/T_n ratios is obtained. Based on their experimental and simulation results, Li et al. (Li et al., 2017) showed that this method can produce reliable D_e estimates up to 5 times the D_0 value of the DRC, which is far beyond the conventional limit of $2D_0$.

We first investigated if individual grains from each of the Enderby samples could be used to develop a SGC. Since the latter should be established using only grains that are considered well-behaved (Li et al., 2016) we applied the same rejection criteria as described above, except that grains with natural signals in saturation were accepted for purposes of establishing the SGC. The L_x/T_x ratios of all of the accepted grains from each of the samples are plotted against the corresponding regenerative doses in Fig. S5a, which shows the large variation in L_x/T_x ratios at any particular dose. We then applied the least-square normalisation (LS-normalisation) procedure of Li et al. (2016) to these data, resulting in significantly reduced scatter and a common SGC for the accepted grains from each of the samples (the best-fit curve in Fig. S5b). Next, the L_n/T_n ratio for each grain was multiplied by a scaling factor determined from the LS-normalisation procedure (Li et al., 2016). The distributions of LS-normalised single-grain L_n/T_n ratios



are displayed in Fig. S6. Enderby-OSL1 (Fig. S6a) and -OSL2 (Fig. S6b) both appear to have two distinct populations of grains, with the vast majority of grains (75% in Enderby-OSL1 and 94% in Enderby-OSL2) comprising one component and the other component consisting of 2–3 grains with much smaller L_n/T_n ratios; we interpret the latter as younger, intrusive grains. We applied the finite mixture model (FMM; (Roberts et al., 2000)) to both distributions determine the weighted-mean L_n/T_n ratio for each component, assuming a two-component mixture. In the model, the L_n/T_n overdispersion value was varied between 0 and 40%, and the lowest Bayes Information Criterion score was used to identify the optimum fit (David et al., 2007; Jacobs et al., 2011; Jacobs et al., 2008).

10 In contrast, Enderby-OSL3 (Fig. S6c) and -OSL4 (Fig. S6d) each appear to consist of a single population of grains, so we determined the weighted-mean L_n/T_n ratio for these two distributions using the central age model (CAM (Galbraith et al., 1999)). The weighted-mean ratio for Enderby-3 is based on just 4 grains, so the relative standard error on this estimate is correspondingly large (35%). The weighted mean L_n/T_n ratios for Enderby-OSL3 and -OSL4, and for the major FMM components of Enderby-OSL1 and -
15 OSL2, were then projected on to the SGC to estimate the corresponding D_e values (Table 3).

The environmental dose rates were estimated from the sediment samples recovered from each tube hole. The total dose rate to K-feldspar grains consists of 4 components: the external gamma, beta and cosmic-ray dose rates, and the internal beta dose rate. The external gamma and beta dose rates were estimated
20 from thick-source alpha counting (Aitken, 1985) and low-level beta counting (Bøtter-Jensen and Mejdahl, 1988; Jacobs and Roberts, 2015). The minor contribution from cosmic rays was estimated from the present-day burial depth of the samples and the latitude, longitude and altitude of the site (Prescott and Hutton, 1994). These external components of the total dose rate were adjusted for water content, assuming
25 a long-term value of 25% (which falls within the range of measured water contents: 17–26%) and a standard error of $\pm 6\%$ to capture any plausible variations in time-averaged water content over the entire period of sample burial. The internal beta dose rate was estimated by assuming K and Rb concentrations of $10 \pm 2\%$ and 400 ± 100 ppm, respectively (Huntley and Hancock, 2001; Smedley et al., 2012). The dosimetry data are summarised in Table 3.



4. Flowline model description

The glacier model employed is essentially the one-dimensional shallow-ice approximation (SIA) flowline model of Golledge & Levy (2011), with two modifications implemented for this study. First, the positive degree-day (PDD) scheme formerly used to calculate surface mass balance (SMB) has been replaced by a simplified energy-balance scheme, following the insolation-temperature melt method (Eq. 16 of Robinson et al., (2010)). This formulation allows for orbital variability to be included in the calculation of SMB by incorporating time-dependent insolation values from the solution of Laskar *et al.*, (2004). Shortwave radiation is specifically accounted for (unlike in the PDD scheme), since SMB is controlled in part by atmospheric transmissivity (the ratio between downward shortwave radiation at the land surface and at the top of the atmosphere). Atmospheric transmissivity is modified from a base value according to surface elevation (Robinson et al., 2010). Spatial and temporal variations in surface albedo are included, with values ranging from 0.7 (snow) to 0.05 (ocean). A fractional albedo value is calculated to account for changes in surface type within each cell during the year. Resultant snowfall is prevented from accumulating on slopes steeper than 35° and instead is redistributed (progressively if necessary) to the next downslope cell.

The second modification to the model is the way in which basal sliding velocities are calculated. Previously we used a triangular averaging scheme (Kamb and Echelmeyer, 1986) to smooth calculated driving stress values, which were then used to calculate sliding velocities using a rate factor and sliding exponent (Eq. 8 of Golledge and Levy (2011)). Here we follow instead the approach of Bueler & Brown (2009), and superpose velocity solutions from the SIA with those from the shallow-shelf approximation (SSA). The latter essentially represents the sliding component of glacier flow, by explicitly incorporating the effects of longitudinal stresses. Our iterative scheme follows MacAyeal (1997): ice thickness and the horizontal velocity gradient are first used to calculate an effective viscosity, from which the horizontal strain rate is determined and then used to update the calculated sliding velocity. We use a coefficient to modify calculated sliding values to account for unknown basal drag.



Data availability

Data will be lodged on the NOAA Paleoclimate Archive

Acknowledgements

5 First and foremost, we must acknowledge the contribution of Dr Richard Jones, who was integral to this work, and is dearly missed. The work was supported by the Australasian Antarctic Expedition 2013-2014, the Australian Research Council (FL100100195, FT120100004, DE130101336 and DP130104156) and the University of New South Wales. MM and JW were supported by Core Funding for Crown Research Institutes, from the New Zealand Ministry of Business, Innovation and Employment's Science and Innovation Group. A major thanks to the captain and crew of the *MV Akademik Shokalskiy*, James
10 MacDiarmid and Ben Fink, and Henk Haazen and Kali Kahn on the *Tiama* for all their help in the field. Research on the New Zealand subantarctic Auckland and Campbell Islands was undertaken under the New Zealand Department of Conservation National Authorisation Numbers 37687-FAU and 39761-RES. Thanks to Jenny Dahl of the Rafter Radiocarbon Laboratory for advising on radiocarbon age NZA4509.

15

References

- Aitken, M. J.: Thermoluminescence dating, Academic press, 1985.
- Anderson, R. F., Ali, S., Bradtmiller, L. I., Nielsen, S. H. H., Fleisher, M. Q., Anderson, B. E., and Burckle, L. H.: Wind-Driven Upwelling in the Southern Ocean and the Deglacial Rise in Atmospheric CO₂, *Science*, 323, 1443-1448, 2009.
- 20 Auclair, M., Lamothe, M., and Huot, S.: Measurement of anomalous fading for feldspar IRSL using SAR, *Radiation measurements*, 37, 487-492, 2003.
- Augustinus, P., Fink, D., Fletcher, M.-S., and Thomas, I.: Re-assessment of the mid to late Quaternary glacial and environmental history of the Boco Plain, western Tasmania, *Quaternary Science Reviews*,
25 160, 31-44, 2017.
- Balco, G.: A surprisingly large marine ice cap at Heard Island during the Last Glacial Maximum. In: *Antarctica: A Keystone in a Changing World e Online Proceedings of the 10th ISAES*. USGS Open-File Report, 2007.
- Bard, E. and Rickaby, R. E. M.: Migration of the subtropical front as a modulator of glacial climate,
30 *Nature*, 460, 380-383, 2009a.
- Bard, E. and Rickaby, R. E. M.: Migration of the subtropical front as a modulator of glacial climate, *Nature*, 460, 380, 2009b.
- Barrows, T. T., Lehman, S. J., Fifield, L. K., and De Deckker, P.: Absence of cooling in New Zealand and the adjacent ocean during the younger dryas chronozone, *Science*, 318, 86-89, 2007.



- Bentley, M. J., Evans, D. J. A., Fogwill, C. J., Hansom, J. D., Sugden, D. E., and Kubik, P. W.: Glacial geomorphology and chronology of deglaciation, South Georgia, sub-Antarctic, *Quaternary Science Reviews*, 26, 644-677, 2007.
- 5 Blegen, N., Tryon, C. A., Faith, J. T., Peppe, D. J., Beverly, E. J., Li, B., and Jacobs, Z.: Distal tephras of the eastern Lake Victoria basin, equatorial East Africa: correlations, chronology and a context for early modern humans, *Quaternary Science Reviews*, 122, 89-111, 2015.
- Bøtter-Jensen, L., Andersen, C., Duller, G. A., and Murray, A. S.: Developments in radiation, stimulation and observation facilities in luminescence measurements, *Radiation Measurements*, 37, 535-541, 2003.
- 10 Bøtter-Jensen, L. and Mejdahl, V.: Assessment of beta dose-rate using a GM multicounter system, *International Journal of Radiation Applications and Instrumentation. Part D. Nuclear Tracks and Radiation Measurements*, 14, 187-191, 1988.
- Bronk Ramsey, C. and Lee, S.: Recent and planned developments of the program OxCal, *Radiocarbon*, 55, 720-730, 2013.
- 15 Brook, M. S.: GLACIATION OF MT ALLEN, STEWART ISLAND (RAKIURA): THE SOUTHERN MARGIN OF LGM GLACIATION IN NEW ZEALAND, *Geografiska Annaler: Series A, Physical Geography*, 91, 71-81, 2009.
- Bueler, E. and Brown, J.: Shallow shelf approximation as a “sliding law” in a thermomechanically coupled ice sheet model, *Journal of Geophysical Research: Earth Surface* (2003–2012), 114, 2009.
- 20 Buylaert, J.-P., Murray, A. S., Thomsen, K. J., and Jain, M.: Testing the potential of an elevated temperature IRSL signal from K-feldspar, *Radiation Measurements*, 44, 560-565, 2009.
- Campbell, I. B.: Soil pattern on Campbell Island, *New Zealand Journal of Science*, 24, 111-135, 1981.
- David, B., Roberts, R. G., Magee, J., Mialanes, J., Turney, C., Bird, M., White, C., Fifield, L. K., and Tibby, J.: Sediment mixing at Nonda Rock: investigations of stratigraphic integrity at an early archaeological site in northern Australia and implications for the human colonisation of the continent, *Journal of Quaternary Science*, 22, 449-479, 2007.
- 25 Denton, G. H., Anderson, R. F., Toggweiler, J. R., Edwards, R. L., Schaefer, J. M., and Putnam, A. E.: The Last Glacial Termination, *Science*, 328, 1652-1656, 2010.
- Duller, G.: Improving the accuracy and precision of equivalent doses determined using the optically stimulated luminescence signal from single grains of quartz, *Radiation Measurements*, 47, 770-777, 2012.
- 30 Duller, G. A.: Single- grain optical dating of Quaternary sediments: why aliquot size matters in luminescence dating, *Boreas*, 37, 589-612, 2008.
- EPICA, C. M.: One-to-one coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444, 195-198, 2006.
- Feathers, J. K.: Single-grain OSL dating of sediments from the Southern High Plains, USA, *Quaternary Science Reviews*, 22, 1035-1042, 2003.
- 35 Feathers, J. K., Holliday, V. T., and Meltzer, D. J.: Optically stimulated luminescence dating of Southern High Plains archaeological sites, *Journal of Archaeological Science*, 33, 1651-1665, 2006.
- Fleming, C. A., Mildenhall, D. C., and Moar, N. T.: QUATERNARY SEDIMENTS AND PLANT MICROFOSSILS FROM ENDERBY ISLAND, AUCKLAND ISLANDS, *Journal of the Royal Society of New Zealand*, 6, 433-&, 1976.
- 40



- Fogwill, C. J. and Kubik, P. W.: A glacial stage spanning the Antarctic Cold Reversal in Torres del Paine (51 degrees S), Chile, based on preliminary cosmogenic exposure ages, *Geografiska Annaler Series a-Physical Geography*, 87A, 403-408, 2005.
- 5 Fogwill, C. J., Turney, C. S. M., Hutchinson, D. K., Taschetto, A. S., and England, M. H.: Obliquity Control On Southern Hemisphere Climate During The Last Glacial, *Scientific Reports*, 5, 11673, 2015.
- Fraser, C. I., Nikula, R., Spencer, H. G., and Waters, J. M.: Kelp genes reveal effects of subantarctic sea ice during the Last Glacial Maximum, *Proceedings of the National Academy of Sciences*, 106, 3249-3253, 2009.
- 10 Galbraith, R. F., Roberts, R. G., Laslett, G. M., Yoshida, H., and Olley, J. M.: Optical dating of single and multiple grains of quartz from Jinmium rock shelter, northern Australia: Part I, experimental design and statistical models, *Archaeometry*, 41, 339-364, 1999.
- Gersonde, R., Crosta, X., Abelmann, A., and Armand, L.: Sea-surface temperature and sea ice distribution of the Southern Ocean at the EPILOG Last Glacial Maximum—a circum-Antarctic view based on siliceous microfossil records, *Quaternary Science Reviews*, 24, 869-896, 2005.
- 15 Golledge, N. and Levy, R.: Geometry and dynamics of an East Antarctic Ice Sheet outlet glacier, under past and present climates, *Journal of Geophysical Research: Earth Surface (2003–2012)*, 116, 2011.
- Golledge, N. R., Mackintosh, A. N., Anderson, B. M., Buckley, K. M., Doughty, A. M., Barrell, D. J. A., Denton, G. H., Vandergoes, M. J., Andersen, B. G., and Schaefer, J. M.: Last Glacial Maximum climate in New Zealand inferred from a modelled Southern Alps icefield, *Quaternary Science Reviews*, 46, 30-45, 2012.
- 20 Guo, Y.-J., Li, B., Zhang, J.-F., Yuan, B.-Y., Xie, F., and Roberts, R. G.: Luminescence ages for three ‘Middle Palaeolithic’ sites in the Nihewan Basin, northern China, and their archaeological and palaeoenvironmental implications, *Quaternary Research*, 85, 456-470, 2016.
- Guo, Y. J., Li, B., Zhang, J. F., Yuan, B. Y., Xie, F., and Roberts, R. G.: New ages for the Upper
- 25 Palaeolithic site of Xibaimaying in the Nihewan Basin, northern China: implications for small- tool and microblade industries in north- east Asia during Marine Isotope Stages 2 and 3, *Journal of Quaternary Science*, 32, 540-552, 2017.
- Hayward, B. W., Scott, G. H., Crundwell, M. P., Kennett, J. P., Carter, L., Neil, H. L., Sabaa, A. T., Wilson, K., Rodger, J. S., Schaefer, G., Grenfell, H. R., and Li, Q.: The effect of submerged plateaux on Pleistocene gyral circulation and sea-surface temperatures in the Southwest Pacific, *Global and Planetary Change*, 63, 309-316, 2008.
- 30 Hein, A. S., Hulton, N. R. J., Dunai, T. J., Sugden, D. E., Kaplan, M. R., and Xu, S.: The chronology of the Last Glacial Maximum and deglacial events in central Argentine Patagonia, *Quaternary Science Reviews*, 29, 1212-1227, 2010.
- 35 Heiri, O., Lotter, A., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results, *Journal of Paleolimnology*, 25, 101-110, 2001.
- Hesse, P. P.: The record of continental dust from Australia in Tasman Sea sediments, *Quat. Sci. Rev.*, 13, 257-272, 1994.
- 40 Hince, B.: Charles Fleming's cape expedition diary, Auckland Islands, 1942-43. CSIRO PUBLISHING 150 OXFORD ST, PO BOX 1139, COLLINGWOOD, VICTORIA 3066, AUSTRALIA, 2007.



- 5 Hodgson, D. A., Graham, A. G. C., Roberts, S. J., Bentley, M. J., Cofaigh, C. O., Verleyen, E., Vyverman, W., Jomelli, V., Favier, V., Brunstein, D., Verfaillie, D., Colhoun, E. A., Saunders, K. M., Selkirk, P. M., Mackintosh, A., Hedding, D. W., Nel, W., Hall, K., McGlone, M. S., Van der Putten, N., Dickens, W. A., and Smith, J. A.: Terrestrial and submarine evidence for the extent and timing of the Last Glacial Maximum and the onset of deglaciation on the maritime-Antarctic and sub-Antarctic islands, *Quaternary Science Reviews*, 100, 137-158, 2014a.
- 10 Hodgson, D. A., Graham, A. G. C., Roberts, S. J., Bentley, M. J., Cofaigh, C. Ó., Verleyen, E., Vyverman, W., Jomelli, V., Favier, V., Brunstein, D., Verfaillie, D., Colhoun, E. A., Saunders, K. M., Selkirk, P. M., Mackintosh, A., Hedding, D. W., Nel, W., Hall, K., McGlone, M. S., Van der Putten, N., Dickens, W. A., and Smith, J. A.: Terrestrial and submarine evidence for the extent and timing of the Last Glacial Maximum and the onset of deglaciation on the maritime-Antarctic and sub-Antarctic islands, *Quaternary Science Reviews*, 100, 137-158, 2014b.
- 15 Hogg, A., Southon, J., Turney, C., Palmer, J., Ramsey, C. B., Fenwick, P., Boswijk, G., Friedrich, M., Helle, G., and Hughen, K.: Punctuated Shutdown of Atlantic Meridional Overturning Circulation during Greenland Stadial 1, *Scientific Reports*, 6, 25902, 2016.
- 20 Hogg, A., Turney, C., Palmer, J., Southon, J., Kromer, B., Bronk Ramsey, C., Boswijk, G., Fenwick, P., Noronha, A., Staff, R., Friedrich, M., Reynard, L., Guetter, D., Wacker, L., and Jones, R.: The New Zealand kauri (*Agathis australis*) research project: A radiocarbon dating intercomparison of Younger Dryas wood and implications for IntCal13, *Radiocarbon*, 55, 2035-2048, 2013.
- 25 Huntley, D. J. and Hancock, R.: The Rb contents of the K-feldspar grains being measured in optical dating, *Ancient TL*, 19, 43-46, 2001.
- Huntley, D. J. and Lamothe, M.: Ubiquity of anomalous fading in K-feldspars and the measurement and correction for it in optical dating, *Canadian Journal of Earth Sciences*, 38, 1093-1106, 2001.
- 30 Jaccard, S., Hayes, C. T., Martínez-García, A., Hodell, D., Anderson, R. F., Sigman, D., and Haug, G.: Two modes of change in Southern Ocean productivity over the past million years, *Science*, 339, 1419-1423, 2013.
- Jacobs, Z., Duller, G. A., and Wintle, A. G.: Interpretation of single grain De distributions and calculation of De, *Radiation Measurements*, 41, 264-277, 2006.
- 35 Jacobs, Z., Meyer, M. C., Roberts, R., Aldeias, V., Dibble, H., and El Hajraoui, M.: Single-grain OSL dating at La Grotte des Contrebandiers ('Smugglers' Cave'), Morocco: improved age constraints for the Middle Paleolithic levels, *Journal of Archaeological Science*, 38, 3631-3643, 2011.
- Jacobs, Z. and Roberts, R. G.: Advances in optically stimulated luminescence dating of individual grains of quartz from archeological deposits, *Evolutionary Anthropology: Issues, News, and Reviews*, 16, 210-223, 2007.
- 40 Jacobs, Z. and Roberts, R. G.: An improved single grain OSL chronology for the sedimentary deposits from Diepkloof Rockshelter, Western Cape, South Africa, *Journal of Archaeological Science*, 63, 175-192, 2015.
- Jacobs, Z., Wintle, A. G., Duller, G. A., Roberts, R. G., and Wadley, L.: New ages for the post-Howiesons Poort, late and final Middle stone age at Sibudu, South Africa, *Journal of Archaeological Science*, 35, 1790-1807, 2008.



- Jones, J. M., Gille, S. T., Goose, H., Abram, N. J., Canziani, P. O., Charman, D. J., Clem, K. R., Crosta, X., de Lavergne, C., and Eisenman, I.: Assessing recent trends in high-latitude Southern Hemisphere surface climate, *Nature Climate Change*, 6, 917-926, 2016.
- 5 Kamb, B. and Echelmeyer, K. A.: Stress-gradient coupling in glacier flow: I. Longitudinal averaging of the influence of ice thickness and surface slope, *Journal of Glaciology*, 32, 267-284, 1986.
- Kaplan, M. R., Douglass, D. C., Singer, B. S., Ackert, R. P., and Caffee, M. W.: Cosmogenic nuclide chronology of pre-last glacial maximum moraines at Lago Buenos Aires, 46°S, Argentina, *Quaternary Research*, 63, 301-315, 2005.
- 10 Kaplan, M. R., Fogwill, C. J., Sugden, D. E., Hulton, N., Kubik, P. W., and Freeman, S.: Southern Patagonian glacial chronology for the Last Glacial period and implications for Southern Ocean climate, *Quaternary Science Reviews*, 27, 284-294, 2008.
- Kohfeld, K., Graham, R., De Boer, A., Sime, L., Wolff, E., Le Quéré, C., and Bopp, L.: Southern Hemisphere westerly wind changes during the Last Glacial Maximum: paleo-data synthesis, *Quaternary Science Reviews*, 68, 76-95, 2013.
- 15 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astronomy & Astrophysics*, 428, 261-285, 2004.
- Li, B., Jacobs, Z., and Roberts, R. G.: Investigation of the applicability of standardised growth curves for OSL dating of quartz from Haaq Fteah cave, Libya, *Quaternary Geochronology*, 35, 1-15, 2016.
- 20 Li, B., Jacobs, Z., Roberts, R. G., Galbraith, R., and Peng, J.: Variability in quartz OSL signals caused by measurement uncertainties: Problems and solutions, *Quaternary Geochronology*, 41, 11-25, 2017.
- Li, B., Jacobs, Z., Roberts, R. G., and Li, S.-H.: Review and assessment of the potential of post-IR IRSL dating methods to circumvent the problem of anomalous fading in feldspar luminescence, *Geochronometria*, 41, 178-201, 2014.
- 25 Li, B. and Li, S.: A reply to the comments by Thomsen et al. on "Luminescence dating of K-feldspar from sediments: a protocol without anomalous fading correction", 2012. 2012.
- Li, B. and Li, S.-H.: Luminescence dating of K-feldspar from sediments: a protocol without anomalous fading correction, *Quaternary Geochronology*, 6, 468-479, 2011.
- Li, W., Wang, R., Xiang, F., Ding, X., and Zhao, M.: Sea surface temperature and subtropical front movement in the South Tasman Sea during the last 800 ka, *Chinese Science Bulletin*, 55, 3338-3344, 30 2010.
- Lowe, J. and Walker, M.: Radiocarbon dating the last glacial-interglacial transition (Ca. 14-9 14C ka BP) in terrestrial and marine records: the need for new quality assurance protocols, *Radiocarbon*, 42, 53-68, 2000.
- MacAyeal, D. R.: EISMINT: Lessons in ice-sheet modeling, Department of Geophysical Sciences, 35 University of Chicago, Chicago, IL, 1832, 1839, 1997.
- Marshall, J. and Speer, K.: Closure of the meridional overturning circulation through Southern Ocean upwelling, *Nature Geoscience*, 5, 171-180, 2012.
- McGlone, M., Wilmschurst, J., and Meurk, C.: Climate, fire, farming and the recent vegetation history of subantarctic Campbell Island, *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 98, 71-84, 2007. 40
- McGlone, M. S.: The Late Quaternary peat, vegetation and climate history of the Southern Oceanic Islands of New Zealand, *Quaternary Science Reviews*, 21, 683-707, 2002.



- McGlone, M. S., Moar, N. T., Wardle, P., and Meurk, C. D.: Late-glacial and Holocene vegetation and environment of Campbell Island, far southern New Zealand, *Holocene*, 7, 1-12, 1997.
- McGlone, M. S., Turney, C. S. M., Wilmshurst, J. M., Renwick, J., and Pahnke, K.: Divergent trends in land and ocean temperature in the Southern Ocean over the past 18,000 years, *Nature Geosci*, 3, 622-626, 2010.
- 5 McGlone, M. S., Wilmshurst, J. M., and Wisler, S. K.: Lateglacial and Holocene vegetation and climatic change on Auckland Island, subantarctic New Zealand, *The Holocene*, 10, 719-728, 2000.
- Mitchell, K. J., Wood, J. R., Scofield, R. P., Llamas, B., and Cooper, A.: Ancient mitochondrial genome reveals unsuspected taxonomic affinity of the extinct Chatham duck (*Pachyanas chathamica*) and resolves divergence times for New Zealand and sub-Antarctic brown teals, *Molecular Phylogenetics and Evolution*, 70, 420-428, 2014.
- 10 Moreno, P. I., Kaplan, M. R., Francois, J. P., Villa-Martinez, R., Moy, C. M., Stern, C. R., and Kubik, P. W.: Renewed glacial activity during the Antarctic cold reversal and persistence of cold conditions until 11.5 ka in southwestern Patagonia, *Geology*, 37, 375-378, 2009.
- 15 Nelson, C. S., Cooke, P. J., Hendy, C. H., and Cuthbertson, A. M.: Oceanographic and climatic changes over the past 160,000 years at Deep Sea Drilling Project Site 594 off southeastern New Zealand, southwest Pacific Ocean, *Paleoceanography*, 8, 435-458, 1993.
- Newnham, R., Lowe, D. J., Giles T.M., and B.V., A.: Vegetation and climate of Auckland, New Zealand, since ca 32 000 cal. yr ago: support for an extended LGM., *Journal of Quaternary Science*, 22, 517-534, 2007.
- 20 Newnham, R. M., Lowe, D. J., and Green, J. D.: PALYNOLOGY, VEGETATION AND CLIMATE OF THE WAIKATO LOWLANDS, NORTH ISLAND, NEW-ZEALAND, SINCE C 18,000 YEARS AGO, *Journal of the Royal Society of New Zealand*, 19, 127-150, 1989.
- Oliver, R. L., Finlay, H. J., and Fleming, C. A.: The geology of Campbell Island, Department of Scientific and Industrial Research, Wellington, New Zealand Cape Expedition Series, Bulletin No. 3., 1950.
- 25 Otto-Bliesner, B. L., Brady, E. C., Clauzet, G., Tomas, R., Levis, S., and Kothavala, Z.: Last glacial maximum and Holocene climate in CCSM3, *Journal of Climate*, 19, 2526-2544, 2006.
- Pahnke, K. and Sachs, J. P.: Sea surface temperatures of southern midlatitudes 0–160 kyr B.P, *Paleoceanography*, 21, n/a-n/a, 2006.
- 30 Pahnke, K., Zahn, R., Elderfield, H., and Schulz, M.: 340,000-year centennial-scale marine record of Southern Hemisphere climatic oscillation, *Science*, 301, 948-952, 2003.
- Panitz, S., Cortese, G., Neil, H. L., and Diekmann, B.: A radiolarian-based palaeoclimate history of Core Y9 (Northeast of Campbell Plateau, New Zealand) for the last 160 kyr, *Marine Micropaleontology*, 116, 1-14, 2015.
- 35 Parrenin, F., Barnola, J. M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer, H., Fujita, S., Jouzel, J., Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V., Narcisi, B., Petit, J. R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R., Steffensen, J. P., Svensson, A., Udisti, R., Waelbroeck, C., and Wolff, E.: The EDC3 chronology for the EPICA Dome C ice core, *Clim. Past*, 3, 485-497, 2007.
- 40 Pedro, J. B., Bostock, H. C., Bitz, C. M., He, F., Vandergoes, M. J., Steig, E. J., Chase, B. M., Krause, C. E., Rasmussen, S. O., Markle, B. R., and Cortese, G.: The spatial extent and dynamics of the Antarctic Cold Reversal, *Nature Geosci*, advance online publication, 2015.



- Porter, S. C.: EQUILIBRIUM-LINE ALTITUDES OF LATE QUATERNARY GLACIERS IN SOUTHERN ALPS, NEW-ZEALAND, *Quaternary Research*, 5, 27-47, 1975.
- Prescott, J. R. and Hutton, J. T.: Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations, *Radiation measurements*, 23, 497-500, 1994.
- 5 Putnam, A. E., Schaefer, J. M., Barrell, D. J. A., Vandergoes, M., Denton, G. H., Kaplan, M. R., Finkel, R. C., Schwartz, R., Goehring, B. M., and Kelley, S. E.: In situ cosmogenic ^{10}Be production-rate calibration from the Southern Alps, New Zealand, *Quaternary Geochronology*, 5, 392-409, 2010.
- Quilty, P. G.: Origin and evolution of the sub-antarctic islands: the foundation, *Papers and Proceedings of the Royal Society of Tasmania*, 141, 35-58, 2007.
- 10 Ramsey, C. B.: *Dealing with Outliers and Offsets in Radiocarbon Dating*, 2009.
- Roberts, R., Bird, M., Olley, J., Galbraith, R., Lawson, E., Laslett, G., Yoshida, H., Jones, R., Fullagar, R., and Jacobsen, G.: Optical and radiocarbon dating at Jinmium rock shelter in northern Australia, *Nature*, 393, 358-362, 1998.
- Roberts, R., Galbraith, R., Olley, J., Yoshida, H., and Laslett, G.: Optical dating of single and multiple
15 grains of quartz from Jinmium rock shelter, northern Australia: part II, results and implications, *Archaeometry*, 41, 365-395, 1999.
- Roberts, R. G., Galbraith, R., Yoshida, H., Laslett, G., and Olley, J. M.: Distinguishing dose populations in sediment mixtures: a test of single-grain optical dating procedures using mixtures of laboratory-dosed quartz, *Radiation Measurements*, 32, 459-465, 2000.
- 20 Robinson, A., Calov, R., and Ganopolski, A.: An efficient regional energy-moisture balance model for simulation of the Greenland Ice Sheet response to climate change, *The Cryosphere*, 4, 129-144, 2010.
- Roche, D. M., Crosta, X., and Renssen, H.: Evaluating Southern Ocean sea-ice for the Last Glacial Maximum and pre-industrial climates: PMIP-2 models and data evidence, *Quaternary Science Reviews*, 56, 99-106, 2012.
- 25 Smedley, R., Duller, G., Pearce, N., and Roberts, H.: Determining the K-content of single-grains of feldspar for luminescence dating, *Radiation Measurements*, 47, 790-796, 2012.
- Steig, E. J., Schneider, D. P., Rutherford, S. D., Mann, M. E., Comiso, J. C., and Shindell, D. T.: Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year, *Nature*, 457, 459-462, 2009.
- 30 Summerhayes, C.: *The marine geology of the Auckland Islands area*, 1967.
- Thiel, C., Buylaert, J.-P., Murray, A., Terhorst, B., Hofer, I., Tsukamoto, S., and Frechen, M.: Luminescence dating of the Stratzing loess profile (Austria)—Testing the potential of an elevated temperature post-IR IRSL protocol, *Quaternary International*, 234, 23-31, 2011.
- 35 Thomsen, K. J., Murray, A. S., Buylaert, J.-P., Jain, M., Hansen, J., and Aubry, T.: Testing single-grain quartz OSL methods using sediment samples with independent age control from the Bordes-Fitte rockshelter (Roches d'Abilly site, Central France), *Quaternary Geochronology*, 31, 77-96, 2016.
- Thomsen, K. J., Murray, A. S., Jain, M., and Bøtter-Jensen, L.: Laboratory fading rates of various luminescence signals from feldspar-rich sediment extracts, *Radiation measurements*, 43, 1474-1486, 2008.
- 40 Turney, C. S., Jones, R. T., Lister, D., Jones, P., Williams, A. N., Hogg, A., Thomas, Z. A., Compo, G. P., Yin, X., and Fogwill, C. J.: Anomalous mid-twentieth century atmospheric circulation change over the South Atlantic compared to the last 6000 years, *Environ. Res. Lett.*, 11, 64009-64022, 2016a.



- Turney, C. S., Kershaw, A. P., Lowe, J. J., van der Kaars, S., Johnston, R., Rule, S., Moss, P., Radke, L., Tibby, J., and McGlone, M. S.: Climatic variability in the southwest Pacific during the Last Termination (20–10kyrBP), *Quaternary Science Reviews*, 25, 886-903, 2006.
- 5 Turney, C. S. M. and Jones, R. T.: Does the Agulhas Current amplify global temperatures during super-interglacials?, *Journal of Quaternary Science*, 25, 839-843, 2010.
- Turney, C. S. M., McGlone, M., Palmer, J., Fogwill, C., Hogg, A., Thomas, Z. A., Lipson, M., Wilmshurst, J. M., Fenwick, P., Jones, R. T., Hines, B., and Clark, G. F.: Intensification of Southern Hemisphere westerly winds 2000-1000 years ago: Evidence from the subantarctic Campbell and Auckland Islands (52-50°S), *Journal of Quaternary Science*, 31, 12-19, 2016b.
- 10 Van der Putten, N., Verbruggen, C., Ochyra, R., Verleyen, E., and Frenot, Y.: Subantarctic flowering plants: pre- glacial survivors or post- glacial immigrants?, *J. Biogeogr.*, 37, 582-592, 2010.
- WAIS, D. P. M.: Precise inter-polar phasing of abrupt climate change during the last ice age, *Nature*, 520, 661-665, 2015.
- Walker, M., Johnsen, S., Rasmussen, S. O., Popp, T., Steffensen, J.-P., Gibbard, P., Hoek, W., Lowe, J., Andrews, J., Björck, S., Cwynar, L. C., Hughen, K., Kershaw, P., Kromer, B., Litt, T., Lowe, D. J., Nakagawa, T., Newnham, R., and Schwander, J.: Formal definition and dating of the GSSP (Global Stratotype Section and Point) for the base of the Holocene using the Greenland NGRIP ice core, and selected auxiliary records, *Journal of Quaternary Science*, 24, 3-17, 2009.
- 15 Watson, E.: Two nivation cirques near Aberystwyth, Wales, *Biuletyn Peryglacjalny*, 15, 79-101, 1966.
- 20 Williams, P. W., McGlone, M., Neil, H., and Zhao, J.-X.: A review of New Zealand palaeoclimate from the Last Interglacial to the global Last Glacial Maximum, *Quaternary Science Reviews*, 110, 92-106, 2015.



Tables

#	Location	Site	Lat (S)	Long (E)	Elevation (m)	LTEL Lab code	Lab code	Depth (cm)	Material	d13C	Uncalibrated age (14C yrs BP)	+/-	Calibrated mean age (yrs BP)	+/-
1	Auckland	Sandy Bay 2	50°29'55.90"S	166°17'23.80"E	5	X13/82	Wk-38429 NZ	119	Peat	n.m	13020	36	15471	107
2	Auckland	North Point Bay Pillar Rock Exposure	50°31'0.09"S	166°10'29.60"E	40	X94/16	8226 WK-	45	Wood	-28.8	9292	98	10439	125
3	Auckland	Dea's Head ¹	50°30'58.29"S	166°12'59.07"E	20	X14/52	40801 NZA	119	Peat	n.m.	10086	31	11550	110
4	Auckland	Red Tape Bog	50°31'28.57"S	166°13'5.55"E	25	X94/12	4608 NZA	440	Silty peat	-27	11080	120	12910	108
5	Auckland	Near O Transect	50°31'29.85"S	166°12'59.73"E	20	X94/11	4607 WK	570	Peat	-27.31	11951	95	13744	129
6	Auckland	O Transect	50°31'37.42"S	166°12'42.60"E	20	X94/15	19756 WK	105	Bulk peat	-27.2	9057	52	10186	70
7	Auckland	Ranui Cove ²	50°31'36.58"S	166°12'42.08"E	20	X94/14	19734 WK-	340	Bulk peat	-28.2	12173	67	13988	108
8	Auckland	Mt Hooker	50°32'12.60"S	166°15'48.48"E	20	X13/10	6 38424 NZA	651	Peat	n.m.	9374	28	10543	56
9	Auckland	Musgrave Harbour	50°32'51.85"S	166°10'42.42"E	275	X94/17	9293 WK-	185	Peat	-27.9	10859	77	12732	61
10	Auckland	Carnley Harbour 2	50°47'13.65"S	165°58'18.71"E	52	X14/50	40795 WK-	175	Peat	n.m.	9222	30	10338	60
11	Auckland	Carnley Harbour 3	50°48'12.47"S	166° 0'43.79"E	n/a	n/a	39916 WK-	29-30	Peat	n.m.	10681	29	12617	41
12	Auckland	Tagua Bay Lower Lake	50°49'39.47"S	166° 0'41.74"E	n/a	n/a	39919 WK-	49.5-50	Peat	n.m	10143	30	11686	120
13	Auckland	Speight Corrie Lower Lake	50°48'54.36"S	166° 3'47.31"E	157	X13/87	38425 WK	288	Peat	n.m.	9717	30	11070	114
14	Auckland	Speight Terrace	50°49'34.90"S	165°59'14.06"E	189	X14/50	41406 WK	108	Silty peat	n.m.	10150	40	11699	135
15	Auckland	Hooker Cliffs ³	50°49'31.78"S	165°59'46.78"E	120	X13/85	39079 M84/1 NZ689	186	Peat	n.m.	12306	48	14200	122
16	Campbell	Col Ridge Crest	52°28'19.16"S	169°11'36.08"E	60	X99/9	1 8 WK	440	Peat	n.m.	12950	200	15290	291
17	Campbell	Homestead Scarp ⁴	52°32'20.23"S	169° 7'37.90"E	200	X99/11	14730 WK	157	Peat	-29.6	9416	57	10597	91
18	Campbell	Homestead Ridge	52°32'55.83"S	169° 8'7.56"E	30	X99/1	19746 AA	390	Bulk peat Wood fragment	-29.2	13648	73	15000	136
19	Campbell	Rocky Bay	52°33'2.45"S	169° 8'1.86"E	50	X99/1	44135 NZ694	945	s	-27.8	12780	120	15128	233
20	Campbell	Mt Honey Saddle ⁴	52°33'40.32"S	169° 5'2.17"E	287	n/a	8 WK	340	Peat	n.m	11700	140	13497	147
21	Campbell		52°34'35.48"S	169° 8'48.22"E	120	X99/3	13466	650	Peat	-27.9	12445	76	14509	215



Table 1: Radiocarbon dating of basal peat from sediment cores across Campbell and Auckland Islands used to model peat growth onset. Column 11: n.m. marks samples where $\delta^{13}\text{C}$ not measured owing to small sample size. Core numbers align with those on Fig. 1. Previously published dates shown by superscript numbers: ¹(McGlone et al., 2000); ²(McGlone, 2002); ³(McGlone et al., 1997); ⁴(McGlone et al., 2010)

5

Sediment sequence	Lab code	Depth (cm)	Material	Uncalibrated age (14C yrs BP)	+/-	Calibrated mean age (yrs BP)	+/-
LLS Corrie	Wk-41404	94	Peat	8359	36	9330	70
LLS Corrie	Wk-41405	100	Peat	8182	34	9090	60
LLS Corrie	Wk-41406	108	Mineral peat	10150	40	11699	135
LLS Corrie	Wk-41407	149	Silty peat	5630	26	n/a	n/a
LLS Corrie	Wk-41408	153.5	Silty peat	7449	40	n/a	n/a
LLS Terrace	Wk-39076	50	Peat	5445	25	6220	50
LLS Terrace	Wk-39077	99	Peat	7279	28	8060	50
LLS Terrace	Wk-39078	170	Peat	11952	36	13710	70
LLS Terrace	Wk-41149	175	Peat	12225	44	14070	80
LLS Terrace	Wk-39079	185	Peat	12306	48	14200	122
LLS Terrace	Wk-41150	196	Sandy peat	11729	36	n/a	n/a
LLS Terrace	Wk-38423	209	Sandy peat	11409	31	n/a	n/a
Pillar Rock	Wk-40800	94	Peat	8106	27	8960	80
Pillar Rock	Wk-40801	119	Peat	10086	31	11549	110
Pillar Rock	Wk-40799	238	Peat	36858	597	41350	500
Pillar Rock	Wk-40802	248	Peat	42234	203	45900	1250

Table 2: Radiocarbon dating for sedimentary sequences taken from Pillar Rock, Musgrave Harbour, and Lower ‘Lake’ Speight cirque and terrace.

10



Sample code	Grain size (μm)	Water content (%) ^a	External dose rate (Gy/ka) ^b			Internal beta dose rate (Gy/ka) ^b	Total dose rate (Gy/ka) ^b	D_e (Gy) ^{b, c}	pIRIR age (ka) ^{b, d}
			Gamma	Beta	Cosmic				
Enderby-1	180–212	25 ± 6 (17)	0.33 ± 0.02	0.59 ± 0.04	0.10 ± 0.01	0.67 ± 0.15	1.69 ± 0.16	678 ± 85 (600 / 9)	400 ± 64
Enderby-2	90–125	25 ± 6 (26)	0.25 ± 0.02	0.53 ± 0.04	0.11 ± 0.01	0.39 ± 0.09	1.28 ± 0.10	441 ± 42 (600 / 32)	343 ± 43
Enderby-3	90–125	25 ± 6 (23)	0.30 ± 0.02	0.62 ± 0.04	0.13 ± 0.02	0.39 ± 0.09	1.44 ± 0.10	408 ± 142 (300 / 4)	284 ± 101
Enderby-4	90–125	25 ± 6 (22)	0.30 ± 0.02	0.61 ± 0.04	0.14 ± 0.02	0.39 ± 0.09	1.45 ± 0.10	598 ± 35 (500 / 39)	413 ± 39

Table 3: Dose rates, equivalent dose (D_e) values and post-infrared IRSL (pIRIR) ages for the Enderby samples. a: The 5 measured water content of each sample is shown in parentheses below the assumed long-term value of 25 ± 6% used for age estimation. We consider the latter value as a realistic estimate of the mean water content (± standard error) over the full period of sample burial (see text). b: All uncertainties are expressed at 1 σ . c: The numbers in parentheses are the number of grains measured / number of grains accepted for D_e estimation. d: The age uncertainty includes a systematic error of 2% to allow for possible bias in the calibration of the laboratory beta source.

10





Figures





Figure 1: Map of Auckland and Campbell Islands. (A) Inset showing location relative to New Zealand mainland. Star shows location of ocean core DSDP 594 used in flowline modelling; (B) Auckland Islands. Black box marks location of inset C; (C) Inset showing northeast Auckland Islands; (D) Campbell Island. Model flowline marked along Perseverance Harbour, red rectangle highlights location of submerged moraines. Locations of basal peat radiocarbon (^{14}C) dates given by core numbers (see main text and Table 1). Key glacial cirques marked by black dots.



5

10

15



20

Figure 2: Digital elevation model of Lower Lake Speight Cirque, captured by Unmanned Aerial Vehicle, overlain on imagery from Google Earth. Moraines are highlighted by dashed white lines, with locations of Cores 14 and 15 given by arrows.




Figure 3: Schematic of exposed sediment section surveyed and sampled at Pillar Rock, Auckland Island. Radiocarbon dates of organic sections given as uncalibrated ^{14}C dates and calibrated mean ages. Not to scale.





Figure 4: Schematic of exposed 'Enderby Formation' on Enderby Island. IRSL locations shown by black circles. Particle size analysis of laminated silts blown up at right.

10

15



5

10

15

20

25

Figure 5: Multibeam surveys of Perseverance Harbour, Campbell Island (A) Multibeam imagery of submarine moraines at Boyack Point and Shoal Point, respectively the north and south terrestrial expressions of the moraines, with Terror Shoal long known as a navigational hazard (Ross, 1847); (B) Multibeam imagery along the length of Perseverance Harbour. No other submerged moraine features were found, including at De la Vire and Davis Points, which could be considered similar terrestrial expressions.




Figure 6: Schematic of sediment cores taken from Lower Lake Speight Cirque and Terrace. Radiocarbon dates given as calibrated ages. Loss-on-ignition (LOI; %) curve shown alongside cores.

5

10




Figure 7: Ensemble of glacier simulations, coloured according to the mean annual air temperature depression used to scale the EPICA Dome C or sea surface temperature timeseries forcing of (A) all 288 flowline model runs and (B) the 102 simulations that fit the radiocarbon constraint. Inset shows the frequency of MAAT depressions in this subset from 3-8°C.




Figure 8: (A) Model output showing best fit of glacial extent to palaeoclimatic and geomorphological data, with mean annual air temperature depression of 5°C, seasonality range of 11°C, temperature-coupled precipitation regime and forced using the EPICA Dome C temperature data. Marine Isotope Stage 4, Last Glacial Maximum and Antarctic Cold Reversal are labelled. (B) Model output with identical climatic parameters to (A), except seasonality which is 8°C.

20

25




Figure 9: Plot showing key Late Pleistocene events on the islands. Organic growth at ~43 ka and onset of peat growth at 15.7 ka are shown by dark grey bars against sea surface temperature reconstructions (Pahnke and Sachs, 2006) and EPICA Dome C δT (Parrenin et al., 2007). The ACR and New Zealand timing of the LGM are highlighted by pale grey bars. Temperature reconstructions from Campbell Island (McGlone et al., 2010) are shown in green (Mount Honey) and red (Homestead Scarp), highlighting minimal ACR temperature depression over the islands despite the strong SST and EPICA Dome C expressions.