

Rogation ceremonies: key to understand past drought variability in northeastern Spain since 1650

*Tejedor E^{1,2}, de Luis M^{1,2}, Barriendos M³, Cuadrat JM^{1,2}, Luterbacher J^{4,5}, Saz MA^{1,2}

¹Dept. of Geography and Regional Planning. University of Zaragoza. Zaragoza. (Spain).

²Environmental Sciences Institute of the University of Zaragoza. Zaragoza. (Spain).

³Department of History. University of Barcelona (Spain).

⁴Department of Geography, Climatology, Climate Dynamics and Climate Change, Justus Liebig University Giessen, Germany

⁵Centre for International Development and Environmental Research, Justus Liebig University Giessen, Germany

*Correspondence to: Miguel Ángel Saz; masaz@unizar.es

ABSTRACT

In the northeast of the Iberian Peninsula, drought recurrence, intensity, persistence and spatial variability have been mainly studied by using instrumental data covering the past ca. 60 years. Fewer studies have reconstructed drought occurrence and variability for the preinstrumental period using documentary evidence and natural proxies. In this study, we compiled a unique dataset of rogation ceremonies, religious acts to ask god for rain, from 13 cities in the northeast of Spain and investigated the annual drought variability from 1650 to 1899 AD. We converted the qualitative information into three regionally different coherent areas (Mediterranean, Ebro Valley and Mountain) with semiquantitative, annually resolved (December to August) drought indices according to the type of religious act. The Mediterranean Drought Index was compared with the instrumental series of Barcelona for the overlapping period (1787-1899) and we discovered a highly significant and stable correlation with the Standard Precipitation Drought Index of May with a 4 months lag ($r=-0.53$; $p<0.001$), asserting the validity of the regional Drought Indices derived from the historical documents as drought proxies. We found common periods with prolonged droughts (during the mid and late 18th century) and extreme drought years (1775, 1798, 1753, 1691 and 1817) associated with more blocking situations. A superposed epoch analysis (SEA) was performed to test the regional hydroclimatic responses after major tropical volcanic eruptions. The SEA shows a significant decrease in drought events one year after the volcanic events, which might be explained by the decrease in evapotranspiration due to decreases in surface temperatures and, consequently, the higher water availability that increases soil moisture. In addition, we discovered a common and significant drought response two years after the Tambora volcanic eruption in the three regional drought indices. Documented information on rogations thus contains important independent information to reconstruct extreme drought events for specific seasons in areas and periods for which instrumental information and other proxies are scarce.

1. Introduction

Water availability is one of the most critical factors for human activities, human wellbeing and the sustainability of natural ecosystems. Drought is an expression of a precipitation deficit, which is often longer than a season, a year or even a decade. Drought leads to water shortages associated with adverse impacts on natural systems and socioeconomic activities, such as reductions in streamflow, crop failures, forest decay or restrictions on urban and irrigation water supplies (Eslamian and Eslamian, 2017). Droughts represent a regular, recurrent process that occurs in almost all climate zones. In the Mediterranean region, the impacts of climate change on water resources are of significant concern (García-Ruiz et al., 2001). Spain is one of the European countries with a large risk of drought caused by high temporal and spatial variability in the distribution of the precipitation (Vicente-Serrano et al., 2014; Serrano-Notivol et al., 2017). Several recent Iberian droughts and their impacts on society and the environment have been documented in the scientific literature (e.g., Dominguez Castro et al., 2012; Trigo et al. 2013; Vicente-Serrano et al. 2014; Russo et al. 2015; Turco et al. 2017). For instance, during the period from 1990 to 1995, almost 12 million people suffered from water scarcity, the loss in agricultural production was an estimated 1 billion Euro, hydroelectric production dropped by 14.5 % and 63% of southern Spain was affected by fires (Dominguez Castro et al., 2012). One of the most recent droughts in Spain lasted from 2004 to 2005 (García-Herrera et al., 2007) and was associated with major socioeconomic impacts (hydroelectricity and cereal production decreased to 40% and 60%, respectively, of the average value).

In the Iberian Peninsula, natural archives including tree-ring chronologies, lake sediments and speleothems have been used to infer drought variability before the instrumental period (Esper et al., 2015; Tejedor et al., 2016, 2017c; Benito et al., 2003, 2008; Pauling et al. 2006; Brewer et al., 2008; Carro-Calvo et al., 2013, Abrantes et al., 2017, Andreu-Hayles et al., 2017). Nevertheless, most of the highly temporally resolved natural proxy-based reconstructions represent high-elevation conditions during specific periods of the year (mainly summer e.g., Tejedor et al., 2017c). Spain has a high amount of documentary-based data with a good degree of continuity and homogeneity for many areas, which allows the derivation of important paleo climate information at different timescales and for various territories. Garcia-Herrera et al. (2003) describe the main archives and discuss the techniques and strategies used to derive climate-relevant information from documentary records. Past drought and precipitation patterns have been inferred by exploring mainly rogation ceremonies and historical records from Catalonia (Martin-Vide and Barriendos 1995; Barriendos, 1997; Barriendos and Llasat, 2003; Trigo et al. 2009), Zaragoza (Vicente-Serrano and Cuadrat, 2007), Andalusia (Rodrigo et al., 1998; 2000), central Spain (Domínguez-Castro et al., 2008; 2012; 2014; 2016) and Portugal (Alcoforado et al. 2000). In northeastern Spain, the most important cities were located on the riversides of the Ebro Valley, which were surrounded by large cropland areas (Fig. 1). Bad wheat and barley harvests triggered socio-economic impacts, including the impoverishment or malnutrition of families, the severe alteration of the market economy, social and political conflicts, marginality, loss of population due to emigration and starvation and diseases and epidemics, such as those caused by pests

(Tejedor, 2017a). Recent studies have related precipitation/drought variability in regions of Spain to wheat yield variability (Ray et al., 2015; Esper et al. 2017). The extent of impacts caused by droughts depends on the socio-environmental vulnerability of an area. This is related to the nature and magnitude of the drought and the social structure of societies, such as agricultural-based societies including trades (Scandlyn et al., 2010; Esper et al. 2017). During the past few centuries, Spanish society has been strongly influenced by the Catholic Church. Parishioners firmly believe in the will of God and the church to provide them with better harvests. They asked God to stop or provoke rain through rogations, a process created by bishop Mamertus in AD 469 (Fierro, 1991). The key factor in evaluating rogation ceremonies for paleo climate research is determining the severity and duration of adverse climatic phenomena based on the type of liturgical act that was organized after the deliberation and decision-making of local city councils (Barriendos, 2005). Rogations are solemn petitions by believers to ask God specific requests (Barriendos 1996, 1997). *Pro pluviam* rogations were conducted to ask for precipitation during a drought, and they therefore provide an indication of drought episodes and clearly identify climatic anomalies and the duration and severity of the event (Martín-Vide & Barriendos, 1995; Barriendos, 2005). In contrast, *pro serenitate* rogations were requests for precipitation to end during periods of excessive or persistent precipitation, which caused crop failures and floods. In the Mediterranean basin, the loss of crops triggered important socio-economic consequences and was related to insufficient rainfall. Rogations were an institutional mechanism to address social stress in response to climatic anomalies or meteorological extremes (e.g. Barriendos, 2005). The municipal and ecclesiastical authorities involved in the rogation process guaranteed the reliability of the ceremony and maintained a continuous documentary record of all rogations. The duration and severity of natural phenomena that stressed society can be reflected by the different levels of liturgical ceremonies that were applied (e.g. Martín-Vide and Barriendos, 1995; Barriendos, 1997; 2005). Through these studies, we learned that the present heterogeneity of drought patterns in Spain also occurred in the past few centuries, in terms of the spatial differences, severity and duration of the events (Martín-Vide, 2001, Vicente-Serrano 2006b). However, a compilation of the main historical document datasets that have been compiled over the past several years is lacking, impeding the creation of a continuous record of drought recurrences and intensities in the northeast of the Iberian Peninsula.

Here we compiled 13 series of historical documentary information of the *pro pluviam* rogation data from the Ebro Valley and the Mediterranean Coast of Catalonia (Fig. 1) from 1438 to 1945 (Tab. 1). Regarding the location of the cities, they cover a wide range spanning from Barcelona, which is near the sea (9 m a.s.l.), and Teruel (915 m a.s.l.) (Fig 1). Although some periods have already been analyzed for certain cities (i.e., Zaragoza in 1600-1900 AD by Vicente-Serrano and Cuadrat, 2007; Zaragoza, Calahorra, Teruel, Vic, Cervera Girona, Barcelona, Tarragona and Tortosa in 1750-1850 AD by Dominguez-Castro et al., 2012; La Seu d'Urgell, Girona, Barcelona, Tarragona, Tortosa and Cervera in 1760-1800 AD by Barriendos and Llasat, 2003), this is the first systematic approach analyzing all existing information for northeastern Spain, including new

unpublished data for Huesca (1557-1860 AD) and Barbastro (1646-1925 AD) and examining the 13 sites jointly for a period of 250 years (1650-1899 AD). We analyzed droughts across the sites and identify extreme drought years and common periods in frequency and intensity. We also analyze statistical links between drought indices and major tropical volcanic events in order to determine the effects of strong eruptions on regional droughts.

2. Methods

2.1. Study area

The study area comprises the northeastern part of Spain, with an area of approximately 100,000 km², and includes three geological units, the Pyrenees in the north, the Iberian Range in the south, and the large depression of the Ebro Valley that separates them (Fig. 1). The Ebro Valley has an average altitude of 200 m a.s.l. The Ebro Valley climate can be characterized as a Mediterranean type climate, with warm summers, cold winters and increasing continental characteristics with distance from the coast. Some geographic aspects determine its climatic characteristics; for example, several mountainous chains isolate the valley from moist winds, preventing precipitation. Thus, in the central areas of the valley, annual precipitation is low, with small monthly variations and an annual precipitation in the central Ebro Valley of approximately 322 mm (AEMET, 2012). In both the Pyrenees and the Iberian Range, the main climatic characteristics are related to a transition from oceanic/continental to Mediterranean conditions in the East. In addition, a gradually higher aridity towards the east and the south is caused by the barrier effect of the most frequent humid air masses (Vicente-Serrano, 2005; López-Moreno & Vicente-Serrano, 2007). Areas above 2000 m a.s.l. receive approximately 2,000 mm of precipitation annually, increasing to 2,500 mm of precipitation in the highest peaks of the mountain range (García-Ruiz, et al., 2001). The annual precipitation in the Mediterranean coast is higher than that in the middle Ebro Valley and ranges from approximately 500 mm in Tortosa to 720 mm in Girona (Serrano-Notivol et al., 2017).

2.2. From historical documents to climate: Development of drought index for each location in NE Spain from 1650 to 1899 AD

Historical documents from 13 cities in the northeast of Spain were compiled into a novel dataset by using a consistent approach (Fig. 1, Tab. 1, Tab. S1). These historical documents are the rogation ceremonies reported in the 'Actas Capitulares' of the municipal archives or main cathedrals. The extension of the consulted documents (described in Table S1) ranges from 461 years of continuous data in Girona, to 120 years in Lleida, with an average of 311 years of data on each station. Rogations not only were religious acts but also were supported by the participation of several institutions; agricultural organizations and municipal and ecclesiastical authorities analyzed the situation and deliberated before deciding to hold a rogation ceremony (Vicente-Serrano and Cuadrat, 2007). Usually, the agricultural organizations would request rogations

when they observed a decrease in rainfall, which could result in weak crop development. Then, municipal authorities would recognize the setback and discuss the advisability of holding a rogation ceremony. Whether a rogation was celebrated or not was not arbitrary, since rogations had a price paid by public coffers. When the municipal authorities decided to hold a rogation, the order was communicated to the religious authorities, who placed the rogation on the calendar of religious celebrations and organized and announced the rogation. Previous studies have reported that winter precipitation is key for the final crop production in dry-farming areas of the Ebro Valley (wheat and barley; Austin et al., 1998a, 1998b; McAneney and Arrué, 1993; Vicente-Serrano and Cuadrat, 2007). In addition to winter rogations, most of the rogations were held during the vegetation growth period (March-May) and harvest period (June-August), since the socio-economic consequences when the harvest was poor were more evident during these periods. Thus, it is reasonable to consider those rogations in an index from December to August.

The qualitative information contained conveyed by the rogations was transformed into a semiquantitative continuous monthly series following the methodology of the Millennium Project (European Commission, IP 017008-Domínguez-Castro et al., 2012). Only *pro pluviam* rogations were included in this study. According to the intensity of the religious act, which were homogenously performed throughout the Catholic territories and triggered by droughts, we categorized the events in 4 levels from low to high intensity: 0, there is no evidence of any kind of ceremony; 1, a simple petition within the church was held; 2, intercessors were exposed within the church; and 3, a procession or pilgrimage took place in the public itineraries, the most extreme type of rogation (see Tab. 2). Although rogations have appeared in historical documents since the late 15th century and were reported up to the mid 20th century, we restricted the common period to 1650-1899 AD, since there are a substantial number of data gaps before and after this period, although some stations do not extent the full period. A continuous drought index (DI) was developed for each site by grouping the rogations at various levels. A simple approach, similar to that of Martín-Vide and Barriendos (1995) and Vicente-Serrano and Cuadrat (2007), was performed. The annual DI values were obtained by determining the weighted average of the number of level 1, 2 and 3 rogations recorded between December and August in each city. The weights of levels 1, 2 and 3 were 1, 2, and 3, respectively. Accordingly, the drought index for each city is a continuous semiquantitative value from 0, indicating the absence of drought, to a maximum of 3 (Figure 2A).

2.3. Clustering station drought to regional drought indices from 1650 to 1899 AD

To develop regional drought indices, we performed a cluster analysis (CA) that separates data into groups (clusters) with minimum variability within each cluster and maximum variability between clusters. We selected the period of common data 1650-1770 to perform the cluster analysis. The main benefit of performing a cluster analysis

(CA) is that it allows similar data to be grouped together, which helps in the identification of common patterns between data elements. To assess the uncertainty in hierarchical cluster analysis, the R package ‘pvclust’ (Suzuki and Shimodaira, 2006) was used. We used the Ward’s method in which the proximity between two clusters is the magnitude by which the summed squared in their joint cluster will be greater than the combined summed square in these two clusters $SS_{12} - (SS_1 + SS_2)$ (Ward, 1963; Everitt et al., 2001). Then, the root of the square difference between co-ordinates of pair of objects is computed with its Euclidian distance. Finally, for each cluster within the hierarchical clustering, quantities called *p-values* are calculated via multiscale bootstrap resampling (1000 times). Bootstrapping techniques does not require assumptions such as normality in original data (Efron, 1979) and thus represents a suitable approach applied to the semiquantitative characteristics of drought indices (DI) derived from historical documents. The *p-value* of a cluster is a value between 0 and 1, which indicates how strongly the cluster is supported by the data. The package ‘pvclust’ provides two types of *p-values*: AU (approximately unbiased *p-value*) and BP (bootstrap probability) *value*. AU *p-value* is computed by multiscale bootstrap resampling and is a better approximation of an unbiased *p-value* than the BP value computed by normal bootstrap resampling. The frequency of the sites falling into their original cluster is counted at different scales, and then the *p-values* are obtained by analyzing the frequency trends. Clusters with high AU values, such as those >0.95 , are strongly supported by the data (Suzuki and Shimodaira, 2006). Therefore, in this study, sites belonging to the same group were merged by means of an arithmetical average (Eq.1).

$$\text{Eq.1 Regional Drought Index } (\bar{x}) = (x_1 + x_2 + x_3 \dots)/n$$

where x_n represents each individual annual drought index, and n is the number of drought indices per cluster. Then, to evaluate the relationship of each site’s rogations, we performed a matrix correlation (Spearman) between the new groups derived from the cluster and each individual drought index for the period of 1650-1899.

2.4. Validation of the regional Drought indices against overlapping instrumental series.

To better understand the relationship with the derive drought indices and the instrumental series, we used the longest instrumental precipitation and temperature series covering the period 1786-2017 AD (Prohom et al., 2012; Prohom et al., 2015) for the city of Barcelona and thus overlap the rogation ceremony’s period from 1786 to 1899 AD. The instrumental series was homogenized and developed including data from cities nearby and along the Mediterranean coast (see Prohom et al., 2015 for details). We then calculated the Standardized Precipitation Index (SPI, McKee et a., 1993) and the Standardized Evapotranspiration and Precipitation Index (SPEI, Begueria et al., 2014) and calculated spearman correlation between DIMED and the SPI/SPEI at different time scales including a maximum lag of 12 months covering the period 1787-1899. To further explore the relationship between the drought indices inferred from historical documents and the instrumental drought indices through time, we performed 30 and 50 years moving correlations.

2.5. Detecting extreme drought years and periods in the northeast of Spain between 1650-1899 AD and links to large-scale volcanic forcing

To identify the extreme drought years, we selected those years above the 99th percentile of each regional drought index and mapped them in order to find common spatial patterns. In addition, the 11-year running mean performed for each drought index helped highlight drought periods within and among the drought indices. Finally, since rogation ceremonies are a response of the population to an extreme event, we performed a superposed epoch analysis (SEA; Panofsky and Brier, 1958) of the three years before and after the volcanic event, using the package 'dplR' (Bunn, 2008) to identify possible effects on the hydroclimatic cycle caused by volcanic eruptions. The largest (Sigl et al., 2015) volcanic eruptive episodes chosen for the analysis were 1815, 1783, 1809, 1695, 1836, 1832, 1884 and 1862. In addition, we performed the SEA only with the largest eruption of this period, the Tambora eruption in the year 1815.

3. Results

3.1. From historical documents to climate: Development of drought index for each location in NE Spain from 1650 to 1899 AD

Performing a weighted average of the monthly data (see methods), we converted the ordinal data into continuous semiquantitative index data. As a result, we developed an annual drought index (from the previous December to the current August) for each of the 13 locations that contains continuous values from 0 to 3 collected from information on the annual mean extreme droughts of each year. The EDCF (Fig.2A) confirmed that the new drought indices can be treated as a continue variable since the Drought Index can take almost infinite values in the range from 0 to 3. Then, to study drought across the region, we performed a cluster analysis including the annual drought indices of the 13 cities. These data were then used to study the hydrological responses after strong tropical eruptions.

3.2. Clustering station drought to regional drought indices from 1650 to 1899 AD

The cluster analysis (CA, see methods) using the DI of the 13 locations for the period of 1650-1899 AD revealed three significantly coherent areas, hereafter known as Mountain, Mediterranean and Ebro Valley (Fig. 3). The first cluster includes cities that are similar in altitude (Teruel, La Seu) and similar in latitude (Barbastro, Lleida, Huesca, Girona, see Fig. 1). The cities within the second and third clusters are near the Ebro River (Calahorra, Zaragoza and Tortosa) or have similar climatic conditions (Cervera, Vic, Barcelona, Tarragona). Clusters two and three suggest (Fig. 3) that the coherence of the grouping can be explained by the influence and proximity of the Mediterranean Sea (Tortosa, Cervera, Tarragona, Vic and Barcelona) and the influence of a more continental climate (Zaragoza and Calahorra). Accordingly, three regional drought indices were developed by combining the individual DIs of each group; DI Mountain (DIMOU), composed of Barbastro, Teruel, Lleida, La Seu, and Girona; DI Mediterranean (DIMED),

composed of Tortosa, Cervera, Tarragona, Vic and Barcelona, and DI Ebro Valley (DIEV), composed of Zaragoza and Calahorra. Resulting drought indices in regional DI series can also varies from 0 to 3 but showing a quite continuous distribution range (Figure 2B).

The Spearman correlation matrix for the period of 1650-1899 AD confirms the high and significant ($p<0.05$) correlations between each individual DI and its corresponding group, asserting the validity of the new DI groups (Fig. 4). The correlations among the cluster drought indices range from 0.76 (between DIEV and DIMED) to $r=0.38$ (between DIEV and DIMOU) and $r=0.42$ (between DIMED and DIMOU). In DIEV, both of the local DIs show similar correlations (Zaragoza, $r=0.73$; Calahorra, $r=0.75$). In the DIMED cluster, the high correlations among the members show a strong coherency. DIMOU is the most heterogeneous cluster, with correlations of $r=0.57$ for Barbastro and $r=0.33$ for La Seu. Although each individual DI within this group and within the DIMOU shows significant correlation, when individual DIs are compared between each other, some correlation values are not significant ($p<0.05$).

3.3. Validation of the regional Drought indices against overlapping instrumental series.

The maximum correlation ($r=-0.53$; $p<0.001$) between the Mediterranean Drought Index and the instrumental SPI over the full 113-year period (1787-1899 AD; Fig.5C) is found for the SPI of May with a lag of 4 months (SPI_{MAY_4} hereafter). Slightly lower, though still significant correlation, is obtained when using the SPEI of May with a lag of 4 months ($SPEI_{MAY_4}$) ($r=-0.50$; $p<0.001$, Fig.5D). The moving correlations between SPI_{MAY_4} and DIMED for 30 and 50 years (Fig.5A; Fig.5B) show high and stable correlation through the full period. The relationship with the $SPEI_{MAY_4}$ is also high and stable throughout the overlapping period, although lower than with SPI_{MAY_4} . The next step (iv) will address the selection of extreme drought years and periods within the 250 years from 1650-1899 AD using information from the cluster analysis.

3.4. Detecting extreme drought years and periods in the northeast of Spain between 1650-1899 AD and links to large-scale volcanic forcing

According to the cluster grouping, the three new spatially averaged drought indices (DIEV, DIMED and DIMOU) are presented in Fig. 6. Mountain DI (DIMOU) had the least number of drought events and a maximum DI of 1.6 in 1650 AD. The Ebro Valley DI (DIEV) had the highest number of droughts (inferred by the highest number of positive index values) followed by the third region (Mediterranean, DI DIMED). The 17th and 18th centuries exhibited a relatively high number of strong droughts (Fig. 6). A drought period, as indicated by the high positive index values over the duration of the DIs in all three series, occurred from 1740 to 1755 AD. The lowest DIs were found at the end of the 19th century; thus, this period experienced a reduced drought frequency. The 11-year running mean shows common periods with low DI values, such as 1706-1717, 1800-1811, 1835-1846 and 1881-1892, which we infer to be 'normal' or without droughts. On the other hand, 1678-1689, 1745-1756, 1770-1781, and 1814-1825 are periods with

continuously high DIs, indicating that significant droughts affected the crops during these periods and intense rogation performances were needed.

In the Ebro Valley, the most extreme years (Fig. 7) (according to the 99% percentile of the years 1650-1899) were 1775 (drought index value of 2.8), 1798 (2.7), 1691 (2.6), 1753 (2.5) and 1817 (2.5). Most of these extreme drought years can also be found in the Mediterranean DI 1753 (2.6), 1775 (2.5), 1737 (2.3), 1798 (2.2) and 1817 (2.2). For the DI Mountain, the extreme drought years occurred in the 17th century: 1650 (1.6), 1680 (1.5), 1701 (1.5) and 1685 (1.4). These extreme drought years are spatially displayed in Fig. 7. In the years 1775 and 1798, the Ebro Valley, Mediterranean and some mountain cites suffered from severe droughts. It is notable that the year 1650 in the Mountain area presented high values of DI, while the other locations had very low DI values (DIEV=0.4; DIMED=0.8).

We performed a superposed epoch analysis (SEA, see methods) to study the drought response over NE Iberia to major volcanic eruptions (Fig. 8a). The figure shows significant decreases ($p<0.05$) in the Ebro Valley and Mediterranean DI values during the year of and one year after volcanic events. We did not find a post-volcanic drought response in the Mountain area. No significant response was found for any of the DIs two or three years after the volcanic eruptions, including the major volcanic eruptions. However, two years after the Tambora eruption in April 1815, there was a significant ($p<0.05$) increase in the three drought indices (DIEV, DIMED and DIMOU) (Fig. 8b), in agreement with findings of Trigo et al. (2009).

4. Discussion

The exploration of historical documents from the main Cathedrals or the municipal city archives, the so called 'Actas Capitulares', yielded the different types and payments of the rogation ceremonies that were performed in drought stress situations. In fact, it is challenging to determine whether the decrease in the number of rogations at the beginning and at the end of the 19th century is due to the lack of droughts, the loss of documents, or a loss of religiosity within these periods. For instance, after the Napoleonic invasion (1808-1814) and the arrival of new liberal ideologies (Liberal Triennial 1820-1823), there was a change in the mentality of people in the big cities. These new liberal ideas were concentrated in the places where commerce and industry began to replace agriculturally based economies, leading to strikes and social demonstrations demanding better labor rights. New societies were less dependent on agriculture; hence, in dry spells, the fear of losing crops was less evident and fewer rogations were performed. In summary, the apparent low frequency of rogations in the 19th century could be explained by a combination of political instability and the loss of religiosity and historical documents. Further limitations of converting qualitative information into quantitative data refer to the fact that, for instance, a drought index of level 2 does not necessarily imply a drought twice as intense as a drought index of level 1. This is an inherent limitation when dealing with historical documents as a climate proxy, and different approaches have been applied in the scientific literature (Vicente-

Serrano and Cuadrat, 2007; Dominguez-Castro et al., 2008). In our paper, we follow the methodology proposed in the Millennium Project (European Commission, IP 017008) and demonstrated in Domínguez-Castro et al., (2012). To that extent, the ECDF helped understanding the nature of the historical documents when transformed into semiquantitative data, which confirm that they can be treated as a continuous variable.

Besides, the drought indices of different cities had similar characteristics, which allowed the grouping. Clustering is a descriptive technique (Soni, 2012), the solution is not unique, and the results strongly rely upon the analyst's choice of parameters and yet, we found three significant ($p < 0.05$) and consistent structures across the drought stations. The fact that the main cities were located along the Ebro River, which is surrounded by vast areas of river orchards and watered crops, could have delayed the occurrence of rogation ceremonies, since the food supply of the region enables better adaptation to droughts. This might also explain the similarities between DIEV and DIMED. In addition, the clusters might not only be collecting climatic information but also diverse agricultural practices or even species. For instance, Cervera and Lleida, sharing similar annual precipitation totals, belong to the Mediterranean and the Mountain Drought Indices respectively. Lleida is located in a valley with an artificial irrigation system since the Muslim period, which is fed by the river Segre (one of the largest tributaries to the Ebro river). The drought in the Pyrenees is connected with a shortage of water for the production of energy in the mills as well as to satisfy irrigated agriculture. However, the irrigation system itself allowed them to manage the resource and resist much longer. Therefore, only the most severe droughts, and even so in an attenuated form, are perceived in the city. Cervera, located in the mountains, in the so-called pre-littoral system and its foothills, has a different precipitation dynamic more sensitive to the arrival of humid air from the Mediterranean. Besides, Lleida had a robust irrigation system that Cervera did not have. The droughts in Cervera are therefore more "Mediterranean" like and thus it seems consistent its presence in the Mediterranean Drought Index.

The Mediterranean Drought Index is then compared with the longest existing instrumental series for the city of Barcelona for the 1787-1899 AD period. To the best of our knowledge this is the first time that rogation ceremonies in the Iberian Peninsula are calibrated with such a long instrumental period. The correlation is maximized in May, the key month for the development of the harvest. In addition, the accumulated of 4 months is confirming the importance of the end of winter and spring precipitation for the appropriate development of the crops. The high DIMED correlation ($r = -0.53$; $p < 0.001$) indicates not only that this cluster is indeed capturing the Mediterranean drought signal, but also that it can indeed be used as a semiquantitative proxy.

In fact, it opens a new line of research that the authors will continue exploring in future studies. We believe that these results highlight the validity of the Drought Indices to be consider as continuous variables. In addition, by performing this analysis we also confirm that the grouping made by the cluster analysis demonstrates spatial coherency among the historical documents.

418 Compared to other drought studies based on documentary sources, the
419 persistent drought phase affecting the Mediterranean and the Ebro Valley areas in the
420 second half of the 18th century is similar to that found in Vicente-Serrano and Cuadrat,
421 (2007) for Zaragoza. The results for the second half of the 18th century also agree with
422 the drought patterns previously described for Catalonia (Barriendos, 1997, 1998;
423 Martín-Vide and Barriendos, 1995). Common drought periods were also found in 1650-
424 1775 for Andalusia (Rodrigo et al., 1999, 2000) and in 1725-1800 for Zamora
425 (Domínguez-Castro et al., 2008). In general, based on documentary sources from
426 Mediterranean countries, the second half of the 18th century has the highest drought
427 persistency and intensity, which may be because there were more blocking situations in
428 this period (Luterbacher et al. 2002, Vicente-Serrano and Cuadrat, 2007). The period of
429 1740-1800 AD coincides with the so-called 'Maldá anomaly period'; a phase
430 characterized by strong climatic variability, including extreme drought and wet years
431 (Barriendos and Llasat, 2003). The 18th century is the most coherent period, including a
432 succession of dry periods (1740-1755), extreme years (1753, 1775 and 1798) and years
433 with very low DIs, which we interpret as normal years. Next, the period from 1814-1825
434 is noteworthy due to its prolonged drought. The causes of this extreme phase are still
435 unknown. However, Prohom et al. (2016) suggested these years experienced a
436 persistent situation of atmospheric blocking and high-pressure conditions.

437 In the Ebro Valley and the Mediterranean area, rogation ceremonies were
438 significantly less frequent in the year of and one year after volcanic eruptions. Such
439 patterns may be explained by the volcanic winter conditions, which are associated with
440 reductions in temperature over the Iberian Peninsula 1-3 years after the eruption
441 (Fischer et al., 2007; Raible et al., 2016). The lower temperature is experienced in spring
442 and summer after volcanic eruptions compared to spring and summer conditions of
443 nonvolcanic years. This might be related to a reduction in evapotranspiration, which
444 reduces the risk of droughts. This reinforces the significance of volcanic events in large-
445 scale climate changes. In addition, the lower temperatures may benefit the soil moisture
446 of croplands.

447 Furthermore, a significant increase in the intensity of the droughts was observed
448 two years after the eruptive Tambora event in the third cluster (Mountain, Fig. 3). The
449 normal conditions in the year of and the year after the Tambora eruption and the
450 increased drought intensity two years after the event are in agreement with recent
451 findings about hydroclimatic responses after volcanic eruptions (Fischer et al., 2007;
452 Wegmann et al., 2014; Rao et al., 2017; Gao and Gao 2017) though based on tree ring
453 data only. In addition, Gao and Gao, (2017) highlight the fact that high latitude eruptions
454 tend to cause drier conditions in western-central Europe two years after the eruptions.
455 Rao et al., (2017) suggested that the forced hydroclimatic response was linked to a
456 negative phase of the East Atlantic Pattern (EAP), which causes anomalous spring uplift
457 over the western Mediterranean. This pattern was also found in our drought index for
458 the Tambora eruption (1815 AD), but no significant pattern was found in the NE of Spain
459 for the other major (according to Sigl et al., 2015) volcanic eruptions. In particular, the
460 mountain areas show less vulnerability to drought compared to the other regions. This

is mainly due to the fact, that mountainous regions experience less evapotranspiration, more snow accumulation and convective conditions that lead to a higher frequency of thunderstorms during the summertime. In addition, the productive system of the mountain areas is not only based on agriculture but also on animal husbandry, giving them an additional source for living in case of extreme drought. This might explain the lower coherence among stations within the DIMOU.

5. Conclusions

We developed a new dataset of historical documents by compiling historical records (rogation ceremonies) from 13 cities in the northeast of the Iberian Peninsula. These records were transformed into semiquantitative continuous data to develop drought indices (DIs). We regionalized them by creating three DIs (Ebro Valle, Mediterranean and Mountain), which cover the period from 1650 to 1899 AD. The intensity of the DI is given by the strength and magnitude of the rogation ceremony, and the spatial extent of the DI is given by the cities where the rogations were held.

Our study highlights three considerations: i) the spatial and temporal resolution of rogations should be taken into account, particularly when studying specific years, since the use of *pro-pluviam* rogations gives information about drought periods and not about rainfall in general. Accordingly, it must be stressed that the drought indices developed here are not precipitation reconstructions; rather, they are high-resolution extreme event reconstructions of droughts spells. The comparison of these results with other continuous proxy records must be carried out with caution (Dominguez-Castro et al., 2008), although here we found a very high and stable correlation with the instrumental series for the overlapping period, which opens new lines of research. ii) The validity of rogation ceremonies as a high-resolution climatic proxy to understand past drought variability in the coastal and lowland regions of the northeastern Mediterranean Iberian Peninsula is clearly supported by our study. This is crucial, considering that most of the high-resolution climatic reconstruction for the northern Iberian Peninsula have been developed using tree-ring records collected from high-elevation sites (>1,600 m a.s.l.) in the Pyrenees (Büntgen et al., 2008, 2017; Dorado-Liñán et al., 2012) and the Iberian Range (Esper et al., 2015, Tejedor et al., 2016, 2017a, 2017b, 2017c), thus inferring the climate of mountainous areas. iii) Particularly in the Mediterranean and in the Ebro Valley areas, imprints of volcanic eruptions are significantly detected in the drought indices derived from the rogation ceremonies. These results suggest that DI is a good proxy to identify years with extreme climate conditions in the past at low elevation sites.

In addition, recent studies have emphasized the great precipitation (González-Hidalgo, et al., 2011; Serrano-Notivol et al., 2017) and temperature variabilities (González-Hidalgo, et al., 2015) within reduced spaces, including those with a large altitudinal gradient, such as our study area. In addition, the rogations' historical data covers a gap within the instrumental measurement record of Spain (i.e., which starts in the 20th century). Hence, rogation data are key to understanding the full range of past climate characteristics (in lowlands and coastal areas) to accurately contextualize the

current climate change. We encourage the use of further studies to better understand past droughts and their influence on societies and ecosystems; learning from the past can help adaptation in the future, especially because climate variability is predicted to increase in the same regions where climate variability historically explained most of the variability in crop yield.

Acknowledgments

Supported by the project 'CGL2015-69985' and the government of Aragon (group Clima, Cambio Global y Sistemas Naturales, BOA 147 of 18-12-2002) and FEDER funds. We would like to thank the support of all the custodians of the historical documents.

Author Contributions statement

E.T., and J.M.C. conceived the study. J.M.C. and M.B. provided the data. E.T. and M.d.L. conducted the data analysis, and E.T. wrote the paper with suggestions of all the authors. All authors discussed the results and implications and commented on the manuscript at all stages.

Competing interests statement

The authors declare no competing interests.

References

- Abrantes, F., Rodrigues, T., Rufino, M., Salgueiro, E., Oliveira, D., Gomes, S., Oliveira, P., Costa, A., Mil-Homens, M., Drago, T., and Naughton, F.: The climate of the Common Era off the Iberian Peninsula, *Clim. Past*, 13, 1901-1918, 2017.
- Alcoforado, M. J., Nunes, M. F., Garcia, J. C., and Taborda, J. P.: Temperature and precipitation reconstruction in southern Portugal during the late Maunder Minimum (AD 1675-1715), *The Holocene*, 10, 333-340, 2000.
- Andreu-Hayles, L., Ummenhofer, C.C., Barriendos, M., Schleser, G.H., Helle, G., Leuenberger, M., Gutierrez, E., Cook, E.R.: 400 years of summer hydroclimate from stable isotopes in Iberian trees, *Clim. Dyn.*, 49, 143, 2017.
- Austin, R. B., Cantero-Martínez, C., Arrúe, J. L., Playán, E., and Cano-Marcellán, P.: Yield-rainfall relationships in cereal cropping systems in the Ebro river valley of Spain, *Eur. J. Agro.*, 8, 239–248, 1998a.
- Austin, R. B., Playán, E., Gimeno, J.: Water storage in soils during the fallow: prediction of the effects of rainfall pattern and soil conditions in the Ebro valley of Spain, *Agric. Water Manag.*, 36, 213–231, 1998b.
- Barriendos, M. Climate and Culture in Spain. Religious Responses to Extreme Climatic Events in the Hispanic Kingdoms (16th-19th Centuries). In Behringer, W., Lehmann H. and C. Pfister (Eds.), *Cultural Consequences of the Little Ice Age* (pp. 379-414). Göttingen, Germany: Vandenhoeck & Ruprecht, 2005.

541 Barriendos, M.: El clima histórico de Catalunya (siglos XIV-XIX) Fuentes, métodos y
542 primeros resultados, *Revista de Geografía*, XXX-XXXI, 69-96, 1996-1997.

543 Barriendos, M., and Llasat, M.C.: The Case of the 'Maldá' Anomaly in the Western
544 Mediterranean Basin (AD 1760–1800): An Example of a Strong Climatic Variability, *Clim.*
545 *Change*, 61, 191-216, 2003.

546 Barriendos, M.: Climatic variations in the Iberian Peninsula during the late Maunder
547 minimum (AD 1675-1715): An analysis of data from rogation ceremonies, *The Holocene*,
548 7, 105-111, 1997.

549 Beguería, S., Vicente-Serrano, S.M., Fergus Reig, Borja Latorre.: Standardized
550 Precipitation Evapotranspiration Index (SPEI) revisited: parameter fitting,
551 evapotranspiration models, kernel weighting, tools, datasets and drought monitoring,
552 *Int. J. Climatol.*, 34: 3001-3023, 2014.

553 Benito, G., Diez-Herrero, A., Fernao, G., and de Villalta, M.: Magnitude and frequency of
554 flooding in the Tagus Basin (Central Spain) over the last millennium, *Clim. Change*, 58,
555 171-192, 2003.

556 Benito, G., Thorndycraft, V. R., Rico, M., Sanchez-Moya, Y., and Sopena, A.: Palaeoflood
557 and floodplain records from Spain: evidence for long-term climate variability and
558 environmental changes, *Geomorph.*, 101, 68–77, 2008.

559 Brewer, S., Alleaume, S., Guiot, J. and Nicault, A.: Historical droughts in Mediterranean
560 regions during the last 500 years: a data/model approach, *Clim. Past*, 3, 55–366, 2007.

561 Bunn, A. G.: A dendrochronology program library in R (dplR), *Dendrochronologia*, 26,
562 115–124, 2008.

563 Büntgen, U., Frank, D., Grudd, H., and Esper, J.: Long-term summer temperature
564 variations in the Pyrenees, *Clim. Dyn.*, 31, 615–631, 2008.

565 Büntgen, U., Krusic, P. J., Verstege, A., Sangüesa Barreda, G., Wagner, S., Camarero, J. J.,
566 Ljungqvist, F. C., Zorita, E., Oppenheimer, C., Konter, O., Tegel, W., Gärtner, H.,
567 Cherubini, P., Reinig, F., Esper, J.: New tree-ring evidence from the Pyrenees reveals
568 Western Mediterranean climate variability since medieval times, *J. Clim.*, 30, 5295–
569 5318, 2017.

570 Carro-Calvo, L., Salcedo-Sanz, S., and Luterbacher, J.: Neural Computation in
571 Paleoclimatology: General methodology and a case study, *Neurocomput.*, 113, 262-268,
572 2013.

573 Dominguez-Castro, F., and García-Herrera, R.: Documentary sources to investigate
574 multidecadal variability of droughts, *Geophys. Res. Lett.*, 42, 13-27, 2016.

575 Domínguez-Castro, F., Santisteban, J. I., Barriendos, M., and Mediavilla, R.:
576 Reconstruction of drought episodes for central Spain from rogation ceremonies
577 recorded at Toledo Cathedral from 1506 to 1900: A methodological approach, *Glob.*
578 *Planet. Change*, 63, 230–242, 2008.

- 580 Domínguez-Castro, F., Ribera, P., García-Herrera, R., Vaquero, J. M., Barriendos, M.,
581 Cuadrat, J. M., and Moreno, J. M.: Assessing extreme droughts in Spain during 1750–
582 1850 from rogation ceremonies, *Clim. Past*, 8, 705–722, 2012.
- 583 Domínguez-Castro, F., de Miguel, J. C., Vaquero, J. M., Gallego, M. C., and García-
584 Herrera, R.: Climatic potential of Islamic chronicles in Iberia: Extreme droughts (AD 711–
585 1010), *The Holocene*, 24, 370–374, 2014.
- 586 Dorado Liñán, I., Büntgen, U., González-Rouco, F., Zorita, E., Montávez, J. P., Gómez-
587 Navarro, J. J., Brunet, M., Heinrich, I., Helle, G., and Gutiérrez, E.: Estimating 750 years
588 of temperature variations and uncertainties in the Pyrenees by tree-ring reconstructions
589 and climate simulations, *Clim. Past*, 8, 919–933, 2012.
- 590 Efron, B: Bootstrap Methods: Another Look at the Jackknife, *Ann. Statist.*, 7, 1, 1–26,
591 1979.
- 592 Eslamian, S., and Eslamian, F. A. (eds): *Handbook of Drought and Water Scarcity.*
593 *Principle of Drought and Water Scarcity.* CRC Press, Taylor & Francis LTD, pp. 607–626,
594 2017.
- 595 Esper, J., Büntgen, U., Denzer, S., Krusic, P. J., Luterbacher, J., Schäfer, R., Schreg, R., and
596 Werner, J.: Environmental drivers of historical grain price variations in Europe, *Clim.*
597 *Res.*, 72, 39–52, 2017.
- 598 Esper, J., Großjean, J., Camarero, J. J., García-Cervigón, A. I., Olano, J. M., González-
599 Rouco, J.F., Domínguez-Castro, F., Büntgen, U.: Atlantic and Mediterranean synoptic
600 drivers of central Spanish juniper growth, *Theor. Appl. Climatol.* 121, 571–579, 2015.
- 601 Everitt, B. S., Landau, S. and Leese, M.; *Cluster Analysis*, Oxford University Press, Inc., 4th
602 Edition, New York; Arnold, London. ISBN 0-340-76119-9, 2001.
- 603 Fierro, A. *Histoire de la météorologie.* Denoël, Paris, 1991.
- 604 Fischer, E. M., Luterbacher, J., Zorita, E., Tett, S. F. B., Casty, C., and Wanner, H.:
605 European climate response to tropical volcanic eruptions over the last half millennium,
606 *Geophys. Res. Lett.*, 34, L05707, 2007.
- 607 Gao, Y., and Gao, C.: European hydroclimate response to volcanic eruptions over the
608 past nine centuries, *Int. J. Climatol.*, 37, 4146–4157, 2017.
- 609 García-Herrera, R., García, R. R., Prieto, M. R., Hernández, E., Gimeno, L., and Díaz, H. F.:
610 The use of Spanish historical archives to reconstruct climate variability, *Bull. Am.*
611 *Meteorol. Soc.*, 84, 1025–1035, 2003.
- 612 García-Herrera, R., Paredes, D., Trigo, R., Trigo, I. F., Hernández, E., Barriopedro, D., and
613 Mendes, M.A.: The Outstanding 2004/05 Drought in the Iberian Peninsula: Associated
614 Atmospheric Circulation, *J. Hydrometeorol.*, 8, 483–498, 2007.

615 García-Ruiz, J. M. (Ed).: Los recursos hídricos superficiales del Pirineo aragonés y su
616 evolución reciente. Logroño, Geofroma, 2001.

617 González-Hidalgo, J. C., Brunetti, M., and de Luis, M.: A new tool for monthly
618 precipitation analysis in Spain: MOPREDAS database (monthly precipitation trends
619 December 1945–November 2005), *Int. J. Climatol.*, 31, 715–731, 2011.

620 Gonzalez-Hidalgo, J. C., Peña-Angulo, D., Brunetti, M., and Cortesi, N.: MOTEDAS: a new
621 monthly temperature database for mainland Spain and the trend in temperature (1951–
622 2010), *Int. J. Climatol.*, 35, 4444–4463, 2015.

623 López-Moreno, J. I., and Vicente-Serrano, S. M.: Atmospheric circulation influence on
624 the interannual variability of snow pack in the Spanish Pyrenees during the second half
625 of the 20th century, *Hydrol. Res.*, 38, 33–44, 2007.

626 Luterbacher, J., Xoplaki, E., Dietrich, D., Rickli, R., Jacobeit, J., Beck, C., Gyalistrias, D.,
627 Schmutz, C., and Wanner, H.: Reconstruction of Sea Level Pressure fields over the
628 Eastern North Atlantic and Europe back to 1500, *Clim. Dyn.*, 18, 545–561, 2002.

629 Martín-Vide, J. and Barriendos, M.: The use of rogation ceremony records in climatic
630 reconstruction: a case study from Catalonia (Spain), *Clim. Change*, 30, 201–221, 1995.

631 Martín-Vide, J., and Fernández, D.: El índice NAO y la precipitación mensual en la España
632 peninsular, *Investigaciones Geográficas*, 26, 41–58, 2001.

633 McAneney, K. J., and Arrúe, J. L.: A wheat-fallow rotation in northeastern Spain: water
634 balance – yield considerations, *Agronomie*, 13, 481–490, 1993.

635 McKee, T.B., Doesken, N.J., Kliest, J.: The relationship of drought frequency and duration
636 to time scales. In: *Proceedings of the 8th Conference on Applied Climatology*, Anaheim,
637 CA, USA, 17–22. American Meteorological Society, Boston, MA, USA, pp 179–184, 1993.

638 Panofsky, H. A., and Brier, G. W.: Some applications of statistics to meteorology,
639 Pennsylvania: University Park, 1958.

640 Pauling, A., Luterbacher, J., Casty, C., and Wanner, H.: Five hundred years of gridded high-
641 resolution precipitation reconstructions over Europe and the connection to large-scale
642 circulation, *Clim. Dyn.*, 26, 387–405, 2006.

643 Prohom, M., Barriendos, M., Aguilar, E., Ripoll, R. : Recuperación y análisis de la serie de
644 temperatura diaria de Barcelona, 1780–2011. *Cambio Climático. Extremos e Impactos*,
645 Asociación Española de Climatología, Serie A, Vol. 8, 207–217, 2012

646 Prohom, M., Barriendos, M. and Sanchez-Lorenzo, A.: Reconstruction and
647 homogenization of the longest instrumental precipitation series in the Iberian Peninsula
648 (Barcelona, 1786–2014), *Int. J. Climatol.*, 36, 3072–3087, 2015.

649 Raible, C. C., Brönnimann, S., Auchmann, R., Brohan, P., ...and Wegman, M.: Tambora
650 1815 as a test case for high impact volcanic eruptions: Earth system effects, *WIREs Clim.*
651 *Change*, 7, 569–589, 2016.

652 Rao, M. P., Cook, B. I., Cook, E. R., D'Arrigo, R. D., Krusic, P. J., Anchukaitis, K. J., LeGrande,
653 A. N., Buckley, B. M., Davi, N. K., Leland, C., and Griffin, K. L.: European and
654 Mediterranean hydroclimate responses to tropical volcanic forcing over the last
655 millennium, *Geophys. Res. Lett.*, 44, 5104–5112, 2017.

656 Ray, D. K., Gerber, J. S., MacDonald, G. K., and West, P. C.: Climate variation explains a
657 third of global crop yield variability, *Nat. Commun.* 6, 5989, 2015.

658 Rodrigo, F. S., Esteban-Parra, M. J., Pozo-Vázquez, D., and Castro-Díez, Y.: A 500-year
659 precipitation record in southern Spain, *Int. J. Climatol.*, 19, 1233- 1253, 1999.

660 Rodrigo, F. S., Esteban-Parra, M. J., Pozo-Vázquez, D., and Castro-Díez, Y.: Rainfall
661 variability in southern Spain on decadal to centennial times scales, *Int. J. Climatol.*, 20,
662 721-732, 2000.

663 Russo, A., Gouveia, C., Trigo, R., Liberato, M.L.R., and DaCamara, C.C.: The influence of
664 circulation weather patterns at different spatial scales on drought variability in the
665 Iberian Peninsula, *Front. Environ. Sci.*, 3, 1, 2015.

666 Scandlyn, J., Simon, C. N., Thomas, D. S. K., Brett, J. Theoretical Framing of worldviews,
667 values, and structural dimensions of disasters. In Phillips BD, Thomas DSK, Fothergill A,
668 Blinn-Pike L, editors. *Social Vulnerability to disasters*. Cleveland: CRC Press Taylor &
669 Francis Group, p. 27-49 (2010).

670 Serrano-Notivol, R., Beguería, S., Saz, M. A., Longares, L. A., and de Luis, M.: SPREAD: a
671 high-resolution daily gridded precipitation dataset for Spain – an extreme events
672 frequency and intensity overview, *Earth Syst. Sci. Data*, 9, 721-738, 2017.

673 Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., ...
674 Woodruff, T. E.: Timing and climate forcing of volcanic eruptions for the past 2,500 years,
675 *Nature*, 523 (7562), 543–549, 2015.

676 Soni, T.: An overview on clustering methods, *IOSR J. Engineering*, 2, 719-725, 2012.

677 Suzuki, R. & Shimodaira, H. Pvcust: an R package for assessing the uncertainty in
678 hierarchical clustering. *Bioinformatics* 22, 1540-1542 (2006).

679 Tejedor, E. Climate variability in the northeast of Spain since the 17th century inferred
680 from instrumental and multiproxy records. PhD thesis. University of Zaragoza. Zaragoza
681 (Spain), 2017a.

682 Tejedor, E., de Luis, M., Cuadrat, J. M., Esper, J., and Saz, M. A.: Tree-ring-based drought
683 reconstruction in the Iberian Range (east of Spain) since 1694, *Int. J. Biometeorol.*, 60,
684 361–372, 2016.

685 Tejedor, E., Saz, M. A., Cuadrat, J. M., Esper, J., and de Luis, M.: Temperature variability
686 in the Iberian Range since 1602 inferred from tree-ring records, *Clim. Past*, 13, 93-105,
687 2017b.

- Tejedor, E., Saz, M. A., Cuadrat, J.M., Esper, J. and de Luis, M.: Summer drought reconstruction in Northeastern Spain inferred from a tree-ring latewood network since 1734, *Geophys. Res. Lett.*, 44, 8492-8500, 2017c.
- Trigo, R. M., Vaquero, J. M., Alcoforado, M. J., Barriendos, M., Taborda, J., García-Herrera, R., and Luterbacher, J.: Iberia in 1816, the year without summer, *Int. J. Climatol.*, 29, 99–115, 2009.
- Trigo, R.M., Añel, J.A., Barriopedro, D., García-Herrera, R., Gimeno, L., Nieto, R., Castillo, R., Allen, M.R., and Massey, N.: The record Winter drought of 2011-12 in the Iberian Peninsula, in *Explaining Extreme Events of 2012 from a Climate Perspective*, *Bull. Am. Meteorol. Soc.*, 94, S41-S45, 2013.
- Turco, M., von Hardenberg, J., AghKouchak, A., Llasat, M. C., Provenzale, A., and Trigo, R.: On the key role of droughts in the dynamics of summer fires in Mediterranean Europe, *Nat. Sci. Rep.* 7, 81, 2017.
- Vicente-Serrano, S. M., and Cuadrat, J. M.: North Atlantic oscillation control of droughts in north-east Spain: Evaluation since 1600 A.D, *Clim. Change*, 85, 357-379, 2007.
- Vicente-Serrano, S. M., López-Moreno, J. I., Beguería, S., Lorenzo-Lacruz, J., Sánchez-Lorenzo, A., García-Ruiz, J. M., Azorín-Molina, E., Morán-Tejeda, E., Revuelto, J., Trigo, R., Coelho, F., and Espejo, F.: Evidence of increasing drought severity caused by temperature rise in southern Europe, *Environ Res Lett.*, 9 (4), 44001, 2014.
- Vicente-Serrano, S. M.: Las sequías climáticas en el valle medio del Ebro: Factores atmosféricos, evolución temporal y variabilidad espacial, *Consejo de Protección de la Naturaleza de Aragón, Zaragoza*, 277 pp, 2005.
- Vicente-Serrano, S. M.: Spatial and temporal analysis of droughts in the Iberian Peninsula (1910–2000), *Hydrol. Sci. J.*, 51, 83–97, 2006.
- Ward, J. H.: Hierarchical Grouping to Optimize an Objective Function, *Journal of the American Statistical Association*, 58, 236–244, 1963.
- Wegmann, M., and Brönnimann, S.: Volcanic influence on European Summer Precipitation through monsoons: possible cause for “years without summer, *J. Clim.*, 27, 3683–3691, 2014.




Figure 1. Location of the historical documents in the northeast of Spain.

730

731

732

733

734

735

736

737

738

739

740

741

Site	Latitude (degrees)	Longitude (degrees)	Altitude (m.a.s.l.)	Start (Years AD)	End	Extension (years)
<i>Zaragoza</i>	41.64	-0.89	220	1589	1945	356
<i>Teruel</i>	40.34	-1.1	915	1609	1925	316
<i>Barbastro</i>	42.03	0.12	328	1646	1925	279
<i>Calahorra</i>	42.3	-1.96	350	1624	1900	276
<i>Huesca</i>	42.13	-0.4	457	1557	1860	303
<i>Girona</i>	42.04	2.93	76	1438	1899	461
<i>Barcelona</i>	41.38	2.17	9	1521	1899	378
<i>Tarragona</i>	41.11	1.24	31	1650	1874	224
<i>Tortosa</i>	40.81	0.52	14	1565	1899	334
<i>La Seu</i>	42.35	1.45	695	1539	1850	311
<i>Vic</i>	41.92	2.25	487	1570	1899	329
<i>Cervera</i>	41.67	1.27	548	1484	1850	366
<i>Lleida</i>	41.61	0.62	178	1650	1770	120

742

Table 1. Historical document characteristics in the northeast of Spain.




Figure 2. The empirical cumulative distribution function (ECDF), used to describe a sample of observations of a given variable. Its value at a given point is equal to the proportion of observations from the sample that are less than or equal to that point. ECDF performed for the local drought indices (A) and the regional drought indices (B).




Figure 3. Dendrogram showing the hierarchical cluster analysis of the drought indices developed from the historical documents for each location. The AU (approximately unbiased p -value) is indicated in red and the BP (bootstrap probability) is presented in green.




Figure 4. Correlation matrix (Spearman) between the individual drought indices and the cluster drought indices for the period of 1650-1899. Values are significant at $p < 0.05$, except those marked with a gray cross, which are not significant.




Figure 5. A) 30y moving correlation between DIMED and the instrumental computed SPI and SPEI. B) Same but 50y moving correlations. C) Correlation (Spearman) between DIMED and SPI_{MAY_4} for the full period (1787-1899). D) Correlation (Spearman) between DIMED and SPEI_{MAY_4} for the full period (1787-1899).




Figure 6. Drought indices of the three clusters, DIMOU (Mountain), DIEV (Ebro Valley) and DIMED (Mediterranean). Vertical orange bars represent the drought index magnitude, 0 denotes normal conditions, and 3 denotes an extreme drought year. The extreme drought index years are also highlighted with a red circle. Extreme volcanic events from Sigl et al., 2015, are shown in the top panel. Vertical pink shadows indicate extreme common (for all three clusters) drought periods, while blue shadows indicate common periods with fewer droughts.




Figure 7. Spatial distribution of the most extreme drought years (based on the 99th percentile of the cluster drought indices). The distribution is ordered top-down. The drought index value (magnitude) for each site within the cluster is also represented.




Figure 8. a) Superposed epoch analysis (SEA) of the three regional drought indices, DIMOU (Mountain), DIEV (Ebro Valley) and DIMED (Mediterranean), with major volcanic events from Sigl et al., 2015. Black shadows show significance at $p < 0.05$, i.e., significantly lower or higher drought index values after the volcanic event. b) SEA of only the Tambora (1815) event showing a significant ($p < 0.05$) increase in the drought index.

Level	Type of ceremony
0	No ceremonies
1	Petition within the church
2	Masses and processions with the intercessor within the church
3	Pilgrimage to the intercessor of other sanctuary or church

Table 2. Rogation levels according to the type of ceremony celebrated.